

Regularity for the CR vector bundle problem II

XIANGHONG GONG AND SIDNEY M. WEBSTER

Abstract. We derive a $C^{k+\frac{1}{2}}$ Hölder estimate for $P\varphi$, where P is either of the two solution operators in Henkin's local homotopy formula for $\bar{\partial}_b$ on a strongly pseudoconvex real hypersurface M in \mathbb{C}^n , φ is a $(0, q)$ -form of class C^k on M , and $k \geq 0$ is an integer. We also derive a C^a estimate for $P\varphi$, when φ is of class C^a and $a \geq 0$ is a real number. These estimates require that M be of class $C^{k+\frac{5}{2}}$, or C^{a+2} , respectively. The explicit bounds for the constants occurring in these estimates also considerably improve previously known such results.

These estimates are then applied to the integrability problem for CR vector bundles to gain improved regularity. They also constitute a major ingredient in a forthcoming work of the authors on the local CR embedding problem.

Mathematics Subject Classification (2010): 32V05 (primary); 32A26, 32T15 (secondary).

1. Introduction

In this paper we will prove the following.

Theorem 1.1. *Let $n \geq 4$. Let M be a strongly pseudoconvex real hypersurface in \mathbb{C}^n of class C^2 . Let ω be an $r \times r$ matrix of continuous $(0, 1)$ -forms on M . Assume that $\bar{\partial}_M \omega = \omega \wedge \omega$. Near each point of M , there exists a non-singular matrix A of Hölder class $C^{1/2}$ satisfying $\bar{\partial}_M A = -A\omega$ and*

- a) $A \in C^a(M)$, if $\omega \in C^a(M)$, $M \in C^{a+2}$ and $a > 0$ is a real number;
- b) $A \in C^{k+\frac{1}{2}}(M)$, if $\omega \in C^k(M)$, $M \in C^{k+\frac{5}{2}}$ and $k > 0$ is an integer.

If ω and A are of class C^0 , the identities $\bar{\partial}_M \omega = \omega \wedge \omega$ and $\bar{\partial}_M A = -A\omega$ are in the sense of currents; see Section 3. This work is a continuation of [5]. For earlier results see [23] and Ma-Michel [15].

We now describe the above result in terms of an integrability problem for CR vector bundles ([23]). Let E be a complex vector bundle of rank r over M with

Research supported in part by NSF grant DMS-0705426.

Received October 19, 2009; accepted February 18, 2010.

a connection D . For a local frame $e = (e_1, \dots, e_r)$, $De_i = \omega_i^j e_j$, where $\omega = (\omega_i^j)$ are connection 1-forms on M ; by a frame change $\tilde{e} = Ae$, $D\tilde{e} = \tilde{\omega}\tilde{e}$ with $\tilde{\omega} = (dA + A\omega)A^{-1}$. The integrability problem is to find an A such that the new connection forms $\tilde{\omega} = (\tilde{\omega}_i^j)$ belong to the ideal $\mathcal{J}(M)$ generated by $(1, 0)$ -forms on M . The integrability condition is that the curvature 2-forms $d\omega - \omega \wedge \omega$ belong to $\mathcal{J}(M)$.

We want to mention a few ingredients in the proof. The integrability problem is local. As in [5, 20], we will use the Henkin local homotopy formula. Let $M \subset \mathbb{C}^n$ be a graph over a domain $D \subset \mathbb{R}^{2n-1}$, given by $y^n = |z'|^2 + \hat{r}(z', x^n)$ with $\hat{r}(0) = \partial\hat{r}(0) = 0$, where $z = (z', z^n)$ are standard coordinates. Set $M_\rho = M \cap \{(x^n)^2 + y^n < \rho^2\}$ and $D_\rho = \pi(M_\rho)$. Suppose that $\overline{D}_{\rho_0} \subset D$ and the \mathcal{C}^2 -norm $\|\hat{r}\|_{\rho_0, 2}$ of \hat{r} on D_{ρ_0} is sufficiently small. For $0 < \rho \leq \rho_0$ and $n \geq 4$, we have the Henkin homotopy formula

$$\varphi = \bar{\partial}_M P\varphi + Q\bar{\partial}_M\varphi \tag{1.1}$$

for $(0, q)$ -forms φ on \overline{M}_ρ with $0 < q < n - 2$. We will prove the following estimates.

(i) Let $a \geq 0$ be a real number. Then

$$\|P\varphi\|_{(1-\sigma)\rho, a} \leq C_a \rho^{-s_*} \sigma^{-s} (\|\varphi\|_{\rho, a} + \|\hat{r}\|_{\rho, a+2} \|\varphi\|_{\rho, 0}).$$

(ii) Let $k \geq 0$ be an integer. Then

$$\begin{aligned} \|P\varphi\|_{(1-\sigma)\rho, k+\frac{1}{2}} &\leq C_k \rho^{-s_*} \sigma^{-s} (1 + \|\hat{r}\|_{\rho, \frac{s}{2}}) \|\varphi\|_{\rho, k} + \|\hat{r}\|_{\rho, k+\frac{s}{2}} \|\varphi\|_{\rho, 0}; \\ \|P\varphi\|_{(1-\sigma)\rho, 1/2} &\leq C \rho^{-1} \sigma^{1-2n} \|\varphi\|_{\rho, 0}, \quad k = 0, \quad q = 1. \end{aligned}$$

(See (10.16) for s, s_* .) We emphasize that the estimates hold for all $0 < \rho \leq \rho_0 \leq 3$ and $0 < \sigma < 1$. Under the coordinates (z', x^n) of M , $\|\cdot\|_{\rho, a}$ denotes the standard \mathcal{C}^a -norm on the domain $D_\rho \subset \mathbb{R}^{2n-1}$. The same estimates hold for Q ; however, the second estimate in (ii), based on a special property of the kernels for $(0, 1)$ forms, is not applicable to Q when it operates on $(0, q + 1)$ form with $q > 0$. See Romero [17] for estimates in Hölder norms for the Heisenberg group case and an example showing necessity of blow-up constants.

The estimate (i) is proved in Proposition 10.1 and (ii) is in Proposition 11.1. The above theorem and two estimates are our main results. With the estimates, we will prove Theorem 1.1 by using a KAM rapid iteration argument as in [5], which avoids the Nash-Moser smoothing techniques.

We now describe some ideas to derive the estimates. The integral operators P, Q are estimated in the same way. Let us focus on $P = P_0 + P_1$, where P_1 is an integral operator over ∂M_ρ and P_0 is over M_ρ . Since we need estimates only on shrinking domains, the boundary integral P_1 can be treated easily. For the interior integral P_0 , via cutoff, the difficulties lie in the case where the $(0, q)$ -form

$\varphi = \sum \varphi_{\bar{J}} d\bar{z}^{\bar{J}}$ has compact support. We will see that coefficients of $(0, q - 1)$ -form $P_0\varphi$ are sums of $\mathcal{K}f(z) = \int_{M_\rho} f(\zeta)k(\zeta, z) dV(\zeta)$, where $f(\zeta) = \varphi_{\bar{J}}(\zeta)U_{\bar{J}}(\zeta)$ and $U_{\bar{J}}$ depend on derivatives of $|\zeta'|^2 + \hat{r}(\zeta', \xi^n)$ of order at most two, and

$$k(\zeta, z) = \frac{r_{\zeta^j} - r_{z^j}}{(r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b}, \quad a = n - q, \quad b = q$$

for $r(z) = -y^n + |z'|^2 + \hat{r}(z', x^n)$; see Section 3 for details.

To deal with the singularity of the kernel along $\zeta = z$, we fix $z \in M$ and apply the approximate Heisenberg transformation $\zeta \rightarrow \zeta_*$ defined by

$$\psi_z: \zeta'_* = \zeta' - z', \quad \xi^n_* = -2ir_z \cdot (\zeta - z).$$

We will show that $\zeta'_*, \xi^n_* = 2 \operatorname{Im}(r_z \cdot (\zeta - z))$ form coordinates of $\psi_z(M)$, and under this coordinate system the integral becomes

$$\mathcal{K}f(z) = \int_{\pi(\psi_z(M_\rho))} f(\psi_z^{-1}(\zeta_*))k_*(\zeta'_*, \xi^n_*, z', x^n) dV(\zeta'_*, \xi^n_*).$$

Here

$$k_*(\zeta'_*, \xi^n_*, z', x^n) = \sum_{|I|=1} E_I(\zeta'_*, \xi^n_*, z', x^n) \hat{k}_{ab}^I(\zeta'_*, \xi^n_*),$$

$$\hat{k}_{ab}^I(\zeta'_*, \xi^n_*) = \frac{(\zeta'_*, \bar{\zeta}'_*, \xi^n_*)^I}{(|\zeta'_*|^2 + i\xi^n_*)(|\zeta'_*|^2 - i\xi^n_*)^b}, \quad a + b = n,$$

where $I = (i_1, \dots, i_{2n-1})$ and we use standard multi-index notation. The coefficients $E_I(\zeta'_*, \xi^n_*, \cdot)$ are of class \mathcal{C}^a if $(\zeta'_*, \xi^n_*) (\neq 0)$ is fixed and $\hat{r} \in \mathcal{C}^{a+2}$. Since f has compact support, one can take derivatives of $\mathcal{K}f$ directly onto f and onto E_I without disturbing the kernels \hat{k}_{ab}^I . The transformation ψ_z has been used by other people. See Bruna-Burgués [1] and Ma-Michel [13]. To obtain estimate (ii), we need to return to the original coordinates after differentiation. This will give us another formula for the derivatives of $\mathcal{K}f$, which allows us to reduce the $\mathcal{C}^{k+\frac{1}{2}}$ -estimates to the Hölder $\frac{1}{2}$ -estimate for new kernels of the same type.

We would like to mention some methods to derive the fundamental $\frac{1}{2}$ -estimate. Kerzman [11] obtained Hölder α -estimates for all $\alpha < \frac{1}{2}$ for $\bar{\partial}$ -solutions, by estimating a Cauchy-Fantappiè form. Folland-Stein [3] used non-isotropic balls and piecewise smooth curves in complex tangential directions to obtain estimates in their spaces on the real hyperquadric. Henkin-Romanov [8] obtained the $\frac{1}{2}$ -estimate for $\bar{\partial}$ -equations on strongly pseudoconvex domains via a type of Hardy-Littlewood lemma (see also Henkin [7] for $\bar{\partial}_b$ on strictly convex boundaries). Our estimate, like the classical Hölder estimate for the Newtonian potential, is still based on a decomposition of domain. However, we delete a *cylinder* about the pole, instead of a (non-isotropic) ball. The radius of the cylinder is optimized for the $\frac{1}{2}$ -exponent

and is yet so large that, when estimating the Hölder $\frac{1}{2}$ -ratio at two points, we can ignore their non-isotropic distance and connect them with a line segment.

In the 5 dimensional case there is an extra term added to the right-hand side of (1.1). As of this writing, it remains unclear whether such a more general homotopy formula can be used. See [21] and Nagel-Rosay [16].

The $\frac{1}{2}$ -estimate for a solution to $\bar{\partial}$ -equations for $(0, 1)$ -forms on strictly pseudoconvex domains with C^2 boundary is obtained by Henkin-Romanov [8]. For $C^{k+\frac{1}{2}}$ -estimates for solutions of $\bar{\partial}$ -equations of degree $(0, 1)$ on strictly pseudoconvex domains with C^m boundary ($m \geq k+4$), see Siu [19]; for $\bar{\partial}$ -equations in higher degree, see Lieb-Range [12]. The C^0 -estimate for a solution to $\bar{\partial}_b$ -equations on M_ρ , without shrinking M_ρ , is given by Henkin [7]. For C^k -estimates for the homotopy formula for $\bar{\partial}_b$ operator on shrinking domains, see [21]. Michel-Ma [13] also obtain C^k -estimates for a modified homotopy formula without shrinking domains, by introducing an extra derivative via $\bar{\partial}_b$.

We want to mention that in estimating (i) and (ii) we need some Hölder inequalities. For the convenience of the reader, we present these inequalities in Appendix A, following the formulation and proofs of Hörmander [9].

The estimates (i) and (ii) will be used to improve regularity in the local CR embedding problem in [6]. To limit the scope of this paper, we leave the estimates in Folland-Stein spaces for future work.

2. Notation and counting derivatives

To simplify notation, set $z' = (z^1, \dots, z^{n-1})$, $z = (z', z^n)$, and

$$x = (\operatorname{Re} z, \operatorname{Im} z') = \pi(z).$$

Analogously, $\xi = (\operatorname{Re} \zeta, \operatorname{Im} \zeta')$. Denote by $|\cdot|$ the Euclidean norms on \mathbb{C}^{n-1} , $\mathbb{C}^{n-1} \times \mathbb{R} = \mathbb{R}^{2n-1}$ and \mathbb{C}^n . Our real hypersurface $M \subset \mathbb{C}^n$ is always a graph over a domain in $\mathbb{C}^{n-1} \times \mathbb{R}$. Let $M_\rho = M \cap \{(x^n)^2 + y^n < \rho^2\}$ and $D_\rho = \pi(M_\rho)$. For the real hyperquadric $M: y^n = |z'|^2$, $\pi(M_\rho)$ is exactly the ball $B_\rho = \{x \in \mathbb{R}^{2n-1}: |x| < \rho\}$. On \mathbb{R}^{2n-1} , we will use the volume-form $dV = d\xi^1 \wedge d\eta^1 \wedge \dots \wedge d\eta^{n-1} \wedge d\xi^n$. On ∂D_ρ , we will need $(2n-2)$ -forms $dV^s = d\xi^1 \wedge \dots \wedge d\hat{\xi}^s \wedge \dots \wedge d\xi^{2n-1}$.

Let $k \geq 0$ be an integer. Denote by $\partial^I u$ a derivative of u of order $|I|$, where I is a standard multi-index. Let $\partial^k u$ denote the set of the k -th order derivatives of u . For a function u on $D \subset \mathbb{R}^{2n-1}$, define

$$\begin{aligned} \|\partial^k u\|_{D,0} &= \sup_{x \in D, |I|=k} |\partial^I u(x)|, & \|u\|_{D,k} &= \max_{0 \leq j \leq k} \|\partial^j u\|_{D,0}, \\ |u|_{D,\alpha} &= \sup_{x,y \in D} \frac{|u(x) - u(y)|}{|x - y|^\alpha}, & 0 < \alpha < 1, \\ \|u\|_{D,k+\alpha} &= \max\{\|u\|_{D,k}, |\partial^I u|_{D,\alpha}: |I| = k\}, & 0 < \alpha < 1. \end{aligned}$$

If $A = (a_i^j)$ is a matrix of C^k functions on D , we define $\|A\|_{D,k} = \max_{i,j} \{\|a_i^j\|_{D,k}\}$. Define $\|A\|_{D,k+\alpha}$ analogously.

To avoid confusion and in the essence of estimates (i)-(ii) in the introduction, the C^a -norm of a function on M_ρ is its C^a -norm on $D_\rho \subset \mathbb{R}^{2n-1}$, defined above.

The $C^a(M_\rho)$ norm of a $(0, q)$ -form $\varphi = \sum \varphi_{\bar{J}} d\bar{z}^{\bar{J}}$ is the maximum of C^a -norms of $\varphi_{\bar{J}}$. The C^a -norm on M_ρ will be denoted by $\|\cdot\|_{C^a(M_\rho)} = \|\cdot\|_{D_\rho,a}$, or simply by $\|\cdot\|_{\rho,a}$ when there is no confusion about M .

Throughout the paper, C_k denotes a constant dependent of k and this dependence will not be expressed sometimes. Constants, such as C, C_1, C_k , might have different values when they reoccur. All constants are independent of M, \hat{r}, ρ, ρ_0 .

Let M be defined by

$$r(z) \stackrel{\text{def}}{=} -y^n + |z'|^2 + \hat{r}(x) = 0, \quad x \in D, \tag{2.1}$$

where $\hat{r} \in C^2(D)$. Throughout the paper, we make the basic assumption

$$\overline{D_{\rho_0}} \subset D, \quad \hat{r}(0) = \hat{r}_z(0) = 0, \quad \epsilon = \|\hat{r}\|_{\rho_0,2} < C_0^{-1}. \tag{2.2}$$

We emphasize that C_0 is a large constant to be adjusted several times. However, it will not depend on any quantity other than n .

Counting derivatives. We need to count derivatives efficiently and use the count to estimate norms. Such a counting scheme is essentially in [9] and we specialize it for two reasons. First, the homotopy formula involves two extra derivatives of the defining function r of M ; second, for each consecutive x -derivative on $\varphi_{\bar{J}} \circ \psi_z^{-1}$, the x -derivative which falls on $\varphi_{\bar{J}}$ yields an extra factor $\partial^2 r(z)$ via the chain rule and $\psi_z(\zeta) = (\zeta' - z', -i2r_z \cdot (\zeta - z))$. We illustrate below how to use the scheme to cope with the two extra derivatives on r and the consecutive derivatives.

Recall that for $l \geq 1$, $\partial^l r(z) = \partial^l r(x)$ is the set of the l -th order derivatives of r . Define

$$\begin{aligned} \partial_*^1 r(\xi, x) &= p\left(\xi, x, (1 + \hat{r}_{x^n} \hat{r}_{\xi^n})^{-1}, r_{z^n}^{-1}, \partial \hat{r}(\xi), \partial \hat{r}(x)\right), \\ \partial_*^2 r(\xi, x) &= p\left(\xi, x, (1 + \hat{r}_{x^n} \hat{r}_{\xi^n})^{-1}, r_{z^n}^{-1}, \partial \hat{r}(\xi), \partial^2 \hat{r}(\xi), \partial \hat{r}(x), \partial^2 \hat{r}(x)\right). \end{aligned} \tag{2.3}$$

Here and in what follows, p is a polynomial with constant coefficients. Its coefficients and degree are bounded in absolute values by a constant depending only on fixed quantities, say k, n . Also, p might be different when it reoccurs. In general, define

$$\partial_*^{2+k} r(\xi, x) = \sum \partial_*^2 r(\xi, x) \partial^{I_1} r(\xi) \cdots \partial^{J_l} r(\xi) \partial^{J_1} r(x) \cdots \partial^{J_l} r(x),$$

where the sum is over finitely many multi-indices I_i, J_i satisfying

$$\sum_{i=1}^j (|I_i| - 2) + \sum_{i=1}^l (|J_i| - 2) \leq k, \quad |I_i| \geq 2, \quad |J_i| \geq 2.$$

We will write $\partial_*^{2+k}r(\xi, x) = \partial_*^{2+k}r(x)$ when it depends only on x . With this abbreviation, we have simple relations

$$\partial_*^{2+k}r \partial_*^{2+j}r = \partial_*^{2+k+j}r, \quad \partial^J \partial_*^{2+k}r = \partial_*^{2+k+|J|}r. \quad (2.4)$$

From [9, Corollary A.6] (see Proposition A.5 in Appendix A), we know that with $D_\rho^2 = D_\rho \times D_\rho$,

$$\prod_{j=1}^m \|f_j\|_{D_\rho^2, k_j+b_j} \leq C_{|a|+|c|+m} \rho^{-b_1-\dots-b_m} \left(\prod_{j=1}^m \|f_j\|_{D_\rho^2, k_j+a_j} + \prod_{j=1}^m \|f_j\|_{D_\rho^2, k_j+c_j} \right)$$

for any non-negative integers k_j and non-negative real numbers a_j, c_j such that (b_1, \dots, b_m) is in the convex hull of $(a_1, \dots, a_m), (c_1, \dots, c_m)$. With the above abbreviation, basic assumption (2.2), and $0 < \rho \leq \rho_0 \leq 3$, one obtains

$$\|\partial_*^{2+k}r\|_{\rho, a} \leq C_{a+k} \rho^{-a-k} \|r\|_{\rho, 2+k+a}, \quad \|r\|_{\rho, 2+k+a} \stackrel{\text{def}}{=} 1 + \|\hat{r}\|_{\rho, 2+k+a} \quad (2.5)$$

for all real numbers $a \geq 0$ and integers $k \geq 0$.

We will also need a chain rule. Recall that $\psi_z: \zeta'_* = \zeta' - z', \zeta_*^n = -2ir_z \cdot (\zeta - z)$ and define

$$\Psi(\xi, x) = (\pi \psi_z(\zeta), x), \quad z = \pi|_M^{-1}(x), \quad \zeta = \pi|_M^{-1}(\xi).$$

Let $0 < \rho \leq \rho_0 \leq 3$. We will show that $B_{\rho/2} \subset D_\rho \subset B_{2\rho}$ and

$$W_\rho = \Psi(D_\rho \times D_\rho) \subset B_{9\rho} \times D_\rho.$$

(See Lemmas 5.1-5.2.) The Jacobean matrix of Ψ depends only on derivatives of r of order ≤ 2 and has determinant $1 + \hat{r}_{x^n} \hat{r}_{\xi^n}$ (by (6.2)). Then the chain rule takes simple forms. Let $(\Psi^{-1})^j$ be the j -th component of Ψ^{-1} . Then

$$\begin{aligned} \partial^I \{(\Psi^{-1})^j\} &= \partial_*^2 r \circ \Psi^{-1}, \quad |I| = 1; \\ \partial^K (f \circ \Psi^{-1}) &= \sum_{|L| \leq |K|} (\partial^L f \partial_*^{2+|K|-|L|} r) \circ \Psi^{-1}. \end{aligned} \quad (2.6)$$

We will show that the Lipschitz constant of Ψ^{-1} on W_ρ is bounded by C (see Lemma 5.2). Then taking Hölder ratio in (2.6) with $k = [a]$ gives us

$$\|f \circ \Psi^{-1}\|_{W_\rho, a} \leq C_k \rho^{-a} (\|f\|_{D_\rho \times D_\rho, a} + \|f\|_{D_\rho \times D_\rho, 0} \|r\|_{\rho, 2+a}). \quad (2.7)$$

Note that the above mentioned $\varphi_{\bar{J}} \circ \psi_z^{-1}$ is a special case.

The above counting scheme via (2.4)-(2.7) and its variants will be used systematically.

3. The Henkin homotopy formula

In this section we recall the homotopy formula by following the formulation in [21]. We discuss the formula for differential forms of low regularity.

Let $M \subset \mathbb{C}^n : r = 0$ be given by (2.1)-(2.2). We first recall a representative $\bar{\partial}_M$ for the $\bar{\partial}_b$ -operator. By definition, a (p, q) -form φ of class \mathcal{C}^a on M is the restriction of some (p, q) -form $\tilde{\varphi}$ of class \mathcal{C}^a in a neighborhood of M . If $a \geq 1$, we define $\bar{\partial}_b\varphi$ to be the restriction of $\bar{\partial}\tilde{\varphi}$ to M . Notice that on M , $\theta = -2i\partial r = \bar{\theta}$ and that $\bar{\partial}_b\varphi$ is well-defined modulo $\bar{\partial}r$ when $0 \leq p < n$, and it is actually well-defined when $p = n$. By a *tangential* $(0, q)$ -form φ , we mean a form $\varphi = \sum_{|I|=q} \varphi_I d\bar{z}^I$ with $d\bar{z}^I = (d\bar{z}^1, \dots, d\bar{z}^{n-1})$. A continuous $(0, q)$ -form φ can be written uniquely as $\varphi' + \varphi'' \wedge \bar{\theta}$ for some tangential forms φ', φ'' . Define

$$\bar{X}_\alpha = \partial_{\bar{z}^\alpha} - \frac{r_{\bar{z}^\alpha}}{r_{\bar{z}^n}} \partial_{\bar{z}^n}, \quad \bar{\partial}_M\varphi = \bar{\partial}_M\varphi' = \sum_{|I|=q} \sum_{1 \leq \alpha < n} \bar{X}_\alpha \varphi'_I d\bar{z}^\alpha \wedge d\bar{z}^I.$$

Then a straightforward computation shows that

$$\bar{\partial}_b\varphi = \bar{\partial}_M\varphi \quad \text{mod } \theta.$$

Each (n, q) -form on M can be written as $\varphi'' \wedge d\bar{z}^1 \wedge \dots \wedge d\bar{z}^{n-1} \wedge \theta$ where φ'' is tangential. If $\varphi'' \in \mathcal{C}^1$, then

$$\bar{\partial}_b(\varphi'' \wedge d\bar{z}^1 \wedge \dots \wedge d\bar{z}^{n-1} \wedge \theta) = \bar{\partial}_M\varphi'' \wedge d\bar{z}^1 \wedge \dots \wedge d\bar{z}^{n-1} \wedge \theta. \quad (3.1)$$

Let φ be a continuous $(0, q)$ -form on a domain $\mathcal{U} \subset M$, and ϕ be a continuous $(0, q-1)$ -form. Suppose that $q \geq 1$. We say that $\bar{\partial}_b\phi = \varphi \quad \text{mod } \theta$ holds on \mathcal{U} as currents, if $\int_M \phi \wedge \bar{\partial}_b\psi = (-1)^q \int_M \varphi \wedge \psi$ for all \mathcal{C}^1 -smooth $(n, n-q-1)$ -forms ψ with compact support in \mathcal{U} .

Write $r_z = r_z(z)$ and $r_\zeta = r_\zeta(\zeta)$. Set $N_0(\zeta, z) = r_\zeta \cdot (\zeta - z)$, $S_0(\zeta, z) = r_z \cdot (\zeta - z)$ and

$$\Omega_{0,q-1}^{+-}(\zeta, z) = \frac{\partial_\zeta r \wedge (r_z \cdot d\zeta) \wedge (\partial_{\bar{\zeta}}\partial_\zeta r)^{n-1-q} \wedge (\partial_{\bar{z}}r_z \wedge d\zeta)^{q-1}}{N_0^{n-q}(\zeta, z)S_0^q(\zeta, z)}, \quad (3.2)$$

$$\Omega_{0,q-1}^{0+-}(\zeta, z) = \frac{d\zeta^n \wedge \partial_\zeta r \wedge (r_z \cdot d\zeta) \wedge (\partial_{\bar{\zeta}}\partial_\zeta r)^{n-2-q} \wedge (\partial_{\bar{z}}r_z \wedge d\zeta)^{q-1}}{(\zeta^n - z^n)N_0^{n-q-1}(\zeta, z)S_0^q(\zeta, z)}. \quad (3.3)$$

Note that $\theta(\zeta)$ annihilates $\Omega_{0,q-1}^{+-}(\zeta, z)$ and $\Omega_{0,q-1}^{0+-}(\zeta, z)$. Let $n \geq 4$ and $0 < q < n-2$. For a tangential $(0, q)$ -form φ on \overline{M}_ρ , we have the homotopy formula

$$\varphi = \bar{\partial}_M(P'_0 + P'_1)\varphi + (Q'_0 + Q'_1)\bar{\partial}_M\varphi, \quad z \in M_\rho. \quad (3.4)$$

Here P'_0, P'_1, Q'_0, Q'_1 are tangential parts of

$$\begin{aligned} P_0\varphi(z) &= c_1 \int_{M_\rho} \varphi \wedge \Omega_{0,q-1}^{+-}(\zeta, z), & P_1\varphi(z) &= c_2 \int_{\partial M_\rho} \varphi \wedge \Omega_{0,q-1}^{0+-}(\zeta, z), \\ Q_0\psi(z) &= c_3 \int_{M_\rho} \psi \wedge \Omega_{0,q}^{+-}(\zeta, z), & Q_1\psi(z) &= c_4 \int_{\partial M_\rho} \psi \wedge \Omega_{0,q}^{0+-}(\zeta, z). \end{aligned} \tag{3.5}$$

From now on, all $(0, q)$ -forms φ on M are tangential. By $\|\varphi\|_{\rho,a}$, we mean the norm $\|\varphi'\|_{\rho,a}$ as defined in Section 2, where $\varphi' = \varphi \bmod \theta$ and φ' is tangential. By $\|P\varphi\|_{\rho,a}$ as used in the introduction, where P is either of solution operators in the homotopy formula, we mean $\|P'\varphi\|_{\rho,a}$. By an abuse of notation, φ stands for forms on M_ρ and $D_\rho = \pi(M_\rho)$.

Next, we describe kernels of P'_j, Q'_j on domain D_ρ via coordinates x . We have

$$r_\zeta \cdot d\zeta \wedge r_z \cdot d\zeta = \sum_{1 \leq j, l \leq n} r_{\zeta^l} (r_{\zeta^j} - r_{z^j}) d\zeta^j \wedge d\zeta^l, \tag{3.6}$$

$$r_\zeta \cdot d\zeta \wedge r_z \cdot d\zeta \wedge d\zeta^n = \sum_{1 \leq \alpha, \beta < n} r_{\zeta^\beta} (r_{\zeta^\alpha} - r_{z^\alpha}) d\zeta^\alpha \wedge d\zeta^\beta \wedge d\zeta^n. \tag{3.7}$$

Assume now that $\zeta, z \in M_\rho$. We compute $d\bar{\zeta}^n$ and $d\bar{z}^n$ in different ways. We keep the latter a $(0, 1)$ -form and find its tangential part. We have

$$d\bar{z}^n = -\frac{r_{\bar{z}}}{r_{\bar{z}^n}} \cdot d\bar{z}^n \bmod \theta(z), \tag{3.8}$$

$$d\zeta^n = (1 + i\hat{r}_{\xi^n})d\xi^n + i2 \operatorname{Re}\{r_{\zeta'} \cdot d\zeta'\} = 2ir_{\bar{\zeta}^n}d\xi^n + i2 \operatorname{Re}\{r_{\zeta'} \cdot d\zeta'\}. \tag{3.9}$$

Note that in (3.9), we have used $r(\zeta) = -\eta^n + |\zeta'|^2 + \hat{r}(\xi) = 0$. In (3.6)-(3.7) and $\bar{\partial}_z r_z \wedge d\zeta$, we use (3.8)-(3.9) to rewrite $d\bar{z}^n$ and $d\zeta^n$, respectively. In $\partial_{\bar{\zeta}} \partial_{\zeta'} r$, we use (3.9) to rewrite $d\zeta^n, d\bar{\zeta}^n$. From (3.2)-(3.3), we obtain on M_ρ

$$P'_0\varphi(x) = \sum_{|I|=q-1} \sum_{|J|=q} \sum_{1 \leq j \leq n} d\bar{z}^{\bar{I}} \int_{D_\rho} A_I^{j\bar{J}}(\xi, x) \frac{\varphi_{\bar{J}}(\xi)(r_{\zeta^j} - r_{z^j})}{(N_0^{n-q} S_0^q)(\zeta, z)} dV(\xi), \tag{3.10}$$

$$P'_1\varphi(x) = \sum_{|I|=q-1} \sum_{|J|=q} \sum_{\alpha, \beta=1}^{n-1} \sum_{s=1}^{2n-1} d\bar{z}^{\bar{I}} \int_{\partial D_\rho} \frac{B_{I_s}^{\alpha\beta\bar{J}}(\xi, x) \varphi_{\bar{J}}(\xi)(r_{\zeta^\alpha} - r_{z^\alpha}) r_{\zeta^\beta}}{(\zeta^n - z^n)(N_0^{n-q-1} S_0^q)(\zeta, z)} dV^s(\xi). \tag{3.11}$$

Here $A_I^{j\bar{J}}$ and $B_{I_s}^{\alpha\beta\bar{J}}$ are polynomials in $(r_\zeta, r_{\bar{\zeta}}, r_{\zeta\bar{\zeta}}, r_{\bar{z}}, 1/r_{\bar{z}^n}, r_{z\bar{z}})$. We make a remark for the case $q = 1$. In this case we need to remove $\sum_{|I|=0}$ in (3.10)-(3.11).

Also, $A_I^{j\bar{J}} \stackrel{\text{def}}{=} A_I^{j\bar{J}}$ and $B_s^{\alpha\beta\bar{J}} \stackrel{\text{def}}{=} B_{I_s}^{\alpha\beta\bar{J}}$ are independent of z . This observation will play a role in the $\frac{1}{2}$ -estimate of $P'\varphi$ when φ is a $(0, 1)$ -form.

See also Chen-Shaw [2] for homotopy formulae. In this paper we replace the strict convexity of defining function r in [2] by the condition (2.2) with $0 < \rho_0 \leq 3$;

see Appendix B for details. We remark that the homotopy formula (3.4) holds as currents, when φ and $\bar{\partial}_b\varphi$ on M_ρ admit continuous extensions to $\overline{M_\rho}$. See Appendix B for a proof by using the Friedrichs approximation theorem. Henkin [7] formulated $\bar{\partial}_b$ in the sense of currents. Shaw [18] also used the Friedrichs approximation for $\bar{\partial}_b$ -solutions.

4. Kernels and the approximate Heisenberg transformation

In this section, we will describe briefly the new kernels when the approximate Heisenberg transformation is applied. The contents of next few sections are indicated at the end of this section.

Recall that in (3.10) functions $A_{\bar{J}}^{j\bar{J}}$ have the form $\partial_*^2 r$. Hence coefficients of $P'_0\varphi$ are sums over $|J|=q$ and $1 \leq j \leq n$ of $\mathcal{K}_{\varphi_{\bar{J}}}(x) = \int_{D_\rho} \varphi_{\bar{J}}(\xi) k_{ab}^j(\xi, x) dV(\xi)$. Here $\zeta, z \in M$ and

$$k(\xi, x) = k_{ab}^j(\xi, x) = \frac{\partial_*^2 r(\xi, x)(r_{\zeta j} - r_{z j})}{(r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b}, \quad a = n - q, \quad b = q.$$

To understand the kernel, let us compute its denominator. Set $\hat{r}(z) = \hat{r}(x)$. On $M \times M$ we have

$$\begin{aligned} r_z \cdot (\zeta - z) &= \bar{z}' \cdot (\zeta' - z') + \frac{i}{2}(\zeta^n - z^n) + \hat{r}_z \cdot (\zeta - z) \\ &= \bar{z}' \cdot (\zeta' - z') - \frac{1}{2}(|\zeta'|^2 - |z'|^2) + \frac{i}{2}(\xi^n - x^n) \\ &\quad - \frac{1}{2}(\hat{r}(\zeta) - \hat{r}(z)) + \hat{r}_z \cdot (\zeta - z) \\ &= i \operatorname{Im}(r_z \cdot (\zeta - z)) - \frac{1}{2}|\zeta' - z'|^2 \\ &\quad - \frac{1}{2}(\hat{r}(\zeta) - \hat{r}(z)) + \operatorname{Re}(\hat{r}_z \cdot (\zeta - z)). \end{aligned}$$

Also

$$\begin{aligned} r_\zeta \cdot (\zeta - z) &= r_z \cdot (\zeta - z) + (r_\zeta - r_z) \cdot (\zeta - z) \\ &= i \operatorname{Im}(r_z \cdot (\zeta - z)) + \frac{1}{2}|\zeta' - z'|^2 + (\hat{r}_\zeta - \hat{r}_z) \cdot (\zeta - z) \\ &\quad - \frac{1}{2}\{\hat{r}(\zeta) - \hat{r}(z) - 2 \operatorname{Re}(\hat{r}_z \cdot (\zeta - z))\}. \end{aligned}$$

The first two terms in $r_z \cdot (\zeta - z)$ and $r_\zeta \cdot (\zeta - z)$ can be simplified simultaneously. Define

$$N(\zeta, z) \stackrel{\text{def}}{=} |\zeta' - z'|^2 + 2i \operatorname{Im}(r_z \cdot (\zeta - z)). \quad (4.1)$$

Then we arrive at the basic relations

$$-2r_z \cdot (\zeta - z) = \overline{N(\zeta, z)} + A(\zeta, z), \quad 2r_\zeta \cdot (\zeta - z) = N(\zeta, z) + B(\zeta, z)$$

with

$$A(\zeta, z) = \hat{r}(\zeta) - \hat{r}(z) - 2\operatorname{Re}(\hat{r}_z \cdot (\zeta - z)), \quad B(\zeta, z) = 2(\hat{r}_\zeta - \hat{r}_z) \cdot (\zeta - z) - A(\zeta, z).$$

By the condition (2.2) on \hat{r} , we will show that D_{ρ_0} is convex and hence

$$|A(\zeta, z)| \leq C\epsilon|\zeta - z|^2, \quad |B(\zeta, z)| \leq C\epsilon|\zeta - z|^2, \quad \zeta, z \in M_{\rho_0}.$$

We will show that $|N(\zeta, z)| \geq C^{-1}|\zeta - z|^2$ on $M_{\rho_0} \times M_{\rho_0}$. Consequently, the kernel of $P'_0\varphi$ is factored as

$$k(\xi, x) = T_1(\xi, x)^{-a} T_2(\xi, x)^{-b} \frac{\partial_*^2 r(\xi, x)(r_{\zeta j} - r_{z j})}{N^a(\zeta, z) \overline{N^b(\zeta, z)}},$$

$$T_1(\xi, x) = 1 + N^{-1}(\zeta, z)B(\zeta, z), \quad T_2(\xi, x) = 1 + \overline{N^{-1}(\zeta, z)}A(\zeta, z)$$

for $\zeta, z \in M_{\rho_0}$. Moreover, $\min\{|T_1(\xi, x)|, |T_2(\xi, x)|\} \geq 1/4$ when (ξ, x) is on $D_{\rho_0} \times D_{\rho_0}$ and off its diagonal. Now, identity (4.1) suggests the following approximate Heisenberg transformation

$$\psi_z: \zeta'_* = \zeta' - z', \quad \zeta_*^n = -2ir_z \cdot (\zeta - z).$$

We will show that for fixed $z \in M_{\rho_0}$, $\zeta'_* = \zeta' - z'$, $\xi_*^n = 2\operatorname{Im}(r_z \cdot (\zeta - z))$ indeed form coordinates of M_{ρ_0} . Thus, with $\zeta, z \in M_{\rho_0}$

$$N(\zeta, z) = |\zeta'_*|^2 + i\xi_*^n \stackrel{\text{def}}{=} N_*(\xi_*),$$

$$k(\xi, x) = \sum_{|I|=1} E_I(\xi_*, x) \xi_*^I N_*(\xi_*)^{-a} \overline{N_*^{-b}(\xi_*)}, \quad a + b = n.$$

When φ has compact support, the decomposition for $k(\xi, x)$ allows us to take x -derivatives of $K\varphi_{\mathcal{J}}(x)$ directly onto coefficients $E_I(\xi_*, x)$ and onto $\varphi_{\mathcal{J}}(\psi_z^{-1}(\xi_*))$, without destroying the integrability of the kernels.

We now describe the contents of next few sections.

We will carry out the details of this section in Sections 5, 6, and 8. In Section 5 we will also study how domains B_ρ , $D_\rho = \pi(M_\rho)$ and $\pi\psi_z(M_\rho)$ are nested. We will need Hölder inequalities in Appendix A for domains D_ρ . Therefore, we will verify that under the basic assumption (2.2) and $0 < \rho \leq \rho_0 \leq 3$, D_ρ is convex and $B_{\rho/2} \subset D_\rho \subset B_{2\rho}$. In Section 6, we will also express the graph $\psi_z(M_{\rho_0})$ for $z \in M_{\rho_0}$ as

$$\eta_*^n = \sum_{|I|=2} h_I(\xi_*, x) \xi_*^I, \quad \xi_* = (\operatorname{Re} \zeta_*, \operatorname{Im} \zeta'_*).$$

The latter will be used to show $|r_z \cdot (\zeta - z)| \geq C^{-1}(\rho\sigma)^2$ for $z \in M_{(1-\sigma)\rho}$ and $\zeta \in \partial M_\rho$. In Sections 9 and 10, we derive the C^a -estimates. The $\frac{1}{2}$ -estimate is in Section 12, following a reduction for $C^{k+\frac{1}{2}}$ -estimates in Section 11.

We conclude this section with a lower bound for $|\zeta^n - z^n|$. By the basic assumption (2.2), D_{ρ_0} is relatively compact in D and hence $(x^n)^2 + y^n = \rho^2$ on ∂M_ρ for $0 < \rho \leq \rho_0$. Then the image of the projection of ∂M_ρ in the z^n -plane is contained in the parabola $(x^n)^2 + y^n = \rho^2$. Therefore, for $z \in M_{(1-\sigma)\rho}$ and $\zeta \in \partial M_\rho$ with $\rho \leq \rho_0 (\leq 3)$, we obtain

$$|\zeta^n - z^n| \geq C^{-1}\rho^2\sigma. \quad (4.2)$$

5. Domains and images under the transformation

Recall that our real hypersurface M is given by (2.1)-(2.2), and

$$M_\rho = M \cap \{(x^n)^2 + y^n < \rho^2\}, \quad D_\rho = \pi(M_\rho) = \{x \in D : |x|^2 + \hat{r}(x) < \rho^2\}.$$

Lemma 5.1. *Let M satisfy (2.2). Suppose that $0 < \rho \leq \rho_0$. Then $\overline{D_\rho}$ is strictly convex with C^2 boundary. Also,*

$$B_{(1-c\epsilon)\rho} \subset D_\rho \subset B_{(1+c\epsilon)\rho}, \quad C^{-1}\rho\sigma \leq \text{dist}(\partial D_{(1-\sigma)\rho}, \partial D_\rho) \leq C\rho\sigma.$$

Moreover, constants C_0, c, C are independent of ρ and \hat{r} .

Proof. Let $\epsilon = \|\hat{r}\|_{\rho_0, 2} < C_0^{-1}$. The strict convexity follows from the positivity of the Hessian of $\phi(x) = (x^n)^2 + y^n = |x|^2 + \hat{r}(x)$ on $\overline{D_\rho}$. Since $0 \in D_\rho$, then $|\hat{r}(x)| \leq C\epsilon|x|^2$ on D_ρ . Now,

$$(1 - C\epsilon)|x|^2 \leq \phi(x) \leq (1 + C\epsilon)|x|^2.$$

In particular, for a possibly larger C , we have $B_{(1-C\epsilon)^{1/2}\rho} \subset D_\rho \subset B_{(1+C\epsilon)^{1/2}\rho}$, since $\overline{D_{\rho_0}} \subset D$ implies that $\phi = \rho^2$ on ∂D_ρ for all $0 < \rho \leq \rho_0$.

Let $x \in \partial D_{(1-\sigma)\rho}$. Then $\phi(x) = ((1-\sigma)\rho)^2$ and $\frac{1}{2}(1-\sigma)\rho < |x| < 2(1-\sigma)\rho$. For $y \in D_\rho$, we have $|\phi(y) - \phi(x)| \leq C\|\partial^1\phi\|_{\rho, 0}|y-x|$. We get

$$|\phi(y)| \leq (1-\sigma)^2\rho^2 + C\|\partial^1\phi\|_{\rho, 0}|y-x| \leq (1-\sigma)^2\rho^2 + C\rho|y-x|.$$

Hence, $\phi(y) < \rho$ for $|y-x| \leq C^{-1}\rho\sigma$. This shows that $\text{dist}(\partial D_{(1-\sigma)\rho}, \partial D_\rho) \geq C^{-1}\rho\sigma$. On the other hand, if $y = (1+t)x \in D_\rho$ and $t > 0$, applying the mean-value-theorem to $\phi((1+s)x)$ for $0 \leq s \leq t$ yields

$$\begin{aligned} \phi(y) &\geq ((1-\sigma)\rho)^2 + x \cdot (y-x) - C\epsilon|y||y-x| \\ &\geq \rho^2(1-\sigma)^2 + t|x|^2 - C'\epsilon\rho t|x| \\ &\geq \rho^2\{(1-\sigma)^2 + \frac{1}{4}t(1-\sigma)^2 - 2C''\epsilon t\}. \end{aligned}$$

This show that $\phi((1+t)x) > \rho^2$ if $(1+t)x \in D_\rho$ and $t > C\sigma$, a contradiction. Therefore, $\text{dist}(\partial D_{(1-\sigma)\rho}, \partial D_\rho) \leq C\rho\sigma$. \square

Recall the approximate Heisenberg transformation

$$\psi_z: \zeta'_* = \zeta' - z', \quad \zeta_*^n = -2ir_z \cdot (\zeta - z).$$

Before applying ψ_z to kernels $k_{ab}^j(\zeta, z)$, we need to know how it transforms M_ρ . Recall our notation $x \in \mathbb{R}^{2n-1}$ and $\xi \in \mathbb{R}^{2n-1}$. Define a map $\tilde{\psi}_x$ by relations

$$\xi_* = \tilde{\psi}_x(\xi) = \pi\psi_z(\zeta), \quad \zeta, z \in M_{\rho_0}.$$

Set $\Psi(\xi, x) = (\tilde{\psi}_x(\xi), x)$. We have the following.

Lemma 5.2. *Let $M: y^n = |z'|^2 + \hat{r}(z', x^n)$ satisfy (2.2) with $0 < \rho_0 \leq 3$. There exist constants C, C_m , independent of \hat{r}, ρ and ρ_0 , such that the following hold.*

(i) *If $x \in D_\rho$ and $0 < \rho \leq \rho_0$ then*

$$\tilde{\psi}_x(D_\rho) \subset B_{9\rho}, \quad \Psi(D_\rho \times D_\rho) \subset B_{9\rho} \times D_\rho.$$

(ii) *For $u, v \in D_{\rho_0} \times D_{\rho_0}$,*

$$C^{-1}|v - u| \leq |\Psi(v) - \Psi(u)| \leq C|v - u|.$$

In particular, if $z \in M_{\rho_0}$ then $\psi_z(M_{\rho_0})$ is a graph over $\pi\psi_z(M_{\rho_0})$.

Proof. Let $R(x) = |z'|^2 + \hat{r}(x)$. By (2.2), $\overline{D_{\rho_0}} \subset D$ and $\epsilon = \|\hat{r}\|_{\rho_0, 2} < C_0^{-1}$. For brevity, set $M = M_{\rho_0}$.

(i). Assume that $0 < \rho \leq \rho_0$. Since D_{ρ_0} is convex, we have $|\hat{r}(x)| \leq C\epsilon|x|^2$ on D_{ρ_0} . The map $\tilde{\psi}_x$ is defined by $\zeta'_* = \zeta' - z'$ and

$$\begin{aligned} \xi_*^n &= \xi^n - x^n + \hat{r}_{x^n}(R(\xi) - R(x)) + 2\text{Im}[R_{z'} \cdot (\zeta' - z')] \\ &= \xi^n - x^n + 2\text{Im}(\overline{z'} \cdot \zeta') + \hat{r}_{x^n}(R(\xi) - R(x)) + 2\text{Im}[\hat{r}_{z'} \cdot (\zeta' - z')]. \end{aligned} \quad (5.1)$$

Let ξ, x be in D_ρ . Recall that $D_\rho \subset B_{(1+C\epsilon)\rho}$. We have

$$\begin{aligned} |x| &< (1+C\epsilon)\rho, \quad |\xi| < (1+C\epsilon)\rho, \quad |\hat{r}_{x^n}| \leq \epsilon, \quad |\hat{r}_{z'}| \leq \epsilon, \\ |R(\xi)| &\leq |\zeta'|^2 + C\epsilon|\xi|^2 < (1+C'\epsilon)\rho^2. \end{aligned}$$

Thus, $|\zeta'_*|^2 = |\zeta' - z'|^2 \leq 2|\zeta'|^2 + 2|z'|^2 < 4(1+C\epsilon)\rho^2$; by (5.1), $|\xi_*^n|^2 \leq (2\rho + 2\rho^2 + C\epsilon\rho)^2$. Since $2\rho^2 \leq 6\rho$ then

$$|\zeta'_*|^2 + |\xi_*^n|^2 \leq 4(1+C\epsilon)\rho^2 + (2\rho + 2\rho^2 + C\epsilon\rho)^2 < (9\rho)^2,$$

if ϵ is sufficiently small. We get $\tilde{\psi}_x(D_\rho) \subset B_{9\rho}$.

(ii). It is obvious that the Lipschitz constant of Ψ on the convex domain $D_{\rho_0} \times D_{\rho_0}$ is bounded by some C . Let $\xi, x, \tilde{\xi}, \tilde{x}$ be in D_{ρ_0} . We need to show that $|\Psi(\tilde{\xi}, \tilde{x}) - \Psi(\xi, x)| \geq C^{-1} |(\tilde{\xi}, \tilde{x}) - (\xi, x)|$. Write $\xi = (\zeta', \xi^n), x = (z', x^n)$. Then

$$|\Psi(\tilde{\xi}, \tilde{x}) - \Psi(\xi, x)| \geq |(\tilde{\zeta}' - \tilde{z}' - \zeta' + z', \tilde{x} - x)| \geq |(\tilde{\zeta}' - \zeta', \tilde{x} - x)|/C.$$

Set $\tilde{\xi}_* = \tilde{\psi}_{\tilde{x}}(\tilde{\xi})$ and $\xi_* = \psi_x(\xi)$. It suffices to show that

$$|\tilde{\xi}_*^n - \xi_*^n| \geq |\tilde{\xi}^n - \xi^n|/C$$

if $|(\tilde{\zeta}' - \zeta', \tilde{x} - x)| < \frac{1}{48} |\tilde{\xi}^n - \xi^n|$. Assume that the latter holds. Recall that $0 < \rho_0 \leq 3$ and $D_{\rho_0} \subset B_{2\rho_0}$. We have $\max\{|z'|, |\zeta'|\} < 2\rho_0$. Now the second identity in (5.1) implies that

$$\begin{aligned} |\tilde{\xi}_*^n - \xi_*^n| &\geq |\tilde{\xi}^n - \xi^n| - |\tilde{x}^n - x^n| - 2|\tilde{\zeta}'||\tilde{z}' - z'| - 2|z'||\tilde{\zeta}' - \zeta'| - C\epsilon |(\tilde{\xi} - \xi, \tilde{x} - x)| \\ &\geq (1 - \frac{1}{48})|\tilde{\xi}^n - \xi^n| - 8\rho_0(|\tilde{\zeta}' - \zeta'| + |\tilde{z}' - z'|) - C'\epsilon |\tilde{\xi}^n - \xi^n|. \end{aligned}$$

Thus, $|\tilde{\xi}_*^n - \xi_*^n| \geq (1 - \frac{1}{48} - \frac{8 \cdot 3 \cdot \sqrt{2}}{48} - C'\epsilon) |\tilde{\xi}^n - \xi^n| \geq |\tilde{\xi}^n - \xi^n|/4$.

That $\psi_z(M_{\rho_0})$ is a graph follows from the injectivity of Ψ . □

6. Estimates on $r_z \cdot (\zeta - z), r_\zeta - r_z$ and $\psi_z(M)$ via Taylor's theorem

To transform $k(\zeta, z)$ via ψ_z , we need expansions of $r_z \cdot (\zeta - z), r_\zeta \cdot (\zeta - z)$ and $r_{\zeta^j} - r_{z^j}$ in new variables ξ_* . We will find these expansions via Taylor's theorem.

Let us recall Taylor's theorem. Assume that $0 < \rho \leq \rho_0 \leq 3$ and \hat{r} satisfies (2.2). So D_ρ is strictly convex. If f is a complex-valued function on the convex set D_ρ , we define $\mathcal{R}_k f$ and $\mathcal{R}_I f$ on $D_\rho \times D_\rho$ by

$$\begin{aligned} \mathcal{R}_k f(y, x) &\equiv f(y) - \sum_{0 \leq j \leq k-1} \frac{1}{j!} \partial_t^j |_{t=0} f(x + t(y-x)) \\ &= \frac{1}{(k-1)!} \int_0^1 (1-t)^{k-1} \partial_t^k f(x + t(y-x)) dt \\ &= \sum_{|I|=k} \mathcal{R}_I f(y, x) (y-x)^I. \end{aligned}$$

Denote by $\mathcal{R}^k f$ the set of coefficients $\mathcal{R}_I f$ with $|I| = k$. For any real number $a \geq 0$,

$$\|\mathcal{R}_I f\|_{D_\rho \times D_\rho, a} \leq C_{a+|I|} \|f\|_{\rho, a+|I|}. \tag{6.1}$$

We will also need to express the remainders in $\xi_* = (\text{Re } \zeta_*, \text{Im } \zeta'_*)$. Let $z \in M_{\rho_0}$ and $\pi(z) = x \in D_{\rho_0}$. By Lemma 5.2, $\psi_z(M_{\rho_0})$ is a graph

$$\eta_*^n = h(\xi_*, x), \quad \xi_* \in \tilde{\psi}_x(D_{\rho_0}).$$

By $\zeta' - z' = \zeta'_*$ and $\zeta_*^n = -2ir_z \cdot (\zeta - z)$, we have

$$\zeta_*^n = \zeta^n - z^n - i\hat{r}_{x^n}(\zeta^n - z^n) - 2iR_{z'} \cdot \zeta'_*.$$

Here $R(x) = |z'|^2 + \hat{r}(x)$. Computing the real and imaginary parts, we get

$$\xi_*^n = \xi^n - x^n + \hat{r}_{x^n}(R(\xi) - R(x)) + 2\text{Im}(R_{z'} \cdot \zeta'_*), \quad (6.2)$$

$$\eta_*^n = |\zeta'_*|^2 + \hat{r}(\xi) - \hat{r}(x) - \hat{r}_{x^n}(\xi^n - x^n) - 2\text{Re}(\hat{r}_{z'} \cdot \zeta'_*). \quad (6.3)$$

In (6.2), replace $R(\xi) - R(x)$ by

$$\sum_{|I|=1} \mathcal{R}_I R(\xi, x)(\xi - x)^I = \sum_{|I|=1} \mathcal{R}_I R(\xi, x)(\xi'_*, \xi^n - x^n)^I$$

and then solve for $\xi^n - x^n$. We get

$$\xi^n - x^n = \sum_{|I|=1} p_I(\xi, x)\xi_*^I, \quad (6.4)$$

where $p_I(\xi, x)$ are of the form

$$p\left(\xi, x, \frac{1}{1 + \hat{r}_{x^n} \mathcal{R}_{(0',1)} \hat{r}(\xi, x)}, \partial \hat{r}(x), \mathcal{R}^1 \hat{r}(\xi, x)\right)$$

for some polynomials p . With these polynomials p , introduce notation

$$\mathcal{R}_*^i g = \sum_{|I|=i} p\left(\xi, x, \frac{1}{1 + \hat{r}_{x^n} \mathcal{R}_{(0',1)} \hat{r}(\xi, x)}, \partial \hat{r}(x), \mathcal{R}^1 \hat{r}(\xi, x)\right) \mathcal{R}_I g(\xi, x). \quad (6.5)$$

Note that reappearing p , $\mathcal{R}_*^i g$ may be different. We now express Taylor remainders in variables ζ'_* , ξ_*^n as follows

$$\mathcal{R}_k f(\xi, x) = \sum_{|L|=k} \mathcal{R}_L f(\xi, x)(\xi - x)^L = \sum_{|L|=k} \mathcal{R}_*^k f(\xi, x)\xi_*^L, \quad (6.6)$$

$$|\mathcal{R}_*^l \hat{r}(\xi, x)| \leq C \|\hat{r}\|_{\rho_0, 2}, \quad l \leq 2, \quad \xi, x \in D_{\rho_0}.$$

We now apply notation (6.5). Using (6.4) in (6.3) we get

$$\eta_*^n = |\zeta'_*|^2 + \sum_{|I|=1} \mathcal{R}_*^1 \hat{r}(\xi, x)\xi_*^I, \quad (6.7)$$

$$\eta_*^n = |\zeta'_*|^2 + \mathcal{R}_2 \hat{r}(\xi, x) = |\zeta'_*|^2 + \sum_{|I|=2} \mathcal{R}_*^2 \hat{r}(\xi, x)\xi_*^I. \quad (6.8)$$

Set $r(z) = -y^n + |z'|^2 + \hat{r}(z', x^n)$. We have defined

$$N(\zeta, z) = |\zeta' - z'|^2 + 2i \text{Im}(r_z \cdot (\zeta - z)), \quad N_*(\xi_*) = |\zeta'_*|^2 + i\xi_*^n.$$

Recall that for $(\zeta, z) \in M_{\rho_0} \times M_{\rho_0}$ we have

$$-2r_z \cdot (\zeta - z) = \overline{N(\zeta, z)} + \mathcal{R}_2 \hat{r}(\xi, x), \quad 2r_\zeta \cdot (\zeta - z) = N(\zeta, z) + B(\zeta, z), \quad (6.9)$$

$$B(\zeta, z) = 2(\hat{r}_\zeta - \hat{r}_z) \cdot (\zeta - z) - \mathcal{R}_2 \hat{r}(\xi, x). \quad (6.10)$$

Define

$$\mathcal{R}_*^i \partial \hat{r} = \sum_{1 \leq j \leq n} \mathcal{R}_*^i \hat{r}_{x_j} + \sum_{1 \leq j < n} \mathcal{R}_*^i \hat{r}_{y_j}. \quad (6.11)$$

We have

$$\begin{aligned} (\hat{r}_\zeta - \hat{r}_z) \cdot (\zeta - z) &= \sum_{|K|=2} \mathcal{R}_*^1 \partial \hat{r}(\xi, x) \xi_*^K, \\ B(\zeta, z) &= \sum_{|L|=2} (\mathcal{R}_*^2 \hat{r}(\xi, x) + \mathcal{R}_*^1 \partial \hat{r}(\xi, x)) \xi_*^L. \end{aligned} \quad (6.12)$$

For the numerator of the kernel, we have

$$r_\zeta - r_z = (\zeta'_*, 0) + \sum_{|I|=1} (\mathcal{R}_*^1 \hat{r}, \dots, \mathcal{R}_*^1 \partial \hat{r})(\xi, x) \xi_*^I. \quad (6.13)$$

In summary, we have proved the following expansions.

Lemma 6.1. *Let $M: y^n = |z'|^2 + \hat{r}(z', x^n)$ satisfy (2.2) with $0 < \rho_0 \leq 3$. Suppose that $\zeta, z \in M_{\rho_0}$ and $\zeta_* = \psi_z(\zeta)$. Then $\psi_z(M_{\rho_0})$ is given by $\eta_*^n = |\zeta'_*|^2 + h(\xi_*, x)$. Moreover,*

$$\begin{aligned} h(\xi_*, x) &= \sum_{|I|=1} \mathcal{R}_*^1 \hat{r}(\xi, x) \xi_*^I = \sum_{|I|=2} \mathcal{R}_*^2 \hat{r}(\xi, x) \xi_*^I, \\ -2r_z \cdot (\zeta - z) &= |\zeta'_*|^2 - i \xi_*^n + \sum_{|I|=2} \mathcal{R}_*^2 \hat{r}(\xi, x) \xi_*^I, \\ 2r_\zeta \cdot (\zeta - z) &= |\zeta'_*|^2 + i \xi_*^n + \sum_{|I|=2} (\mathcal{R}_*^2 \hat{r}(\xi, x) + \mathcal{R}_*^1 \partial \hat{r}(\xi, x)) \xi_*^I, \\ r_\zeta - r_z &= (\zeta'_*, 0) + \sum_{|I|=1} (\mathcal{R}_*^1 \hat{r}_{z^1}, \dots, \mathcal{R}_*^1 \hat{r}_{z^n})(\xi, x) \xi_*^I. \end{aligned}$$

We emphasize that $\mathcal{R}_*^i g$, $\mathcal{R}_*^i \partial \hat{r}$, defined by (6.5) and (6.11), might be different when they reoccur.

Remark 6.2. Notice that (6.2)-(6.8) are valid if we fix $\zeta \in M_{\rho_0}$ and vary $z \in M_{\rho_0}$. Therefore, the image of M_{ρ_0} under the map $z \rightarrow \psi_z(\zeta)$ is still given by (6.7) and (6.8), where z varies in M_{ρ_0} .

Here are immediate consequences of the above expansions:

$$\left| \frac{2r_\zeta \cdot (\zeta - z)}{N(\zeta, z)} - 1 \right| < \frac{1}{2}, \quad \left| \frac{2r_z \cdot (\zeta - z)}{N(\zeta, z)} + 1 \right| < \frac{1}{2}, \quad \frac{|N(z, \zeta)|}{|N(\zeta, z)|} < 4 \quad (6.14)$$

for $\zeta, z \in M_{\rho_0}$ with $\zeta \neq z$ and $\rho_0 \leq 3$.

Lemma 6.3. *Let M satisfy (2.2) and let $0 < \rho \leq \rho_0 \leq 3$. Set $d(\zeta, z) = |r_z \cdot (\zeta - z)|$. Then*

$$\begin{aligned} C^{-1}|\zeta - z|^2 &\leq d(\zeta, z) \leq C(d(\zeta, v) + d(v, z)), & \zeta, z, v \in M_{\rho_0}; \\ d(\zeta, z) &\geq C^{-1}\rho^2\sigma^2, & z \in M_{(1-\sigma)\rho}, \zeta \in \partial M_\rho, 0 < \sigma < 1. \end{aligned}$$

Proof. M is defined by $r = -y^n + |z'|^2 + \hat{r}(z', x^n) = 0$ and $\epsilon = \|\hat{r}\|_{\rho_0, 2} < C_0^{-1}$. Let ζ, z, v be in M_{ρ_0} . Set $\zeta_*' = \zeta' - z'$, $\zeta_*^n = -2ir_z \cdot (\zeta - z)$. Recall that D_{ρ_0} is convex.

(i). By definition,

$$N(\zeta, z) = |\zeta' - z'|^2 + 2i \operatorname{Im}(r_z \cdot (\zeta - z)) = |\zeta_*'|^2 + i\xi_*^n.$$

By Lemma 6.1, we have

$$\psi_z M_{\rho_0} : \eta_*^n = |\zeta_*'|^2 + \sum_{|I|=2} \mathcal{R}_*^2 \hat{r}(\xi, x) \xi_*^I.$$

On $D_{\rho_0} \times D_{\rho_0}$, $|\mathcal{R}_*^2 \hat{r}| \leq C\epsilon$, and $|\xi_*| < C$. Then $|\zeta_*^n|^2 = |\xi_*^n|^2 + |\eta_*^n|^2 \leq C'_0 |\xi_*^n|^2$. Therefore, $|\zeta^n - z^n|^2 = |\frac{i}{2r_z^n} \zeta_*^n - \frac{r_z'}{r_z^n} \cdot \zeta_*'|^2 \leq C|N(\zeta, z)|$. Also, $|\zeta' - z'|^2 \leq |N(\zeta, z)|$. We conclude that $|N(\zeta, z)| \geq C^{-1}|\zeta - z|^2$ for $\zeta, z \in M_{\rho_0}$. By (6.14) we have $d(\zeta, z) \geq |\zeta - z|^2/C$. Now

$$\begin{aligned} |r_\zeta \cdot (\zeta - z)| &\leq |(r_\zeta - r_v) \cdot (z - v)| + |r_v \cdot (z - v)| + |r_\zeta \cdot (v - \zeta)| \\ &\leq |r_\zeta - r_v|^2 + |z - v|^2 + d(z, v) + d(v, \zeta) \leq C(d(\zeta, v) + d(v, z)). \end{aligned}$$

Thus $d(\zeta, z) \leq C'(d(\zeta, v) + d(z, v))$.

(ii). By Lemma 5.1, $\operatorname{dist}(\partial D_{(1-\sigma)\rho}, \partial D_\rho) \geq C^{-1}\rho\sigma$. Then $d(\zeta, z) \geq C^{-1}(\rho\sigma)^2$ follows from $d(\zeta, z) \geq C^{-1}|\zeta - z|^2$. \square

7. Outline of C^a estimates

Let M satisfy (2.1)-(2.2) and let $0 < \rho \leq \rho_0 \leq 3$. Recall from (3.10)-(3.11) that

$$\begin{aligned} P'_0 \varphi(x) &= \sum_{|I|=q-1} \sum_{|J|=q} \sum_{1 \leq j \leq n} d\bar{z}^I \int_{D_\rho} A_I^{j\bar{J}}(\xi, x) \frac{\varphi_{\bar{J}}(\xi)(r_{\zeta^j} - r_{z^j})}{(N_0^{n-q} S_0^q)(\zeta, z)} dV(\xi), \\ P'_1 \varphi(x) &= \sum_{|I|=q-1} \sum_{|J|=q} \sum_{\alpha, \beta=1}^{n-1} \sum_{s=1}^{2n-1} d\bar{z}^I \int_{\partial D_\rho} \frac{B_{I_s}^{\alpha\beta\bar{J}}(\xi, x) \varphi_{\bar{J}}(\xi)(r_{\zeta^\alpha} - r_{z^\alpha}) r_{\zeta^\beta}}{(\zeta^n - z^n)(N_0^{n-q-1} S_0^q)(\zeta, z)} dV^s(\xi). \end{aligned}$$

Here $\zeta, z \in M_\rho$, and $A_I^{j\bar{J}}, B_{I_s}^{\alpha\beta\bar{J}}$ are polynomials in $(r_\zeta, r_{\bar{\zeta}}, r_{\zeta\bar{\zeta}}, r_{z'}, r_{z''}, r_{z''}^{-1}, r_{z\bar{z}})$.

We emphasize that norms are defined on $D_\rho = \pi(M_\rho)$ via coordinates x . Let us indicate how to obtain C^a estimates. We will give estimates on shrinking domains $M_{(1-\sigma)\rho}$. For $\zeta \in \partial M_\rho$ and $z \in M_{(1-\sigma)\rho}$, we obtain $\min\{|r_\zeta \cdot (\zeta - z)|, |r_z \cdot (\zeta - z)|\} \geq C^{-1}(\rho\sigma)^2$ by Lemma 6.3 and $|\zeta^n - z^n| \geq C^{-1}\rho^2\sigma$ by (4.2). This will allow us to estimate the C^a -norm of boundary integral $P'_1\varphi$ by passing derivatives over the integral sign and differentiating the kernels directly.

We now deal with the interior integrals $P'_0\varphi$. Using a partition of unity, we can find a smooth function $\chi = \chi_{\sigma,\rho}$, which is 1 on $D_{\rho(1-\sigma/2)}$ and zero off $D_{\rho(1-\sigma/4)}$, such that $\|\chi\|_{\rho,a} \leq C_a(\rho\sigma)^{-a}$. On D_ρ , decompose

$$\varphi_0 = \chi\varphi, \quad \varphi_1 = \varphi - \varphi_0, \quad P'_0\varphi = P'_0\varphi_0 + P'_0\varphi_1.$$

Now $P'_0\varphi_1$ can be estimated on $M_{(1-\sigma)\rho}$ by differentiating the kernels directly, since φ_1 is supported in $\overline{M_\rho} \setminus M_{(1-\frac{1}{2}\sigma)\rho}$. The only non-trivial integral is $P'_0\varphi_0$, for which φ_0 has compact support in $\overline{M_{(1-\frac{1}{4}\sigma)\rho}}$. To estimate the latter, we will apply the transformation ψ_z for the integral and then differentiate the new integral.

The estimate for $P'_0\varphi_0$ is the most technical part. We deal with this estimate first in Sections 8 and 9. The estimates for boundary and cutoff terms are in Section 10.

We now conclude this section with estimates of some integrals.

Lemma 7.1. *Let $n \geq 2$. Let a be a real number, and $J = (j_1, \dots, j_{2n-1})$ be a multiindex of non-negative integers. Let $\beta = (j_{2n-1} + |J| - 2a) + 2n - 1$ and $0 < \rho_1 \leq \rho_0 < \infty$. Then*

$$\int_{|z'| \leq \rho_1, |x^n| < \rho_0} \frac{|(z', \bar{z}', x^n)^J|}{\|z'\|^2 + |ix^n|^a} dV \leq \begin{cases} C\rho_1^{1+\beta}, & -1 < \beta < 2n - 3, \\ \tilde{C}\rho_1^{2n-2} \log(2 + \frac{\rho_0}{\rho_1}), & \beta \geq 2n - 3; \end{cases}$$

$$\int_{\rho_1 \leq |z'| \leq \rho_0, |x^n| < \rho_0} \frac{|(z', \bar{z}', x^n)^J|}{\|z'\|^2 + |ix^n|^a} dV \leq \begin{cases} C|\rho_1^{1+\beta} - \rho_0^{1+\beta}|, & \beta \neq 1, \beta < 2n - 3, \\ C \log(\rho_0/\rho_1), & \beta = -1, \\ \tilde{C}(\rho_0 - \rho_1), & \beta \geq 2n - 3. \end{cases}$$

Here C depends only on n and β , and \tilde{C} depends on ρ_0 too.

Proof. Using $|x^n| \leq |z'|^2 + |x^n|$ and $|z_j| \leq (|z'|^2 + |x^n|)^{1/2}$, we may assume, without changing β , that $J = 0$ and

$$b(z', x^n) = \frac{|(z', \bar{z}', x^n)^J|}{\|z'\|^2 + |ix^n|^a} = \frac{1}{(|z'|^2 + |x^n|)^a}.$$

Assume first that $\beta = 2n - 1 - 2a < 2n - 3$, i.e. $a > 1$. Using polar coordinates, we get

$$\int_{|z'| \leq \rho_1, |x^n| \leq \rho_0} b(z', x^n) dV \leq C \int_0^{\rho_1} dr \int_0^{\rho_0} \frac{r^{2n-3} dx^n}{(r^2 + |x^n|)^a} \leq C' \int_0^{\rho_1} r^\beta dr.$$

Also, $\int_{\rho_1 \leq |z'| \leq \rho_0, |x^n| < \rho_0} b(z', x^n) dV \leq C' \int_{\rho_1}^{\rho_0} r^\beta dr$. The estimates in this case follow.

Assume now that $\beta \geq 2n - 3$, i.e. $a \leq 1$. Reducing it to $a = 1$ via \tilde{C} , we get

$$\int_{|z'| \leq \rho_1, |x^n| \leq \rho_0} b(z', x^n) dV \leq \tilde{C} \int_0^{\rho_1} r^{2n-3} \log\left(1 + \frac{\rho_0}{r^2}\right) dr.$$

Also, $\int_{\rho_1 \leq |z'| \leq \rho_0, |x^n| < \rho_0} b(z', x^n) dV \leq \int_{\rho_1}^{\rho_0} \tilde{C} dr$. The estimates are obtained by a simple computation. \square

8. New kernels and two formulae for derivatives

In this section we express the kernels by using Lemma 6.1 and derive two formulae for derivatives of $P_0\varphi$, where φ has compact support in D_ρ . The first formula will be used for C^a estimates and the second is for $C^{k+\frac{1}{2}}$ estimates.

Recall that with $\zeta, z \in M_\rho$ the coefficients of $P'_0\varphi(x)$ are sums of

$$\mathcal{I}(x) = \int_{D_\rho} \varphi_{\mathcal{J}}(\xi) \frac{\partial_*^2 r(\xi, x)(r_{\zeta j} - r_{z j})}{(N_0^a S_0^b)(\zeta, z)} dV(\xi)$$

over $1 \leq j \leq n$ and $|J| = q$, where $a = n - q$ and $b = q$. Set $f(\xi, x) = \varphi_{\mathcal{J}}(\xi)$.

We have defined $\partial_*^{2+k}r$ in Section 2. Now, CHANGE NOTATION and let

$$\begin{aligned} \partial_*^2 r &= p\left(\xi, x, (1 + \hat{r}_{x^n} \hat{r}_{\xi^n})^{-1}, r_{z^n}^{-1}, (1 + \hat{r}_{x^n} \mathcal{R}_{(0',1)} \hat{r}(\xi, x))^{-1}, \partial \hat{r}(x), \partial \hat{r}(\xi), \mathcal{Q}(\xi, x)\right), \\ \mathcal{Q}(\xi, x) &= \left(\partial^2 \hat{r}(x), \partial^2 \hat{r}(\xi), \mathcal{R}_*^1 \partial \hat{r}(\xi, x), \mathcal{R}_*^1 \hat{r}(\xi, x), \mathcal{R}_*^2 \hat{r}(\xi, x)\right), \end{aligned}$$

where p is a polynomial. Again, $\mathcal{R}_*^1 \partial \hat{r}$, $\mathcal{R}_*^2 \hat{r}$, defined by (6.11) and (6.5), and p might be different when they reoccur; for instance, $(\mathcal{R}_*^1 r)^2$ may be the product of two different $\mathcal{R}_*^1 r$'s. Define

$$\begin{aligned} \partial_*^{2+k} r(\xi, x) &= \sum \partial_*^2 r \cdot \partial^{l_1} \mathcal{Q}_1(\xi, x) \cdots \partial^{l_j} \mathcal{Q}_j(\xi, x), \quad j \geq 0, \\ \partial_*^2 \hat{r}(\xi, x) &= \sum \partial_*^2 r \cdot \mathcal{Q}_1(\xi, x) \cdots \mathcal{Q}_j(\xi, x), \quad j \geq 1, \end{aligned}$$

where $\sum_{l=1}^j |l_l| \leq k$, $\mathcal{Q}_l \in \mathcal{Q}$ and both sums have finitely many terms. Hence, we have simple relations

$$\partial_*^{2+k} r \partial_*^{2+j} r = \partial_*^{2+k+j} r, \quad \partial^J \partial_*^{2+k} r = \partial_*^{2+k+|J|} r.$$

The chain rule takes the form

$$\begin{aligned} \partial_{\xi_*, x}^I \Psi^{-1} &= \partial_*^2 r \circ \Psi^{-1}, \quad |I| = 1, \\ \partial_{\xi_*, x}^J \{(f \partial_*^{2+k} r) \circ \Psi^{-1}\} &= \sum_{|L| \leq |J|} (\partial^L f \cdot \partial_*^{2+k+|J|-|L|} r) \circ \Psi^{-1}. \end{aligned} \tag{8.1}$$

New kernels. Set $(\xi_*, x) = \Psi(\xi, x)$ with $\zeta, z \in M_{\rho_0}$. Recall that $N_*(\xi_*) = |\xi_*'|^2 + i\xi_*^n$. By Lemma 6.1,

$$\begin{aligned} N_0(\zeta, z) &\equiv 2r_\zeta \cdot (\zeta - z) = N_*(\xi_*)\hat{T}_1(\xi_*, x), \\ S_0(\zeta, z) &\equiv -2r_z \cdot (\zeta - z) = \overline{N_*(\xi_*)}\hat{T}_2(\xi_*, x), \\ \hat{T}_1(\xi_*, x) &= T_1 \circ \Psi^{-1}(\xi_*, x) = 1 + \sum_{|J|=2} \partial_*^2 \hat{r} \circ \Psi^{-1}(\xi_*, x) N_*^{-1}(\xi_*) \xi_*^J, \end{aligned} \quad (8.2)$$

$$\begin{aligned} \hat{T}_2(\xi_*, x) &= T_2 \circ \Psi^{-1}(\xi_*, x) = 1 + \sum_{|J|=2} \partial_*^2 \hat{r} \circ \Psi^{-1}(\xi_*, x) \overline{N_*^{-1}(\xi_*)} \xi_*^J, \\ \partial_*^2 r(\zeta, z)(r_{\zeta j} - r_{zj}) &= \sum_{|I|=1} \partial_*^2 r \circ \Psi^{-1}(\xi_*, x) \xi_*^I. \end{aligned} \quad (8.3)$$

Note that $|\partial_*^2 \hat{r}| \leq C\epsilon$ and $|\hat{T}_j(\xi_*, x) - 1| < 1/2$ when $\xi_* \neq 0$. Thus, we obtain

$$\frac{\partial_*^2 r(\zeta, z)(r_{\zeta j} - r_{zj})}{(N_0^a S_0^b)(\zeta, z)} = \sum_{|I|=1} \left\{ \partial_*^2 r \circ \Psi^{-1} \cdot \hat{T}_1^{-a} \hat{T}_2^{-b} \right\}(\xi_*, x) \hat{k}_{ab}^I(\xi_*), \quad (8.4)$$

$$\hat{k}_{ab}^I(\xi_*) = \xi_*^I N_*^{-a}(\xi_*) \overline{N_*^{-b}(\xi_*)}, \quad a = n - q, \quad b = q. \quad (8.5)$$

First formula of derivatives of \mathcal{I} . Recall that $\Psi(\xi, x) = (\tilde{\psi}_x(\xi), x)$, $\xi_* = \tilde{\psi}_x(\xi)$ and $(\tilde{\psi}_x^* dV)(\xi_*) = (\partial_*^1 r) \circ \Psi^{-1}(\xi_*, x) dV(\xi_*)$. By (8.4) and $\partial_*^2 r \partial_*^1 r = \partial_*^2 r$, we obtain

$$\mathcal{I}(x) = \sum_{|I|=1} \int_{\tilde{\psi}_x(D_\rho)} \left\{ (f \partial_*^2 r) \circ \Psi^{-1} \cdot \hat{T}_1^{-a} \hat{T}_2^{-b} \right\}(\xi_*, x) \hat{k}_{ab}^I(\xi_*) dV(\xi_*), \quad (8.6)$$

where $a = n - q, b = q$. By Lemma 7.1 with $\beta \geq 0, \hat{k}_{ab}^I \in L_{\text{loc}}^1$. For each $x \in D_\rho$, the integrand has compact support in $\tilde{\psi}_x(D_\rho) \subset B_{9\rho}$. To compute $\partial^k \mathcal{I}(x)$, we extend the integrand of $\mathcal{I}(x)$ to be zero on $B_{9\rho} \setminus \tilde{\psi}_x(D_\rho)$. The integral is over the fixed domain $B_{9\rho}$. So we can interchange the integral sign with ∂_x . The derivatives of \mathcal{I} have the form

$$\begin{aligned} \partial^K \mathcal{I}(x) &= \sum_{j+k'+l+m=|K|} \sum_{|J|=j} \sum_{|K'|=k'} \sum_{|L|=l} \sum_{|I|=1} \\ &\int_{B_{9\rho}} \left\{ (\partial^L f \partial_*^{2+m} r) \circ \Psi^{-1} \cdot \partial_x^J \hat{T}_1^{-a} \cdot \partial_x^{K'} \hat{T}_2^{-b} \right\}(\xi_*, x) \cdot \hat{k}_{ab}^I(\xi_*) dV(\xi_*). \end{aligned} \quad (8.7)$$

By (8.2), the first-order x -derivatives of \hat{T}_1^{-a} have the form

$$\begin{aligned} \partial_x^I \hat{T}_1^{-a}(\xi_*, x) &= \sum_{|J'|=1} \sum_{|L'|=2} \{ \hat{T}_1^{-a-1} \partial_x^{J'} (\partial_*^2 r \circ \Psi^{-1}) \}(\xi_*, x) N_*^{-1}(\xi_*) \xi_*^{L'} \\ &= \sum_{|L'|=2} (\partial_*^3 r \circ \Psi^{-1} \hat{T}_1^{-a-1})(\xi_*, x) N_*^{-1}(\xi_*) \xi_*^{L'}. \end{aligned}$$

Take derivatives consecutively and use the product rule. We can write

$$\partial_x^J \hat{T}_1^{-a}(\xi_*, x) = \sum_{s \leq |J|} \sum_{|L'|=2s} \left\{ (\partial_*^{2+|J|} r) \circ \Psi^{-1} \cdot \hat{T}_1^{-a-s} \right\}(\xi_*, x) \frac{\xi_*^{L'}}{N_*^s(\xi_*)}. \quad (8.8)$$

Analogously, $\partial_x^{K'} \hat{T}_2^{-b} (|K'| = k')$ can be written as

$$\partial_x^{K'} \hat{T}_2^{-b}(\xi_*, x) = \sum_{t \leq |K'|} \sum_{|L''|=2t} \left\{ (\partial_*^{2+|K'|} r) \circ \Psi^{-1} \cdot \hat{T}_2^{-b-t} \right\}(\xi_*, x) \frac{\xi_*^{L''}}{N_*^t(\xi_*)}. \quad (8.9)$$

Let $I' = I + L' + L''$, $a' = a + s$, $b' = b + t$. We have

$$N_*^{-s}(\xi_*) \xi_*^{L'} \overline{N_*^{-t}(\xi_*)} \xi_*^{L''} \hat{k}_{ab}^{I'} = \hat{k}_{a'b'}^{I'}, \quad 2a' + 2b' - |I'| = 2n - 1, \quad a' + b' \leq n + |K|.$$

By (8.8)-(8.9), we get

$$\begin{aligned} \partial^K \mathcal{I}(x) &= \sum_{|L| \leq |K|} \sum_{a \leq a' \leq a + |K|} \sum_{b \leq b' \leq b + |K|} \sum_{2a' + 2b' - |I'| = 2n - 1} \quad (8.10) \\ &\int_{B_{\theta\rho}} \frac{(\partial^L f \partial_*^{2+|K|-|L|} r) \circ \Psi^{-1}(\xi_*, x)}{(\hat{T}_1^{-a'} \hat{T}_2^{-b'}) (\xi_*, x)} \hat{k}_{a'b'}^{I'}(\xi_*) dV(\xi_*). \end{aligned}$$

Recall that $f(\xi, x) = \varphi_{\bar{J}}(\xi)$ has compact support in D_ρ . Since $|I'| = 2a' + 2b' - 2n + 1$, Lemma 7.1 with $\beta \geq 0$ implies $\hat{k}_{a'b'}^{I'} \in L_{\text{loc}}^1$. By the dominated convergence theorem, we see that $\partial^K \mathcal{I}$ are continuous. Note that this also implies that if $\hat{r} \in C^{k+2}(\overline{D_{\rho_0}})$ and $\varphi \in C^k(\overline{D_{\rho_0}})$, then $P' \varphi \in C^k(D_{\rho_0})$.

Second formula of derivatives of \mathcal{I} . We return to the original coordinates by letting $(\xi_*, x) = \Psi(\xi, x)$ and $\zeta, z \in M_\rho$. So $dV(\xi_*) = \partial_*^1 r dV(\xi)$. Also

$$\begin{aligned} \hat{T}_1^{-a'} \circ \Psi(\xi, x) &= N^{a'}(\zeta, z) N_0^{-a'}(\zeta, z), \quad \hat{T}_2^{-b'} \circ \Psi(\xi, x) = \overline{N^{b'}(\zeta, z)} S_0^{-b'}(\zeta, z), \\ \hat{k}_{a'b'}^{I'} \circ \tilde{\psi}_x(\xi) &= \frac{(\text{Re}(\zeta' - z'), \text{Im}(\zeta' - z'), 2 \text{Im}(r_z \cdot (\zeta - z)))^{I'}}{N(\zeta, z)^{a'} \overline{N^{b'}(\zeta, z)}}. \end{aligned}$$

Multiply the same sides of the three identities and expand the last numerator. Then $\{\hat{T}_1^{-a'} \hat{T}_2^{-b'} \hat{k}_{a'b'}^{I'}\} \circ \Psi$ is a linear combination of

$$k_{a'b'}^{I''}(\xi, x) \stackrel{\text{def}}{=} \frac{(\zeta' - z', \overline{\zeta' - z'}, \text{Im}(r_z \cdot (\zeta - z)))^{I''}}{N_0^{a'}(\zeta, z) S_0^{b'}(\zeta, z)}.$$

Since $|I''| = |I'| = 2a' + 2b' + 1 - 2n$, then

$$|I''| - 2a' - 2b' - 1 + 2n = 0, \quad |I''| \leq 2|K| + 1, \quad a' + b' \leq n + |K|. \quad (8.11)$$

By (8.10), derivatives of $\mathcal{I}(x)$ have the form

$$\partial^K \mathcal{I}(x) = \sum_{|L| \leq |K|} \sum_{a \leq a' \leq a + |K|} \sum_{b \leq b' \leq b + |K|} \sum_{1 + 2a' + 2b' - 2n \leq |I''| \leq 2|K| + 1} \int_{D_\rho} \partial^L \varphi_{\bar{J}}(\xi) \partial_*^{2+|K|-|L|} r(\xi, x) \hat{k}_{a'b'}^{I''}(\xi, x) dV(\xi). \tag{8.12}$$

Here $\varphi_{\bar{J}}$ has compact support in D_ρ . Obviously, the $\partial^L \varphi_{\bar{J}}$ in (8.12) do not depend on x . This simple observation will be crucial for the $\frac{1}{2}$ -estimate.

The reader might want to acquaint with the counting scheme in Section 2 and Hölder inequalities on domains D_ρ in Appendix A; see Proposition A.5.

9. C^a -estimates, case of compact support

In this section, we derive the C^a -estimate for $P_0\varphi_0$ where

$$\varphi_0 = \chi\varphi, \quad \|\chi\|_{\rho, a} \leq C_a(\rho\sigma)^{-a} \tag{9.1}$$

and χ is supported in D_ρ . We also derive an estimate for $P_0\varphi$ when φ itself has compact support in D_ρ .

Proposition 9.1. *Let $k \geq 0$ be an integer and $0 \leq \alpha < 1$. Let $M: y^n = |z'|^2 + \hat{r}(z', x^n)$ satisfy (2.2) and $0 < \rho \leq \rho_0 \leq 3$. Let φ_0 be a tangential form as in (9.1). Then*

$$\|P'_0\varphi_0\|_{\rho, k+\alpha} \leq C_k \rho^{1-k-\alpha} \sigma^{-k-\alpha} (\|\varphi\|_{\rho, k+\alpha} + \|\varphi\|_{\rho, 0} \|\hat{r}\|_{\rho, k+2+\alpha}). \tag{9.2}$$

If φ has compact support in D_ρ and is tangential, then

$$\|P'_0\varphi\|_{\rho, k+\alpha} \leq C_k \rho^{1-k-\alpha} (\|\varphi\|_{\rho, k+\alpha} + \|\varphi\|_{\rho, 0} \|\hat{r}\|_{\rho, k+2+\alpha}). \tag{9.3}$$

The same estimate holds for Q'_0 .

Proof. Recall that by applying (8.10) to $f = \chi\varphi_{\bar{L}}$, k -th derivatives of a coefficient of $P'_0\varphi_0$ are sums of finitely many

$$\mathcal{I}_k(x) = \int_{B_{9\rho}} \frac{u \circ \Psi^{-1}(\xi_*, x)}{\hat{T}_1^{a'} \hat{T}_2^{b'}(\xi_*, x)} \hat{k}_{a'b'}^{I'}(\xi_*) dV(\xi_*).$$

Here $a = q, b = n - q, a \leq a' \leq a + k, b \leq b' \leq b + k$ and

$$u(\xi, x) \stackrel{\text{def}}{=} \partial^I \chi(\xi) \partial^J \varphi_{\bar{L}}(\xi) \partial_*^{2+|I|} r(\xi, x), \quad |I| = i, |J| = j, \quad i + j + |I| = k, \\ \hat{k}_{a'b'}^{I'}(\xi_*) = N^{-a'}(\xi_*) \overline{N^{-b'}(\xi_*)} \xi_*^{I'}, \quad 2a' + 2b' = |I'| + 2n - 1.$$

To obtain (9.2), we estimate the C^α -norm of \mathcal{I}_k .

By the definition of $\partial_*^{2+l}r$, we have

$$\|\partial_*^{2+l}r\|_{\rho,\alpha} \leq C \sum_{l_1+\dots+l_l \leq l} \|r\|_{2+l_1+\alpha} \cdots \|r\|_{2+l_l}.$$

Thus

$$\begin{aligned} & \|\partial^i \chi\|_{\rho,0} \|\partial^j \varphi_{\overline{L}}\|_{\rho,0} \|\partial_*^{2+l}r\|_{\rho,\alpha} \\ & \leq C(\rho\sigma)^{-i} \sum_{l_1+\dots+l_l \leq l} \|\varphi_{\overline{L}}\|_{\rho,j} \|r\|_{2+l_1+\alpha} \cdots \|r\|_{2+l_l} \\ & \leq C'(\rho\sigma)^{-i} \rho^{-l-j-\alpha} (\|\varphi_{\overline{L}}\|_{\rho,l+j+\alpha} + \|r\|_{2+l+j+\alpha} \|\varphi_{\overline{L}}\|_{\rho,0}). \end{aligned}$$

Here the last inequality is obtained by Proposition A.5. Also

$$\begin{aligned} & \|\partial^i \chi\|_{\rho,\alpha} \|\partial^j \varphi_{\overline{L}}\|_{\rho,0} \|\partial_*^{2+l}r\|_{\rho,0} \\ & \leq C(\rho\sigma)^{-i-\alpha} \rho^{-l-j} (\|\varphi_{\overline{L}}\|_{\rho,l+j} + \|r\|_{2+l+j} \|\varphi_{\overline{L}}\|_{\rho,0}), \\ & \|\partial^i \chi\|_{\rho,0} \|\partial^j \varphi_{\overline{L}}\|_{\rho,\alpha} \|\partial_*^{2+l}r\|_{\rho,0} \\ & \leq \frac{C}{(\rho\sigma)^i \rho^{l+j+\alpha}} (\|\varphi_{\overline{L}}\|_{\rho,l+j+\alpha} + \|r\|_{2+l+j+\alpha} \|\varphi_{\overline{L}}\|_{\rho,0}). \end{aligned}$$

Therefore,

$$\|u\|_{D_{\rho}^2,\alpha} \leq C_k \rho^{-k-\alpha} \sigma^{-k-\alpha} (\|\varphi\|_{\rho,k+\alpha} + \|r\|_{\rho,k+2+\alpha} \|\varphi\|_{\rho,0}), \quad (9.4)$$

$$\|u\|_{D_{\rho}^2,0} \|r\|_{\rho,2+\alpha} \leq C_k \rho^{-k-\alpha} \sigma^{-k} (\|\varphi\|_{\rho,k+\alpha} + \|r\|_{\rho,k+2+\alpha} \|\varphi\|_{\rho,0}). \quad (9.5)$$

By Lemma 5.2, we know that $W_{\rho} = \Psi(D_{\rho} \times D_{\rho}) \subset B_{9\rho} \times D_{\rho}$ and

$$|\Psi^{-1}(v) - \Psi^{-1}(u)| \leq C|v - u|, \quad u, v \in W_{\rho}. \quad (9.6)$$

Assume that $0 \leq \alpha < 1$. Fix $\xi_* \in B_{9\rho} \setminus \{0\}$ and $x_1, x_2 \in D_{\rho}$. Assume first that $\xi_j = \tilde{\psi}_{x_j}^{-1}(\xi_*)$ are in D_{ρ} for $j = 1, 2$. First, by (9.6)

$$|\xi_2 - \xi_1| \leq C|x_2 - x_1|.$$

Now by (8.2)-(8.3), we obtain $|\hat{T}_j(\xi_*, x)| \geq 1/4$ and

$$|\hat{T}_j(\xi_*, x_2) - \hat{T}_j(\xi_*, x_1)| \leq C|\partial_*^2 r(\xi_2, x_2) - \partial_*^2 r(\xi_1, x_1)| \leq \|r\|_{\rho,2+\alpha}|x_2 - x_1|^{\alpha}.$$

Thus

$$\begin{aligned} \Delta & = |\Delta(x_2) - \Delta(x_1)| \stackrel{\text{def}}{=} \left| \frac{u \circ \Psi^{-1}(\xi_*, x_2)}{\hat{T}_1^{a'} \hat{T}_2^{b'}(\xi_*, x_2)} - \frac{u \circ \Psi^{-1}(\xi_*, x_1)}{\hat{T}_1^{a'} \hat{T}_2^{b'}(\xi_*, x_1)} \right| \\ & \leq C\|u\|_{\rho,\alpha}|x_2 - x_1|^{\alpha} + C|u(\xi_2, x_2)((\partial_*^2 r)(\xi_2, x_2) - (\partial_*^2 r)(\xi_1, x_1))| \\ & \leq C(\|u\|_{\rho,\alpha} + \|u\|_{\rho,0} \|\hat{r}\|_{\rho,2+\alpha})|x_2 - x_1|^{\alpha}. \end{aligned}$$

By (9.4)-(9.5) we get

$$\Delta \leq C_k \rho^{-k-\alpha} \sigma^{-k-\alpha} (\|\varphi\|_{\rho, k+\alpha} + \|r\|_{\rho, k+2+\alpha} \|\varphi\|_{\rho, 0}) |x_2 - x_1|^\alpha. \quad (9.7)$$

The above holds trivially if ξ_1, ξ_2 are both not in D_ρ , in which case $\Delta = 0$. If $\xi_2 \in D_\rho$ and $\xi_1 \notin D_\rho$, we replace x_1 by a point x_3 in the line segment $[x_1, x_2]$, for which $\xi_3 = \tilde{\psi}_{x_3}^{-1}(\xi_*) \in \partial D_\rho$. Then $\Delta = |\Delta(x_2)| = |\Delta(x_2) - \Delta(x_3)|$ and (9.7) still holds. By Lemma 7.1 (with $\rho_1 = \rho_0 = 9\rho$ and $\beta \geq 0$), $\int_{B_{9\rho}} |\hat{k}'_{a', b'}| dV \leq C\rho$. Combining the above estimates yields (9.2).

The case that $\varphi_0 = \varphi$ is simpler, and it does not involve σ . So we can remove all powers of σ in (9.4)-(9.5), (9.7), and (9.2). The latter becomes (9.3). \square

We compute the $\mathcal{C}^{\frac{1}{2}}$ norm of $\partial^I \chi \partial^J \varphi_{\bar{L}} \partial_*^{2+I} r$ for a later use. Here it is crucial to avoid the Hölder $\frac{1}{2}$ -norm of $\partial^J \varphi_{\bar{L}}$.

Proposition 9.2. *Let $u(\xi, x) = \partial^I \chi(\xi) \partial^J \varphi_{\bar{L}}(\xi) \partial_*^{2+I} r(\xi, x)$ where χ has compact support in D_ρ and satisfies $\|\chi\|_{\rho, a} \leq C(\rho\sigma)^{-a}$. Let $|I| + |J| + l = k$. Then*

$$\|u(\xi, \cdot)\|_{\rho, \frac{1}{2}} \leq C_k (\rho\sigma)^{-k-\frac{1}{2}} (\|\varphi_{\bar{L}}\|_{\rho, k} \|r\|_{\rho, \frac{5}{2}} + \|\varphi_{\bar{L}}\|_{\rho, 0} \|\hat{r}\|_{\rho, k+\frac{5}{2}}). \quad (9.8)$$

Proof. Fix $\xi \in D_\rho$. The $\varphi_{\bar{L}}$ appearing in $u(\xi, x)$ depends only on ξ . Therefore, for $\|u(\xi, \cdot)\|_{\rho, 1/2}$, we only use the sup norm of $\partial^J \varphi_{\bar{L}}(\xi)$. Then

$$\begin{aligned} \|u(\xi, \cdot)\|_{\rho, 1/2} &\leq C_k ((\rho\sigma)^{-i-\frac{1}{2}} \|\varphi_{\bar{L}}\|_{\rho, j} \|r\|_{\rho, l+2} + (\rho\sigma)^{-i} \|\varphi_{\bar{L}}\|_{\rho, j} \|r\|_{\rho, l+\frac{5}{2}}) \\ &\leq C'_k (\rho\sigma)^{-i-\frac{1}{2}} \rho^{-j-l} (\|\varphi_{\bar{L}}\|_{\rho, j+l} + \|\varphi_{\bar{L}}\|_{\rho, 0} \|r\|_{\rho, 2+j+l}) \\ &\quad + C'_k (\rho\sigma)^{-i} \rho^{-j-l-\frac{1}{2}} (\|\varphi_{\bar{L}}\|_{\rho, 0} \|r\|_{\rho, j+l+\frac{5}{2}} + \|\varphi_{\bar{L}}\|_{\rho, j+l} \|r\|_{\rho, \frac{5}{2}}), \end{aligned}$$

where $|I| = i$, $|J| = j$, $|L| = l$ and the last two terms are obtained by Proposition A.5 in which we take $d_1 = 0$ and $d_2 = \frac{5}{2}$. Simplifying yields (9.8). \square

10. Boundary integrals, end of \mathcal{C}^a -estimates

In this section we will estimate the boundary integrals $P'_1 \varphi$ and cutoff term $P'_0 \varphi_1$, where φ_1 vanishes on $D_{(1-\frac{1}{2}\sigma)\rho}$. Estimates (10.14) and (10.15) below will be used again for the $\mathcal{C}^{k+\frac{1}{2}}$ estimate.

Recall (3.10)-(3.11) that for $\zeta, z \in M_\rho$

$$\begin{aligned} P'_0 \varphi(x) &= \sum_{|I|=q-1} \sum_{|J|=q} \sum_{1 \leq j \leq n} d\bar{z}^{\bar{I}} \int_{D_\rho} A_I^{j\bar{J}}(\xi, x) \frac{\varphi_{\bar{J}}(\xi)(r_{\zeta^j} - r_{z^j})}{(N_0^{n-q} S_0^q)(\zeta, z)} dV(\xi), \\ P'_1 \varphi(x) &= \sum_{|I|=q-1} \sum_{|J|=q} \sum_{\alpha, \beta=1}^{n-1} \sum_{s=1}^{2n-1} d\bar{z}^{\bar{I}} \int_{\partial D_\rho} \frac{B_I^{\alpha\beta\bar{J}}(\xi, x) \varphi_{\bar{J}}(\xi)(r_{\zeta^\alpha} - r_{z^\alpha}) r_{\zeta^\beta}}{(\zeta^n - z^n)(N_0^{n-q-1} S_0^q)(\zeta, z)} dV^s(\xi). \end{aligned}$$

Here $A_I^{j\bar{J}}$ and $B_{I\bar{S}}^{\alpha\beta\bar{J}}$ are polynomials in $(r_\zeta, r_{\bar{\zeta}}, r_{\zeta\bar{\zeta}}, r_{\bar{\zeta}'}, r_{\zeta\bar{\zeta}}^{-1}, r_{z\bar{z}})$. And $N_0(\zeta, z) = r_\zeta \cdot (\zeta - z)$, $S_0(\zeta, z) = r_z \cdot (\zeta - z)$. For $\zeta, z \in M_\rho$, set

$$k(\xi, x) \stackrel{\text{def}}{=} k_I^{j\bar{J}}(\xi, x) = A_I^{j\bar{J}}(\xi, x) \frac{(r_{\zeta^j} - r_{z^j})}{(N_0^{a_0} S_0^{b_0})(\zeta, z)}, \quad a_0 + b_0 = n, \quad b_0 = q,$$

$$l(\xi, x) \stackrel{\text{def}}{=} l_{I\bar{S}}^{\alpha\beta\bar{J}}(\xi, x) = B_{I\bar{S}}^{\alpha\beta\bar{J}}(\xi, x) \frac{r_{\zeta^\beta} (r_{\zeta^\alpha} - r_{z^\alpha})}{(\zeta^n - z^n)(N_0^{a_0-1} S_0^{b_0})(\zeta, z)}.$$

Recall that $\chi = \chi_{\sigma, \rho}$ is a smooth function, which is 1 on $D_{\rho(1-\sigma/2)}$ and zero off $D_{\rho(1-\sigma/4)}$, and $\|\chi\|_{\rho, a} \leq C_a(\rho\sigma)^{-a}$. On D_ρ , set

$$\varphi_0 = \chi\varphi, \quad \varphi_1 = \varphi - \varphi_0.$$

Assume that M satisfies (2.1)-(2.2). Thus $\|\hat{r}\|_{\rho_0, 2} < 1/C_0$. Assume that $0 < \rho \leq \rho_0 \leq 3$. Set $R = |z'|^2 + \hat{r}(x)$. Let π_s be the projection from ∂D_ρ into the subspace $\xi^s = 0$. Since \bar{D}_ρ is bounded and strictly convex, then π_s is a 2-to-1 map from ∂D_ρ onto $\pi_s(\partial D_\rho) = \pi(\bar{D}_\rho)$. Actually, π_s sends $\pi_s^{-1}(\partial(\pi_s(D_\rho)))$ one-to-one and onto $\partial(\pi_s(D_\rho))$. Let $\text{vol}(\pi_s D_\rho)$ be the volume of $\pi_s(D_\rho)$ calculated via the volume-form dV^s . Recall that D_ρ is contained in $B_{2\rho}$. If f is a continuous function on D_ρ , then

$$\left| \int_{\partial D_\rho} f(\xi) dV^s(\xi) \right| \leq 2 \text{vol}(\pi_s D_\rho) \|f\|_{\rho, 0} \leq 2 \text{vol}(B_{2\rho} \cap \mathbb{R}^{2n-2}) \|f\|_{\rho, 0} \quad (10.1)$$

$$\leq C\rho^{2n-2} \|f\|_{\rho, 0}.$$

Since the projection of ∂D_ρ in any coordinate hyperplane is contained in a ball of radius 2ρ , by the Fubini theorem one can verify that

$$\text{vol}(D_\rho \setminus D_{(1-\frac{1}{2}\sigma)\rho}) \leq (2n-1) \cdot C\rho\sigma \cdot \rho^{2n-2} \leq C'\rho^{2n-1}\sigma. \quad (10.2)$$

Let $k \geq 0$ be an integer and $0 \leq \alpha < 1$. Fix $\zeta \in M_\rho \setminus M_{(1-\frac{1}{2}\sigma)\rho}$ and vary $z \in M_{(1-\sigma)\rho}$. By Lemma 6.3, we have

$$|r_\zeta \cdot (\zeta - z)| \geq C^{-1}(\rho\sigma)^2, \quad |r_z \cdot (\zeta - z)| \geq C^{-1}(\rho\sigma)^2. \quad (10.3)$$

We also have $|r_{\zeta^j} - r_{z^j}| \leq C\|\zeta - z\| \leq C'|r_\zeta \cdot (\zeta - z)|^{1/2}$. Hence

$$|r_{\zeta^j} - r_{z^j}| |N_0|^{-a} |S_0|^{-b} \leq C|N_0|^{\frac{1}{2}-a-b} \leq C'(\rho\sigma)^{1-2a-2b} \quad (10.4)$$

if $a + b \geq 1/2$. Using $|f(x_2) - f(x_1)| \leq C\|f\|_{\rho, 1}|x_2 - x_1| \leq C\|f\|_{\rho, 1}(|x_2| + |x_1|)^{1-\alpha}|x_2 - x_1|^\alpha$, we get

$$\|f\|_{\rho, \alpha} \leq \|f\|_{\rho, 0} + C\|f\|_{\rho, 1}\rho^{1-\alpha}.$$

Therefore, for $1 \leq j \leq n$ and $0 < \alpha < 1$,

$$\|r_{\zeta j} - r_{z_j}(\cdot)\|_{\rho, \alpha} \leq C\rho^{1-\alpha}, \quad |r_{\zeta\beta}| \leq C\rho. \quad (10.5)$$

Also for $x_1, x_2 \in D_\rho$,

$$\begin{aligned} \left| \frac{1}{f(x_2)} - \frac{1}{f(x_1)} \right| &= \frac{|f(x_1) - f(x_2)|^{1-\alpha}}{|f(x_2)f(x_1)|} |f(x_1) - f(x_2)|^\alpha \\ &\leq C2^{1-\alpha} \|1/f\|_{\rho, 0}^{1+\alpha} \|f\|_{\rho, 1}^\alpha |x_2 - x_1|^\alpha. \end{aligned}$$

Combining it with Hölder ratio $|1/f^a|_{\rho, \alpha} \leq C_a \|1/f\|_{\rho, 0}^{a-1} |1/f|_{\rho, \alpha}$ for $a \geq 1$, we get

$$\|1/f^a\|_{\rho, \alpha} \leq \|1/f^a\|_{\rho, 0} + C_a \|1/f\|_{\rho, 0}^{a+\alpha} \|f\|_{\rho, 1}^\alpha, \quad a \geq 1. \quad (10.6)$$

Now, by (10.3)

$$\|N_0(\zeta, \cdot)^{-a}\|_{(1-\sigma)\rho, \alpha} + \|S_0(\zeta, \cdot)^{-a}\|_{(1-\sigma)\rho, \alpha} \leq C(\rho\sigma)^{-2(a+\alpha)}, \quad a \geq 1. \quad (10.7)$$

Note that $A_J^{\bar{J}}$ has the form $\partial_*^2 r$. By Proposition A.5, we have

$$\|\partial_*^2 r(\xi, \cdot)\|_{\rho, a} \cdot \|\partial_*^2 r(\xi, \cdot)\|_{\rho, b} \leq C_{a,b} \rho^{-a-b} \|r\|_{\rho, 2+a+b}. \quad (10.8)$$

We have

$$\partial_x^K k(\xi, x) = \sum_{a+b+c+|L|=|K|} \sum_{|L|=0,1} \frac{\partial_*^{2+c} r(\xi, x) \partial_x^L (r_{\zeta j} - r_{z_j})}{(N_0^{a_0+a} S_0^{b_0+b})(\zeta, z)},$$

where $a_0 + b_0 = n$. Fix ξ and vary x . For the summand, we estimate the C^α -norms of two terms in the numerator by (10.3)-(10.5), and use (10.7) for the reciprocals of two terms in the denominator. Set $|K| = k$ and $|L| = d$. Recall that $d = 0$ or 1 . We get

$$\begin{aligned} &\left\| \frac{\partial_*^{2+c} r(\xi, \cdot) \partial_x^L (r_{\zeta j} - r_{z_j}(\cdot))}{(N_0^{a_0+a} S_0^{b_0+b})(\zeta, \cdot)} \right\|_{(1-\sigma)\rho, \alpha} \\ &\leq \frac{C}{(\rho\sigma)^{2(a+a_0+b+b_0)}} \left\{ (\rho\sigma)^{1-d} \rho^{-c-\alpha} \|r\|_{\rho, 2+c+\alpha} \right. \\ &\quad \left. + \rho^{(1-d)(1-\alpha)} \rho^{-c-d\alpha} \|r\|_{\rho, 2+c+d\alpha} + 2(\rho\sigma)^{1-d} (\rho\sigma)^{-2\alpha} \rho^{-c} \|r\|_{\rho, 2+c} \right\}. \end{aligned}$$

The worst term in terms of powers of ρ, σ occurs when $a + b = k, c = d = 0$. We see that

$$\|k(\xi, \cdot)\|_{(1-\sigma)\rho, k+\alpha} \leq C_k (\rho\sigma)^{-s_1} \|r\|_{\rho, k+2+\alpha}, \quad \xi \in D_\rho \setminus D_{(1-\frac{1}{2}\sigma)\rho} \quad (10.9)$$

with $s_1 = 2n - 1 + 2k + 2\alpha$.

To estimate the boundary term, by (4.2) we have

$$|\zeta^n - z^n| \geq C^{-1} \rho^2 \sigma \quad (10.10)$$

for $z \in M_{(1-\sigma)\rho}$ and $\zeta \in \partial M_\rho$. Using (10.6), we get for $\zeta \in \partial M_\rho$

$$\|(\zeta^n - z^n(\cdot))^{-a}\|_{(1-\sigma)\rho, \alpha} \leq C(\rho^2 \sigma)^{-a-\alpha}, \quad a \geq 1, \quad 0 \leq \alpha \leq 1. \quad (10.11)$$

Note that $B_{\bar{I}S}^{\alpha\beta\bar{J}}$ has the form $\partial_*^2 r$. We have

$$\partial_x^K l(\xi, x) = \sum_{a+b+d+c=|K|} \sum_{c_1+|L|=c} \sum_{|L|=0,1} \frac{\partial_*^{2+c_1} r \cdot r_{\zeta\beta} \cdot \partial_x^L (r_{\zeta j} - r_{zj})}{(\zeta^n - z^n)^{1+d} (N_0^{a_0-1+a} S_0^{b_0+b})(\zeta, z)}.$$

Set $|K| = k$ and $|L| = c_2$. We now estimate the C^α -norm in the x variables. Fix $\zeta \in \partial D_\rho$. Using (10.3)-(10.5), (10.8) for three terms in the numerator and (10.7), (10.10)-(10.11) for the reciprocals of three terms in the denominator, we obtain

$$\begin{aligned} \|l(\xi, \cdot)\|_{(1-\sigma)\rho, k+\alpha} &\leq C_k \sum_{a+b+d+c_1+c_2=k} \sum_{c_2=0,1} (\rho\sigma)^{-2(a_0+a+b_0+b+d)} \sigma^{1+d} \rho. \\ &\cdot \left\{ (\rho\sigma)^{1-c_2} \rho^{-c_1-\alpha} \|r\|_{\rho, 2+c_1+\alpha} + \rho^{(1-c_2)(1-\alpha)} \rho^{-c_1-c_2\alpha} \|r\|_{\rho, 2+c_1+c_2\alpha} \right. \\ &\quad \left. + \rho^{-c_1} (\rho\sigma)^{1-c_2} ((\rho^2\sigma)^{-\alpha} + 2(\rho\sigma)^{-2\alpha}) \|r\|_{\rho, 2+c_1} \right\}. \end{aligned} \quad (10.12)$$

The worst term in terms of powers of ρ, σ occurs when $c_1 = c_2 = d = 0, a+b = k$. This shows that for each $\zeta \in \bar{M}_\rho \setminus M_{(1-\frac{1}{2})\rho}$, we have

$$\|l(\xi, \cdot)\|_{(1-\sigma)\rho, k+\alpha} \leq C_k (\rho\sigma)^{-s_2} \|r\|_{\rho, k+2+\alpha} \quad (10.13)$$

with $s_2 = 2(n+k-1+\alpha)$.

By (10.1) and (10.13), we estimate the boundary term by

$$\|P'_1 \varphi\|_{(1-\sigma)\rho, k+\alpha} \leq C_k \rho^{2n-2} (\rho\sigma)^{-s_2} \|r\|_{\rho, k+\alpha+2} \|\varphi\|_{\rho, 0}. \quad (10.14)$$

Estimating the cutoff term by (10.9) and (10.2), we obtain

$$\|P'_0 \varphi_1\|_{(1-\sigma)\rho, k+\alpha} \leq C_a \rho^{2n-1-s_1} \sigma^{1-s_1} \|r\|_{\rho, k+2+\alpha} \|\varphi\|_{\rho, 0}. \quad (10.15)$$

Define $s \stackrel{\text{def}}{=} \max\{s_1-1, s_2, k+\alpha\}$ and $s_* \stackrel{\text{def}}{=} \max\{s_1-2n+1, s_2-2n+2, k+\alpha-1\}$. Thus for $a = k + \alpha$, we get

$$s = 2(a+n-1), \quad s_* = 2a. \quad (10.16)$$

Combining (9.2), (10.14)-(10.15), we get the following.

Proposition 10.1. *Let $n \geq 4$, and let $a \geq 0$ be a real number. Let $M: y^n = |z'|^2 + \hat{r}(z', x^n)$ satisfy (2.2). Let P' be either of P', Q' in the homotopy formula $\varphi = \bar{\partial}_M P' \varphi + Q' \bar{\partial}_M \varphi$ on M_ρ . Assume that $0 < \rho \leq \rho_0 \leq 3$. Then for a tangential form φ*

$$\|P'\varphi\|_{D_{(1-\sigma)\rho}, a} \leq C_a \rho^{-s_*} \sigma^{-s} (\|\varphi\|_{D_{\rho, a}} + \|\varphi\|_{D_{\rho, 0}} \|\hat{r}\|_{D_{\rho, a+2}}),$$

where $0 < \sigma < 1$, s, s_* are given by (10.16).

11. Reduction of $C^{k+\frac{1}{2}}$ -estimates to $C^{\frac{1}{2}}$ -estimate. Summary

We want to prove the following $C^{k+\frac{1}{2}}$ estimates.

Proposition 11.1. *Let $n \geq 4$, and let $k \geq 0$ be an integer. Let $M: y^n = |z'|^2 + \hat{r}(z', x^n)$ satisfy (2.2). Let P' be one of P', Q' in the homotopy formula $\varphi = \bar{\partial}_M P' \varphi + Q' \bar{\partial}_M \varphi$ on M_ρ . Then for $0 < \rho \leq \rho_0 \leq 3$, $0 < \sigma < 1$, and a tangential $(0, q)$ form φ*

$$\begin{aligned} \|P'\varphi\|_{D_{(1-\sigma)\rho}, k+\frac{1}{2}} &\leq \frac{C_k}{\rho^{2k+1} \sigma^{2n+2k-1}} (\|r\|_{D_{\rho, \frac{5}{2}}} \|\varphi\|_{D_{\rho, k}} + \|\hat{r}\|_{D_{\rho, k+\frac{5}{2}}} \|\varphi\|_{D_{\rho, 0}}), \\ \|P\varphi\|_{D_{(1-\sigma)\rho}, \frac{1}{2}} &\leq C\rho^{-1} \sigma^{1-2n} \|\varphi\|_{D_{\rho, 0}}, \quad q = 1. \end{aligned}$$

Proof. Fix a positive integer k . To estimate the $C^{k+\frac{1}{2}}$ -norm of $P'\varphi = (P'_0 + P'_1)\varphi$, we first recall estimates (10.14) and (10.15) for the boundary and cutoff terms

$$\|(P'_0\varphi_1, P'_1\varphi)\|_{(1-\sigma)\rho, k+\frac{1}{2}} \leq C_a \rho^{-s_*} \sigma^{-s} \|r\|_{\rho, k+\frac{5}{2}} \|\varphi\|_{\rho, 0}. \tag{11.1}$$

Here $s = 2n + 2k - 1$ and $s_* = 2k + 1$ are computed by (10.16) for $a = k + \frac{1}{2}$. It remains to estimate the $C^{k+\frac{1}{2}}$ -norm of $P'_0\varphi_1$, where $\varphi_1 = \chi\varphi$ and χ has compact support in D_ρ and $\|\chi\|_{\rho, a} \leq C_a(\rho\sigma)^{-a}$. The proof will be completed later. For the rest of proof, we reduce it to the special case of $k = 0$.

The second formula (8.12) says that the coefficients of $\partial^k P'_0\varphi_0$ are sums of

$$\mathcal{K}u(x) = \int_{D_\rho} u(\xi, x) k(\xi, x) dV \tag{11.2}$$

with functions $u(\xi, x)$ and kernels $k(\xi, x)$ of the form

$$u(\xi, x) = \partial^E \chi(\xi) \partial^F \varphi_{\bar{J}}(\xi) \partial_*^{2+l} r(\xi, x), \quad |E| + |F| + l = k, \tag{11.3}$$

$$k(\xi, x) \stackrel{\text{def}}{=} k_{ab}^l(\zeta, z) = \frac{(\zeta' - z', \bar{\zeta}' - \bar{z}', \text{Im}(r_z \cdot (\zeta - z)))^l}{(r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b}, \quad \zeta, z \in M_\rho. \tag{11.4}$$

Moreover, by (8.11), the non-negative integers $a, b, I = (i_1, \dots, i_{2n-1})$ satisfy

$$|I| - 2a - 2b \geq 1 - 2n, \quad |I| \leq 2k + 1, \quad a + b \leq n + k. \quad (11.5)$$

By (9.8) in Proposition 9.2, we have

$$\|u(\xi, \cdot)\|_{\rho, \frac{1}{2}} \leq Ck(\rho\sigma)^{-k-\frac{1}{2}} (\|\varphi\|_{\rho, k} \|r\|_{\rho, \frac{5}{2}} + \|\varphi\|_{\rho, 0} \|\hat{r}\|_{\rho, k+\frac{5}{2}}). \quad (11.6)$$

In Section 12 we will prove that if $u \in L^\infty(D_\rho \times D_\rho)$ then

$$\|\mathcal{K}u\|_{D_\rho, 1/2} \leq C \sup_{\xi \in D_\rho} \|u(\xi, \cdot)\|_{D_\rho, 1/2}. \quad (11.7)$$

Combining it with (11.1) and (11.6) yields the first estimate in the proposition.

We now consider the case that φ is a tangential $(0, 1)$ -form. We return to (3.10)-(3.11) and look at a special property of the kernels of $P'\varphi$. Recall that in this case

$$P'_0\varphi(x) = \sum_{1 \leq \gamma < n} \sum_{1 \leq j \leq n} \int_{D_\rho} A^{j\bar{\gamma}}(\xi, x) \frac{\varphi_{\bar{\gamma}}(\xi)(r_{\zeta j} - r_{z j})}{(N_0^{n-1} S_0)(\zeta, z)} dV(\xi),$$

$$P'_1\varphi(x) = \sum_{1 \leq \alpha, \beta, \gamma < n} \sum_{1 \leq s < 2n} \int_{\partial D_\rho} \frac{B_s^{\alpha\beta\bar{\gamma}}(\xi, x) \varphi_{\bar{\gamma}}(\xi)(r_{\zeta^\alpha} - r_{z^\alpha})r_{\zeta^\beta}}{(\zeta^n - z^n)(N_0^{n-2} S_0)(\zeta, z)} dV^s(\xi).$$

Here $\zeta, z \in M_\rho$. Also, $A^{j\bar{\gamma}}$ and $B_s^{\alpha\beta\bar{\gamma}}$ are polynomials in $r_\zeta, r_{\bar{\zeta}}, r_{\zeta\bar{\zeta}}$. In particular, they are independent of z . Moreover, in the kernels there are only the first-order derivatives r_{zj} in the z variable. Then all norms of r in (10.12)-(10.14), in which $k = 0$, can be replaced by $\|r\|_{\rho, 2} < C$ and the estimate (10.14) for the boundary term becomes

$$\|P'_1\varphi\|_{(1-\sigma)\rho, \alpha} \leq C\rho^{-2\alpha}\sigma^{2-2n-2\alpha}\|\varphi\|_{\rho, 0}, \quad 0 \leq \alpha \leq 1.$$

Absorb $A_I^{j\bar{\gamma}}(\xi)$ into $\varphi_{\bar{\gamma}}(\xi)$. With $\zeta, z \in M_\rho$ the kernels of interior integral $P'_0\varphi$ have the form

$$k(\xi, x) = \frac{r_{\zeta j} - r_{z j}}{(r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b}, \quad a + b = n. \quad (11.8)$$

We will show that (11.7) holds for this new kernel. Applying it to

$$u(\xi, x) = A^{j\bar{\gamma}}(\xi)\varphi_{\bar{\gamma}}(\xi), \quad \zeta \in M$$

we obtain $\|P'_0\varphi_{0,1}\|_{\rho, 1/2} \leq C\|\varphi_{0,1}\|_{\rho, 0}$. This shows the second estimate of the proposition.

The proof of Proposition 11.1 is thus complete, by assuming (11.7) in which $\mathcal{K}u$, given by (11.2), has a kernel of the form (11.4)-(11.5) or (11.8). \square

We want to unify the two kernels (11.4), which satisfies (11.5), and (11.8). For (11.4), we write

$$\text{Im}(r_z \cdot (\zeta - z)) = \frac{1}{2i} r_z \cdot (\zeta - z) - \frac{1}{2i} \overline{r_z \cdot (\zeta - z)}.$$

Then kernel (11.4) is a linear combination of

$$k(\xi, x) \stackrel{\text{def}}{=} \frac{(\zeta - z, \overline{\zeta - z}, r_\zeta - r_z)^I}{(r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b (r_\zeta \cdot (\zeta - z))^c (r_z \cdot (\zeta - z))^d}, \quad (11.9)$$

where $\zeta, z \in M$, a, b, c, d are now possibly negative integers, I is a $(3n)$ -tuple of nonnegative integers, and

$$|I| - 2(a+b+c+d) \geq 1 - 2n, \quad |I| \leq 2k + 1, \quad |a| + |b| + |c| + |d| \leq n + 3k + 2. \quad (11.10)$$

Indeed, the $I = (i_1, \dots, i_{2n-1})$ in (11.5) is a multi-index of nonnegative integers. Hence, (11.5) and $i_{2n-1} \geq 0$ implies (11.10). Obviously, (11.8) is of the form (11.9). In Section 12, for the new kernel (11.9) with the condition (11.10) we will prove (11.7), *i.e.*

$$\left\| \int_{D_\rho} u(\xi, \cdot) k(\xi, \cdot) dV(\xi) \right\|_{D_\rho, 1/2} \leq C \sup_{\xi \in D_\rho} \|u(\xi, \cdot)\|_{D_\rho, 1/2}, \quad (11.11)$$

where $u \in L^\infty(D_\rho \times D_\rho)$.

Summary. We would like to summarize our observations on $\mathcal{C}^{k+\frac{1}{2}}$ -estimates to explain how the general estimate (11.11) gives us an actual $\frac{1}{2}$ -gain in special cases.

a) $k \geq 1$ and $q \geq 1$. In our applications of (11.11) to the $\mathcal{C}^{k+\frac{1}{2}}$ -estimates, $u(\xi, x)$, arising from the k -th order derivatives of $P'_0\varphi$ after applying the approximate Heisenberg transformation, is of the form $\partial^{k-j} \varphi_{\mathcal{J}}(\xi) \partial_*^{j+2} r(\xi, x)$ where φ has compact support in D_ρ ; see the second formula of the derivatives in Section 8. In particular, the $\mathcal{C}^{\frac{1}{2}}$ -norm of $u(\xi, x)$ in x variables involves only $\|\varphi\|_{\rho, k}$ (and $\|r\|_{\rho, k+\frac{5}{2}}$). Thus the estimate for $\|P'_0\varphi\|_{\rho, k+\frac{1}{2}}$ gains $\frac{1}{2}$ in Hölder exponent from $\|\varphi\|_{\rho, k}$ at the expense of two extra derivatives in $\|r\|_{\rho, k+\frac{5}{2}}$.

b) $k = 0$ and $q = 1$. Apply (11.11) to $(0, 1)$ forms. In this case, we will not need to differentiate the kernels and apply cutoff. Both simplify the estimate considerably. However, it is crucial that when (11.11) is applied, $u(\xi, x)$ has the form $A^{j\bar{j}}(\xi) \varphi_{\bar{j}}(\xi)$; in particular, it is *independent* of x . Therefore, the $\frac{1}{2}$ -estimate of u in the x -variables and hence $\frac{1}{2}$ -estimate of $P'_0\varphi$ only require $r \in \mathcal{C}^2$.

c) While the $\frac{1}{2}$ -estimate of $P'_0\varphi$ is on the original domain (Proposition 12.1 below), the estimate for the boundary term $P'_1\varphi$, which only requires $r \in \mathcal{C}^2$ for the same reason as in b), is on shrinking domains.

12. $\frac{1}{2}$ -estimate

We will derive the $\frac{1}{2}$ -estimate, by modifying a standard argument for Hölder estimates. This will complete the proof of Proposition 11.1 via (11.6) and (11.11). We restate the latter as the following.

Proposition 12.1. *Let M be a real hypersurface of class C^2 and satisfy (2.2). Assume that $0 < \rho \leq \rho_0 \leq 3$. Let a, b, c, d be (possibly negative) integers and I be a $(3n)$ -tuple of nonnegative integers. Suppose that $|I| - 2(a + b + c + d) \geq 1 - 2n$. Let*

$$k(\xi, x) = \frac{(\zeta - z, \overline{\zeta - z}, r_\zeta - r_z)^I}{(r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b (r_\zeta \cdot (\zeta - z))^c (r_z \cdot (\zeta - z))^d}, \quad \zeta, z \in M,$$

$$\mathcal{K}u(x) = \int_{M_\rho} u(\xi, x) k(\xi, x) dV.$$

Then for $u \in L^\infty(D_\rho \times D_\rho)$,

$$\|\mathcal{K}u\|_{D_\rho, \frac{1}{2}} \leq C \left(\rho \sup_{\xi \in D_\rho} \|u(\xi, \cdot)\|_{D_\rho, \frac{1}{2}} + \sup_{\xi \in D_\rho} \|u(\xi, \cdot)\|_{D_\rho, 0} \right).$$

Proof. Recall that D_ρ is convex. It will be convenient to regard $\mathcal{K}u$ as a function on M . The $\frac{1}{2}$ -Hölder ratios on M_ρ and πM_ρ are equivalent, since $|\pi(z_2 - z_1)| \geq |z_2 - z_1|/C$ for $z_1, z_2 \in M_\rho$. By abuse of notation, we write $u(\xi, x)$ as $u(\zeta, z)$ with $\zeta, z \in M$ and apply the same change of notation to $k(\xi, x), \mathcal{K}u(x)$.

Fix z_1, z_2 in M_ρ . We have

$$\begin{aligned} \mathcal{K}u(z_2) - \mathcal{K}u(z_1) &= \int_{M_\rho} u(\zeta, z_2) (k(\zeta, z_2) - k(\zeta, z_1)) dV \\ &\quad + \int_{M_\rho} (u(\zeta, z_2) - u(\zeta, z_1)) k(\zeta, z_1) dV = \mathcal{I} + \mathcal{I}'. \end{aligned}$$

To estimate \mathcal{I}' , we need to estimate $\int_{M_\rho} |k(\xi, z_1)| dV(\xi)$. So we apply the approximate Heisenberg transformation $\zeta_* = \psi_{z_1}(\zeta)$. (However, we want to refrain from use of Taylor remainder expansions, such as (8.4)-(8.5), for the kernel. This avoids the requirement of $r \in C^{5/2}$.) Let $\zeta \in M_\rho$. Using the fundamental theorem of calculus, we get $\eta^n - y_1^n = A(\xi, x_1) \cdot (\xi - x_1)$ with $|A| < C$. By (6.4), we have $\xi^n - x_1^n = p(\xi, x_1) \cdot \xi_*$ with $|p| < C$. Therefore,

$$|\zeta - z_1| \leq C|\xi_*|.$$

By Lemma 6.1, we have

$$|r_\zeta - r_{z_1}| \leq C|\xi_*|, \quad 1/C \leq \frac{|r_{z_1} \cdot (\zeta - z_1)|}{\|\zeta_*'\|^2 + i\xi_*^n} \leq C, \quad 1/C \leq \frac{|r_\zeta \cdot (\zeta - z_1)|}{\|\zeta_*'\|^2 + i\xi_*^n} \leq C.$$

Therefore, for $\zeta \in M_\rho$

$$|k(\zeta, z_1)| \leq \frac{C|\xi_*^I|^I}{\|\zeta'_*\|^2 + i|\xi_*^n|^{a+b+c+d}} \leq \sum_{|J|=|I|} \frac{C|\xi_*^J|}{\|\zeta'_*\|^2 + i|\xi_*^n|^{a+b+c+d}}.$$

By Lemma 5.2, $\pi(\psi_{z_1}(M_\rho)) \subset B_{9\rho}$. Applying Lemma 7.1 with $\rho_1 = \rho_0 = 9\rho$ and $\beta \geq 0$, we get

$$\int_{M_\rho} |k(\zeta, z_1)| dV \leq \sum_{|J|=|I|} \int_{B_{9\rho}} \frac{C|\xi_*^J|}{|\xi_*^n + i|\zeta'_*|^2|^{a+b+c+d}} dV \leq C'\rho. \tag{12.1}$$

Since $u(\zeta, z)$ is of class $\mathcal{C}^{1/2}$ in the z -variable, one gets

$$|\mathcal{I}| \leq C|z_2 - z_1|^{1/2} \rho \sup_{\zeta \in D_\rho} \|u(\zeta, \cdot)\|_{\rho, 1/2}.$$

To estimate the integral \mathcal{I}' , it suffices to show that

$$\int_{M_\rho} |k(\zeta, z_2) - k(\zeta, z_1)| dV \leq C\delta^{1/2}, \quad \delta = |z_2 - z_1|.$$

As mentioned in the introduction we decompose M_ρ into a cylinder and its complement. Consider the cylinder $M_\rho \cap \{|\zeta' - z'_1| < \rho_1\}$, where $\rho_1 = \min(9\rho, C_*|z_2 - z_1|^{\frac{1}{2}})$ with $C_* > 1$ to be determined. Notice that the radius of cylinder is about $|z_2 - z_1|^{1/2}$, which is much large than $|z_2 - z_1|$, the Euclidean distance between z_1, z_2 .

We have

$$\begin{aligned} \int_{M_\rho} |k(\zeta, z_2) - k(\zeta, z_1)| dV &\leq \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3, \\ \mathcal{I}_1 &= \int_{M_\rho \cap \{|\zeta' - z'_1| < \rho_1\}} |k(\zeta, z_1)| dV, \quad \mathcal{I}_2 = \int_{M_\rho \cap \{|\zeta' - z'_1| < \rho_1\}} |k(\zeta, z_2)| dV, \\ \mathcal{I}_3 &= \int_{M_\rho \cap \{|\zeta' - z'_1| \geq \rho_1\}} |k(\zeta, z_2) - k(\zeta, z_1)| dV. \end{aligned}$$

By an analogy of (12.1), we have

$$\mathcal{I}_1 \leq \sum_{|J|=|I|} C \int_{|\zeta'_*| < \rho_1, |\zeta_*| < 9\rho} \frac{|\xi_*^J|}{|\xi_*^n + i|\zeta'_*|^2|^{a+b+c+d}} dV.$$

By Lemma 7.1 with $\beta \geq 0$ and $\rho_0 = 9\rho$ and $\rho_1 = \min(9\rho, C_*|z_2 - z_1|^{\frac{1}{2}})$, we get $\mathcal{I}_1 \leq C\rho_1 \leq CC_*|z_2 - z_1|^{1/2}$. Note that

$$M_\rho \cap \{|\zeta' - z'_1| < \rho_1\} \subset M_\rho \cap \{|\zeta' - z'_2| < \rho_1 + |z_2 - z_1|\}.$$

Applying the estimate for \mathcal{I}_1 , we get $\mathcal{I}_2 \leq C(\rho_1 + |z_2 - z_1|) \leq C'C_*|z_2 - z_1|^{1/2}$.

With $C_* > 1$ to be determined, we have $\mathcal{I}_3 = 0$ if $\rho_1 = 9\rho$. Therefore, we may assume that $\rho_1 = C_*|z_2 - z_1|^{\frac{1}{2}}$.

To estimate \mathcal{I}_3 , we ignore any non-isotropic distance and connect (z'_1, x_1^n) and (z'_2, x_2^n) by a line segment in the convex domain D_ρ . Let $z(t)$ be the graph of the line segment in M . Let ζ be a point of M_ρ which is not in the cylinder. Then

$$|\zeta'_*| = |\zeta' - z'_1| \geq \rho_1 = C_*|z_2 - z_1|^{1/2}.$$

We have

$$\begin{aligned} |r_{z(t)} \cdot (\zeta - z(t)) - r_{z_1} \cdot (\zeta - z_1)| &\leq C_0|z_2 - z_1|, \\ |r_{z_1} \cdot (\zeta - z_1)| &\geq C_1|\zeta - z_1|^2 \geq C_1|\zeta' - z'_1|^2 \geq C_1C_*^2|z_2 - z_1|, \end{aligned}$$

where the second inequality comes from Lemma 6.3. We now fix $C_* > 1$ such that $C_1C_*^2 > 2C_0$. (As remarked earlier, if $C_*|z_2 - z_1|^{\frac{1}{2}} > 9\rho$, we already have $\mathcal{I}_3 = 0$.) We obtain

$$|r_{z_1} \cdot (\zeta - z_1)|/2 \leq |r_{z(t)} \cdot (\zeta - z(t))| \leq 2|r_{z_1} \cdot (\zeta - z_1)|.$$

Recall that $\zeta_* = \psi_{z_1}(\zeta)$ and $C^{-1}||\zeta'_*|^2 + i\xi_*^n| \leq |r_{z_1} \cdot (\zeta - z_1)| \leq C||\zeta'_*|^2 + i\xi_*^n|$. Using $|r_{z(t)} \cdot (\zeta - z(t))|/C \leq |r_\zeta \cdot (\zeta - z(t))| \leq C|r_{z(t)} \cdot (\zeta - z(t))|$, we get

$$C^{-1}||\zeta'_*|^2 + i\xi_*^n| \leq |r_\zeta \cdot (\zeta - z(t))| \leq C||\zeta'_*|^2 + i\xi_*^n|, \quad (12.2)$$

$$C^{-1}||\zeta'_*|^2 + i\xi_*^n| \leq |r_{z(t)} \cdot (\zeta - z(t))| \leq C||\zeta'_*|^2 + i\xi_*^n|. \quad (12.3)$$

Write $k(\zeta, z) = \frac{p(\zeta, z)}{q(\zeta, z)}$ with $p(\zeta, z) = (\zeta' - z', \overline{\zeta' - z'}, r_\zeta - r_z)^I$ and

$$q(\zeta, z) = (r_\zeta \cdot (\zeta - z))^a (r_z \cdot (\zeta - z))^b \overline{(r_\zeta \cdot (\zeta - z))^c} \overline{(r_z \cdot (\zeta - z))^d}.$$

For $\zeta \in M_\rho$ and $C_* > 1$ we have

$$\begin{aligned} |\zeta - z(t)| &\leq |\zeta - z_1| + |z(t) - z_1| \leq |\zeta - z_1| + C|z_2 - z_1| \\ &\leq |\zeta - z_1| + CC_*^{-2}|\zeta' - z'_1|^2 \leq C'|\zeta - z_1|. \end{aligned}$$

By (6.4), $|\xi^n - x_1^n| \leq C|\xi_*|$. Thus $|\zeta - z_1| \leq C|\xi_*|$. Therefore, $|\zeta - z(t)| \leq C|\xi_*|$ and

$$|p(\zeta, z(t))| \leq C|\zeta - z(t)|^{|I|} \leq C'|\xi_*|^{|I|}.$$

By (12.2)-(12.3), we have

$$|\{q(\zeta, z(t))\}^{-1}| \leq \frac{C}{||\zeta'_*|^2 + i\xi_*^n|^{a+b+c+d}}.$$

It is easy to see that $|z'(t)| \leq C|x_2 - x_1| \leq C|z_2 - z_1|$. Then $|\partial_t q(\zeta, z(t))^{-1}|$ does not exceed the sum of

$$\frac{C|z_2 - z_1|}{|(r_\zeta \cdot (\zeta - z(t)))^{a'} (r_{z(t)} \cdot (\zeta - z(t)))^{b'} \overline{(r_\zeta \cdot (\zeta - z(t)))^{c'}} (r_{z(t)} \cdot (\zeta - z(t)))^{d'}|},$$

where $a' + b' + c' + d' = a + b + c + d + 1$. Applying the product rule, we get

$$|\partial_t p(\zeta, z(t))| \leq C|z_2 - z_1| \sum_{|I'|=|I|-1} |(\zeta - z(t), \overline{\zeta - z(t)}, r_\zeta - r_{z(t)})^{I'}|.$$

Therefore,

$$|\partial_t p(\zeta, z(t))| \leq C|z_2 - z_1| |(\zeta'_*, \xi_*^n)|^{|I|-1}, \left| \partial_t \frac{1}{q(\zeta, z(t))} \right| \leq \frac{C|z_2 - z_1|}{\|\zeta'_*\|^2 + i\xi_*^n|^{a+b+c+d+1}}.$$

By the mean-value-theorem, we get for $\zeta \in M_\rho \cap \{|\zeta' - z'| > \rho\}$

$$\begin{aligned} |k(\zeta, z_2) - k(\zeta, z_1)| &\leq C|z_2 - z_1| \left\{ \frac{|\xi_*^n|^{|I|}}{\|\zeta'_*\|^2 + i\xi_*^n|^{a+b+c+d+1}} + \frac{|\xi_*^n|^{|I|-1}}{\|\zeta'_*\|^2 + i\xi_*^n|^{a+b+c+d}} \right\} \\ &\leq C'|z_2 - z_1| \frac{|\xi_*^n|^{|I|}}{\|\zeta'_*\|^2 + i\xi_*^n|^{a+b+c+d+1}} \\ &\leq \sum_{|J|=|I|} \frac{C'|z_2 - z_1| |\xi_*^J|}{\|\zeta'_*\|^2 + i\xi_*^n|^{a+b+c+d+1}}. \end{aligned}$$

Applying Lemma 7.1 with $\rho_1 = C_*|z_2 - z_1|^{1/2} < 9\rho = \rho_0$ and $\beta \geq -2$, we get

$$\mathcal{I}_3 \leq C|z_2 - z_1| \rho_1^{-1} \leq C|z_2 - z_1|^{1/2}. \quad \square$$

13. Proof of Theorem 1.1

We now prove Theorem 1.1, following a KAM argument in [5]. We restate the theorem.

Theorem. *Let $M: r = 0$ be a strongly pseudoconvex real hypersurface of class \mathcal{C}^2 in \mathbb{C}^n with $n \geq 4$. Let ω be a continuous $r \times r$ matrix of $(0, 1)$ -forms on M satisfying the integrability condition $\bar{\partial}_b \omega = \omega \wedge \omega \pmod{\bar{\partial} r}$. Near each point of M there exists a non-singular matrix $A \in \mathcal{C}^{1/2}(M)$ such that $\bar{\partial}_b A = -A\omega \pmod{\bar{\partial} r}$. Moreover, if a is a positive real number, M is of class \mathcal{C}^{a+2} and $\omega \in \mathcal{C}^a(M)$, there is a solution $A \in \mathcal{C}^a(M)$; if k is a positive integer, $\omega \in \mathcal{C}^k$ and $M \in \mathcal{C}^{k+\frac{5}{2}}$, there is a solution $A \in \mathcal{C}^{k+\frac{1}{2}}(M)$.*

Note that not all solutions have the same regularity. If u is a continuous CR function vanishing nowhere on M , then uA is still a solution.

Non-isotropic dilations. The non-isotropic dilation $T_\delta: z \rightarrow (\sqrt{\delta}z', \delta z^n)$ with $\delta > 0$ does not preserve the real hypersurface M . However, it is obvious that it sends $M^\delta = T_\delta^{-1}M$ onto M . By abuse of notation, we will denote T_δ its restriction to $\mathbb{C}^{n-1} \times \mathbb{R}$.

We want to show that M can be put in the form (2.1)-(2.2) after a second order normalization and a non-isotropic dilation.

Let M be a graph $y^n = |z'|^2 + \hat{r}(z', x^n)$ over D . Recall that $M_\rho = M \cap \{(x^n)^2 + y^n < \rho^2\}$ and

$$D_\rho = \pi(M_\rho) = \left\{ (z', x^n) \in D: |z'|^2 + |x^n|^2 + \hat{r}(z', x^n) < \rho^2 \right\}.$$

Set $r(z) = -y^n + \hat{r}(z', x^n)$. Define

$$\hat{r}^\delta(z) = \delta^{-2}\hat{r}(\delta z', \delta^2 x^n), \quad r^\delta(z) = \delta^{-2}r(\delta z', \delta^2 x^n).$$

Then $M^\delta = T_\delta^{-1}M$ is the graph over $D^\delta = T_\delta^{-1}D$, given by

$$y^n = |z'|^2 + \hat{r}^\delta(z', x^n).$$

For $0 < \delta < 1$, we have

$$M_\rho^\delta = M^\delta \cap \{|x^n|^2 + y^n < \rho^2\} \subset M^\delta \cap \{\delta^2|x^n|^2 + y^n < \rho^2\}.$$

So $D_\rho^\delta \stackrel{\text{def}}{=} \pi(M_\rho^\delta) \subset T_\delta^{-1}D_\rho$ for $0 < \delta < 1$.

In Theorem 1.1, we need a local solution A . By a change of local holomorphic coordinates, we may assume that $\hat{r}(0) = \partial\hat{r}(0) = \partial^2\hat{r}(0) = 0$. Therefore

$$\|\hat{r}^\delta\|_{D_{1,2}^\delta} < \epsilon, \quad \overline{D_1^\delta} \subset D^\delta,$$

if δ is sufficiently small.

Integrability conditions. We will find our solution through a sequence of frame changes. We also need a small norm of initial ω via dilation. Therefore, we need to verify that the integrability condition is preserved under dilation and frame changes.

Recall that for $\varphi = \sum_{|I|=q} \varphi_I d\bar{z}^I$, we define $\bar{\partial}_M \varphi = \sum_{|I|=q, 1 \leq \alpha < n} X_{\bar{\alpha}} \varphi_I d\bar{z}^{\bar{\alpha}} \wedge d\bar{z}^I$ for $X_{\bar{\alpha}} = \partial_{\bar{z}^\alpha} - r_{\bar{z}^\alpha}/r_{z^n} \partial_{z^n}$. A direct computation shows that $\delta T_\delta^{-1} X_{\bar{\alpha}} = X_{\bar{\alpha}}^\delta \equiv \partial_{\bar{z}^\alpha} - r_{\bar{z}^\alpha}^\delta/r_{z^n}^\delta \partial_{z^n}$. Thus, $T_\delta^* \bar{\partial}_M = \bar{\partial}_{M^\delta} T_\delta^*$. This shows that the formal integrability condition is invariant under dilation, *i.e.* $\bar{\partial}_{M^\delta} \omega^\delta = \omega^\delta \wedge \omega^\delta$. If $\omega = \sum \varphi_J d\bar{z}^J$ is a tangential $(0, 1)$ -form on M , then $T_\delta^* \omega$ is a tangential $(0, 1)$ -form on M^δ . Moreover, if $\omega \in \mathcal{C}^a(M)$, then

$$T_\delta^* \omega = \delta \sum_{\alpha=1}^{n-1} \varphi_{\bar{\alpha}} \circ T_\delta d\bar{z}^{\bar{\alpha}}, \quad \lim_{\delta \rightarrow 0} \|\omega^\delta\|_{\mathcal{C}^a(D_1^\delta)} = 0. \tag{13.1}$$

In other words, we have achieved the smallness of ω via dilation alone.

We now consider the integrability condition under a frame change. We are given an $r \times r$ matrix of continuous $(0, 1)$ -forms ω on M . We assume that $\bar{\partial}_b \omega = \omega \wedge \omega \pmod{\bar{\partial}r}$. Without loss of generality, we may assume that ω are tangential. The formal integrability condition is that as currents

$$\bar{\partial}_M \omega = \omega \wedge \omega. \quad (13.2)$$

Recall that our goal is to find a non-singular matrix A which solves

$$\bar{\partial}_M A + A\omega = 0. \quad (13.3)$$

We will consider only the solution A such that both A and $\bar{\partial}_M A$ are continuous. For any such A , the transformation $\omega \rightarrow \tilde{\omega} = (\bar{\partial}_M A + A\omega)A^{-1}$ preserves the integrability condition (13.2). Indeed, differentiating $\tilde{\omega}A \equiv \bar{\partial}_M A + A\omega$ and then using $\omega = A^{-1}\tilde{\omega}A - A^{-1}\bar{\partial}_M A$, $\bar{\partial}_M \omega = \omega \wedge \omega$, we verify $\bar{\partial}_M \tilde{\omega} = \tilde{\omega} \wedge \tilde{\omega}$ by

$$\begin{aligned} (\bar{\partial}_M \tilde{\omega})A - \tilde{\omega} \wedge \bar{\partial}_M A &= \bar{\partial}_M A \wedge \omega + A\bar{\partial}_M \omega = \bar{\partial}_M A \wedge (A^{-1}\tilde{\omega}A - A^{-1}\bar{\partial}_M A) \\ &\quad + (\tilde{\omega}A - \bar{\partial}_M A) \wedge (A^{-1}\tilde{\omega}A - A^{-1}\bar{\partial}_M A) \\ &= \tilde{\omega} \wedge \tilde{\omega}A - \tilde{\omega} \wedge \bar{\partial}_M A. \end{aligned}$$

Assume that the matrix ω is of class \mathcal{C}^a . In what follows, all constants, including δ , will depend on a . For simplicity this dependence will not be indicated sometimes. However, constants C_0 , δ_* and ϵ do not depend on a .

Proof of Theorem 1.1. We need to find a non-singular matrix $A = I + B$, defined near the origin of M , such that

$$\bar{\partial}_M B + \omega + B\omega = 0.$$

It suffices to find a $\delta > 0$ and a non-singular matrix A^δ defined near $0 \in M^\delta$ such that $\bar{\partial}_{M^\delta} \omega^\delta + A^\delta \omega^\delta = 0$. Then $A^\delta \circ T_\delta^{-1}$ is a solution to the original equation.

Take $\rho_0 = 1$, $\sigma_j = 2^{-j-1}$ and $\rho_{j+1} = (1 - \sigma_j)\rho_j$. Then $\rho_\infty = \lim_{j \rightarrow \infty} \rho_j > 0$. We will assume that $0 < \delta \leq \delta_*$. We want to apply our estimates for P' and Q' . So we choose $\delta_* \in (0, 1]$ such that the homotopy formula holds on M_ρ^δ for $\rho \in (0, 1]$ and $\delta \in (0, \delta_*]$. For $\rho_\infty < \rho \leq 1$ we have

$$\|P'\varphi\|_{\mathcal{C}^a(M_{(1-\sigma)\rho}^\delta)} \leq C\sigma^{-s}\|\varphi\|_{\mathcal{C}^{a+2}(M_\rho^\delta)}, \quad (13.4)$$

where P' is either of operators P' , Q' in the homotopy formula on M_ρ^δ . We emphasize that the constant C_a is independent of $\delta \in (0, \rho_*)$ and $\rho \in (\rho_\infty, 1]$. We have also absorbed $\|r^\delta\|_{\mathcal{C}^{a+2}(D_{\rho_j}^\delta)}$ into C_a , since $\hat{r}^\delta(z', x^n) = \delta^{-2}\hat{r}(\delta z', \delta^2 x^n)$ and $\hat{r}(0) = \partial\hat{r}(0) = 0$ imply that

$$\|\hat{r}^\delta\|_{\mathcal{C}^{a+2}(D_{\rho_j}^\delta)} \leq C\|\hat{r}\|_{\mathcal{C}^2(D_{\delta\rho_0})} + \delta^a\|\hat{r}\|_{\mathcal{C}^{a+2}(D_{\delta\rho_0})} < C_a, \quad 0 < \delta < \delta_*.$$

Set $M_j = M_{\rho_j}^\delta$ and $\|\cdot\|_{\rho_{j+1},a} = \|\cdot\|_{C^a(M_{j+1})}$. We have $M_{j+1} \subset M_0$. Let $\omega_0 = T_\delta^* \omega$, restricted on M_0 . On M_j we have the homotopy formula $\varphi = \bar{\partial}_{M^\delta} P'_j \varphi + Q'_j \bar{\partial}_{M^\delta} \varphi$, where $P'_j = P'_{M_j}$ and $Q'_j = Q'_{M_j}$.

Using $\bar{\partial}_{M^\delta} \omega_0 = \omega_0 \wedge \omega_0$, we arrive at the equation

$$\bar{\partial}_{M^\delta} (B_0 + P'_0 \omega_0) + Q'_0 (\omega_0 \wedge \omega_0) + B_0 \omega_0 = 0,$$

where P'_0, Q'_0 are applied entrywise to the matrices. We use the approximate solution

$$B_0 = -P'_0 \omega_0.$$

Assume that $A_0 = I + B_0$ is invertible. We repeat this procedure and get $B_j = -P'_j \omega_j$ and

$$\omega_{j+1} = (\bar{\partial}_M A_j + A_j \omega_j) A_j^{-1} = \{Q'_j (\omega_j \wedge \omega_j) - (P'_j \omega_j) \omega_j\} (I - P'_j \omega_j)^{-1}. \quad (13.5)$$

Here, we need all $A_j = I + B_j$ to be non-singular on M_j . We want to show that $\lim_{j \rightarrow \infty} A_j A_{j-1} \cdots A_0$ is a solution.

We now estimate $\|B_j\|_{\rho_{j+1},a}$ and $\|\omega_{j+1}\|_{\rho_{j+1},a}$.

For an $r \times r$ matrix $B = (b_i^j)$ of functions on M_ρ we define $\|B\|_{\rho,a} = \max\{\|b_i^j\|_{\rho,a}\}$. If B, D are two such matrices, we have

$$\|DB\|_{\rho,0} \leq r^2 \|D\|_{\rho,0} \|B\|_{\rho,0}, \quad \|B^l\|_{\rho,0} \leq r^l \|B\|_{\rho,0}^l.$$

Assume that $\rho_\infty < \rho \leq \rho_0$. We want to show that if $\|B\|_{\rho,0} \leq \frac{1}{2r}$, then

$$\|D\|_{\rho,a} \leq c_a \|B\|_{\rho,a}, \quad 0 \leq a < \infty, \quad (13.6)$$

where $c_a > 1$ depends on a, r, n . Let $A^{-1} = I + D$. We know that $D = \sum_{l \geq 1} (-1)^l B^l$ and

$$\|D\|_{\rho,0} \leq \frac{r \|B\|_{\rho,0}}{1 - r \|B\|_{\rho,0}} \leq 2r \|B\|_{\rho,0},$$

which is (13.6) with $a = 0$. Since I is constant, then

$$\begin{aligned} D(z_2) - D(z_1) &= A(z_2)^{-1} - A(z_1)^{-1} = A(z_1)^{-1} (A(z_1) - A(z_2)) A(z_2)^{-1} \\ &= A(z_1)^{-1} (B(z_1) - B(z_2)) A(z_2)^{-1}. \end{aligned}$$

Using (13.6) with $a = 0$, we get $\|D\|_{\rho,a} \leq C \|B\|_{\rho,a}$ if $0 < a \leq 1$. Assume that (13.6) holds when $[a] < k$. Let $[a] = k \geq 1$. Applying the product rule to $(I + B)D = -B$ and multiplying from left by $(I + B)^{-1} = I + D$, we get

$$\partial D = (I + D)(\partial B)D - (I + D)\partial B.$$

By Proposition A.4 and the induction assumption, we obtain

$$\|D\|_{\rho,a} \leq C_a\{(1 + \|B\|_{\rho,a-1})\|B\|_{\rho,1} + \|B\|_{\rho,a}\} \leq C'_a\|B\|_{\rho,a},$$

which proves (13.6).

Since $B_j = -P'_j\omega_j$, by (13.4) we have

$$\|B_j\|_{\rho_{j+1},a} \leq c_a^*\sigma_j^{-s_a}\|\omega_j\|_{\rho_j,a}, \quad c_a^* > 1. \quad (13.7)$$

We want to achieve $\|B_j\|_{\rho_{j+1},0} \leq \frac{1}{2r}$. So it suffices to obtain

$$\|\omega_j\|_{\rho_j,a} \leq \frac{\sigma_j^{s_a}}{2rc_a^*c_a} = b_j, \quad j = 0, 1, 2, \dots \quad (13.8)$$

By (13.6)-(13.8), we have $\|(I+B)^{-1}\|_{\rho_j,a} \leq 1 + \|D\|_{\rho_j,a} \leq 2$. Using (13.4)-(13.5) and the estimates on matrix products, we get

$$\begin{aligned} \|\omega_{j+1}\|_{\rho_{j+1},a} &\leq 2 \cdot r^2 \{ \|Q'_j(\omega_j \wedge \omega_j)\|_{\rho_{j+1},a} \\ &\quad + r^2 \|B_j\|_{\rho_{j+1},a} \|\omega_j\|_{\rho_{j+1},a} \} \leq C_a^*\sigma_j^{-s_a}\|\omega_j\|_{\rho_j,a}^2. \end{aligned} \quad (13.9)$$

Assume that $\|\omega_0\|_{\rho_0,a} \neq 0$. Otherwise the theorem holds trivially. Define

$$\hat{b}_{j+1} = C_a^*\sigma_j^{-s_a}\hat{b}_j^2, \quad \hat{b}_0 = \|\omega_0\|_{\rho_0,a}.$$

By (13.1), we choose a dilation T_δ with $\delta \in (0, \delta_*]$ such that $\omega_0 = T_\delta\omega$ satisfies

$$\hat{b}_0 = \|\omega_0\|_{\rho_0,a} \leq b_0.$$

Then $r_j = \hat{b}_{j+1}/\hat{b}_j$ satisfies

$$\begin{aligned} r_0 &= C_a^*\sigma_0^{-s_a}\hat{b}_0, \quad r_j = 2^{s_a}(r_{j-1})^2, \\ r_1/r_0 &= 2^{s_a}r_0, \quad r_{j+1}/r_j = (r_j/r_{j-1})^2. \end{aligned}$$

This shows that r_{j+1}/r_j , and hence r_j, \hat{b}_j , converge rapidly, if \hat{b}_0 is sufficiently small. Specifically,

$$r_{j+1}/r_j = (r_1/r_0)^{2^j}, \quad r_j = r_0(r_1/r_0)^{2^j-1}, \quad \hat{b}_j = \hat{b}_0 r_0^j (r_1/r_0)^{2^j-j-1}.$$

Recall that $\sigma_j = 2^{-j-1}$. Clearly, $\|\omega_j\|_{\rho_j,a} \leq \hat{b}_j \leq \hat{b}_0 r_0^j \leq \frac{2^{-s_a}}{2rc_a^*c_a}(2^{-s_a})^j = b_j$, provided

$$\hat{b}_0 \leq \frac{2^{-s_a}}{2rc_a^*c_a}, \quad r_0 \leq 2^{-s_a}, \quad r_1/r_0 = 2^{s_a}r_0 \leq \frac{1}{2}.$$

Therefore, we have shown that if $\omega \in \mathcal{C}^a$ and $M \in \mathcal{C}^{2+a}$ then $(I + B_j) \cdots (I + B_0)$ converges in \mathcal{C}^a -norm on $\cap D_{\rho_j}^\delta$ to an invertible matrix A_∞ . When $a = k$ is an integer and $M \in \mathcal{C}^{k+5/2}$, or when $M \in \mathcal{C}^2$ for $k = 0$, we have

$$\|B_j\|_{\rho_{j+1}, k+1/2} \leq c_k \sigma_j^{-s_k} \|\omega_j\|_{\rho_j, k}.$$

Since $\|\omega_j\|_{\rho_j, k}$ tends to zero rapidly, it is obvious that B_j converges to 0 rapidly in $\mathcal{C}^{k+1/2}$ on $\cap D_{\rho_j}^j$. This shows that A^∞ is of class $\mathcal{C}^{k+1/2}$.

To complete the proof, we check that $\bar{\partial}_{M^\delta} A^\infty + A^\infty \omega_0 = 0$. Recall that for $A_j = I + B_j$, we have $\bar{\partial}_{M^\delta} A_j + A_j \omega_j = w_{j+1} A_j$. Thus

$$\begin{aligned} \omega_{j+1} A_j A_{j-1} &= (\bar{\partial}_{M^\delta} A_j) A_{j-1} + A_j \omega_j A_{j-1} \\ &= (\bar{\partial}_{M^\delta} A_j) A_{j-1} + A_j (\bar{\partial}_{M^\delta} A_{j-1} + A_{j-1} \omega_{j-1}) \\ &= \bar{\partial}_{M^\delta} (A_j A_{j-1}) + A_j A_{j-1} \omega_{j-1}. \end{aligned}$$

Inductively, we get $\bar{\partial}_{M^\delta} (A_j \cdots A_0) + A_j \cdots A_0 \omega_0 = \omega_{j+1} A_j \cdots A_0$. Taking the limits, we get $\bar{\partial}_{M^\delta} A^\infty + A^\infty \omega_0 = 0$ on $M_{\rho_\infty}^\delta$. When $k = 0$, the derivatives in the sense of currents are continuous and the above computation is valid as currents. \square

In the above argument, we obtain the rapid convergence of B_j, ω_j in \mathcal{C}^a norm in one step when a is finite. One can also establish a rapid convergence of ω_j, B_j first in \mathcal{C}^0 -norm and then in higher order derivatives. See [23], [5] for details.

Appendix

A. Hölder inequalities

The main purpose of this appendix is to present some Hölder inequalities on domains in \mathbb{R}^m . We do not claim any originality in deriving these inequalities. In fact, we will just modify formulation and proofs of Hörmander [9]. The inequalities in [9] are for a fixed convex domain. In our applications, we need to allow the domain D_ρ to vary. Therefore, we will derive them in full details and omit simple repetitions only.

We say that a domain D in \mathbb{R}^m has the *cone property* if the following hold: (i) Given two points p_0, p_1 in D there exists a piecewise \mathcal{C}^1 curve $\gamma(t)$ in D such that $\gamma(0) = p_0$ and $\gamma(1) = p_1$, $|\gamma'(t)| \leq C_* \|p_1 - p_0\|$ for all t except finitely many values. The diameter of D is less than C_* . (ii) For each point $x \in \bar{D}$, D contains a cone V with vertex x , opening $\theta > C_*^{-1}$ and height $h > C_*^{-1}$.

We will denote $C_*(D)$ a constant $C_* > 1$ satisfying (i) and (ii).

In this appendix, by a cone $V = V(\theta, h, v)$ with vertex at the origin, opening $\theta > 0$ and height $h > 0$, and centered at positive v axis where v is a unit vector, we mean

$$V = \{t \in \mathbb{R}^m : v \cdot t > \theta^{-1} \|t - (v \cdot t)v\|, v \cdot t < h\}, \tag{A.1}$$

where $\|t\| = (|t^1|^2 + \dots + |t^m|^2)^{\frac{1}{2}}$. Note that $x + V$ is a cone with vertex at x . Note that each cone satisfying (ii) contains a ball of radius at least $C_*(D)/C$. Let D_1, D_2 be domains of the cone property, and let V_1, V_2 be cones in (ii) for D_1, D_2 respectively. Assume that the vertex of V_i is p_i . Then $D_1 \times D_2$ contains the convex hull of (p_1, p_2) and a ball of radius depending only on $C_*(D_1), C_*(D_2)$. Therefore, $D_1 \times D_2$ still has the cone property.

For the rest of the appendix, until Proposition A.5, we assume that the domain D has the cone property unless stated otherwise. The constants in all Hölder inequalities will depend on m and $C_*(D)$. For simplicity this dependence will not be expressed sometimes.

Let $k \geq 0$ be an integer. For a complex-valued function u on $D \subset \mathbb{R}^m$, define

$$\begin{aligned} \|\partial^k u\|_{D,0} &= \sup_{x \in D, |I|=k} |\partial^I u(x)|, \quad \|u\|_{D,k} = \max_{0 \leq j \leq k} \|\partial^j u\|_{D,0}, \\ |u|_{D,\alpha} &= \sup_{x,y \in D} \frac{|u(x) - u(y)|}{|x - y|^\alpha}, \quad 0 < \alpha \leq 1, \\ \|u\|_{D,k+\alpha} &= \max\{\|u\|_{D,k}, |\partial^I u|_{D,\alpha} : |I| = k\}, \quad 0 < \alpha < 1. \end{aligned}$$

If $A = (a_i^j)$ is a matrix of functions on D , we define $\|A\|_{D,k+\alpha} = \max\{\|a_i^j\|_{D,k+\alpha}\}$. We also define

$$|u|_{D,k+\alpha} = \max_{|I|=k} |\partial^I u|_{D,\alpha}, \quad 0 < \alpha \leq 1.$$

By (i) of the cone property and the fundamental theorem of calculus

$$C_0^{-1} |u|_{D,k} \leq \|\partial^k u\|_{D,0} \leq |u|_{D,k}, \quad k = 1, 2, \dots, \tag{A.2}$$

provided that $u \in C^k(D)$. It will be convenient to use both $|u|_{D,k}$ and $\|\partial^k u\|_{D,0}$.

Lemma A.1. *Let $D \subset \mathbb{R}^m$ satisfy (i) of the cone property. Let f be a C^1 map from D into \mathbb{R}^m . There exists a constant $C_0 > 1$ such that if $|f' - I| < C_0^{-1}$ on D , then f is a C^1 diffeomorphism from D onto D' . Moreover, $\|f^{-1} - I\|_{D',1} \leq C \|f - I\|_{D,1}$.*

Proof. Take two points p_0, p_1 in D . By assumption there exists a piecewise C^1 curve γ in D such that $\gamma(0) = p_0, \gamma(1) = p_1$, and $|\gamma'(t)| \leq C_* \|p_1 - p_0\|$. Write $f = I + \tilde{f}$. We have

$$f(p_1) - f(p_0) = p_1 - p_0 - \int_0^1 \nabla \tilde{f}(\gamma(t)) \cdot \gamma'(t) dt.$$

Since $\|\tilde{f}'\|_{D,0} < C_0^{-1}$, then for C_0 sufficiently large we get

$$\frac{1}{2}\|p_1 - p_0\| \leq \|f(p_1) - f(p_0)\| \leq 2\|p_1 - p_0\|. \tag{A.3}$$

Now it is obvious that f is a C^1 diffeomorphism from D onto D' and that D' satisfies (i) of the cone property. Let $\tilde{g} = f^{-1} - I$. Then $\tilde{g} \circ f = -\tilde{f}$. In particular, $\|\tilde{g}\|_{D',0} \leq \|\tilde{f}\|_{D,0}$, and by (A.3), $|\tilde{g}|_{D',1} \leq C|\tilde{f}|_{D,1}$. \square

We are ready to derive Hölder inequalities. We start with two lemmas in [9] for our domains.

Lemma A.2. *Let $0 < a < b$. Let D satisfy the cone property. If $|u|_{D,a} \leq 1$ and $|u|_{D,b} \leq 1$ then $|u|_{D,c} \leq C$ for $a < c < b$, where C depends only on a, b and $C_*(D)$.*

Proof. For simplicity, denote $|u|_{D,a}$ by $|u|_a$. Since the diameter of D is bounded by some constant C , it is obvious that $|u|_c \leq C'|u|_{c'}$ if $k < c < c' \leq k + 1$. Therefore, it suffices to prove the inequality when c is an integer.

Let $V \subset D$ be a cone as stated in the cone property. We may assume that $0 \in V$.

If a is an integer, we get $|\partial^a u| \leq 1$ on V . If a is not an integer, let P be its Taylor polynomial of degree $[a]$ at 0 . Set $v = u - P$. Since $\partial^I P$ are constants for all $|I| \geq [a]$, then $|v|_{c'} = |u|_{c'}$ for all $c' > [a]$. Therefore, we may assume that $\partial^I u(0) = 0$ for all $|I| = [a]$. Now $|u|_a \leq 1$ implies that $|\partial^I u| < C_0$ for all $|I| = [a]$.

We want to use the mean-value-theorem repeatedly. Let us first look at the one-variable case. When f is C^k on $[0, 1]$ and $\|f\|_0 \equiv \|f\|_{[0,1],0} < C_0$, there is a point t in $[0, 1]$ such that $|f'(t)| \leq 2C_0$. One can divide $[0, 1]$ into a sufficient number of equal parts and find a point $t_j \in [0, 1]$ such that $|f^{(j)}(t_j)| < C'$ for $j = 1, \dots, k$. If $|f|_a \leq 1$ for some $a \in (k, k + 1]$, we obtain further that $\|f^{(k)}\|_0 < C_k$. Consequently, $\|f^{(j)}\|_0 < C_j$ hold for $j = k, k - 1, \dots, 0$.

Return to our case. Fix a polydisc Δ^m in V with side larger than C^{-1} . Fix I with $|I| = [a]$. Using $|\partial^I u| < C_0$ for $|I| = [a]$ and $|u|_b \leq 1$, by the one-variable argument we obtain $|\partial^j \partial^I u| < C$ on Δ^m for $j = [b] - [a], \dots, c - [a]$. So $|\partial^j u| < C$ on Δ^m for $j = c, \dots, [b]$. If $[b] < b, |u|_b \leq 1$ implies that $|\partial^{[b]} u| < C'$ on D . If $b = [b]$ the assumption and (A.2) implies $|\partial^b u| \leq 1$. Using a path connecting a point in D to Δ^m and $|\partial^j u| < C$ on Δ^m for $j = [b], \dots, c$, we get $|\partial^j u| \leq C''$ on D for $j = [b] - 1, \dots, c$. Using (A.2) again, we get $|u|_c \leq C$. \square

Example. Let $f(x) = x + x^3$ and $D = [0, \epsilon]$. Then $f'(0) = f'(x) - \int_0^x f''(t) dt$ and

$$\epsilon f'(0) = \int_0^\epsilon f'(x) dx - \int_0^\epsilon \int_0^x f''(t) dt dx = f(\epsilon) - f(0) - \int_0^\epsilon \int_0^x f''(t) dt dx.$$

This shows that $1 = |f'(0)| \leq C_1 \|f\|_0 + C_2 \|f''\|_0 \leq 2\epsilon C_1 + 6\epsilon C_2$. However, $|C_1| + |C_2|$ tends to ∞ as $\epsilon \rightarrow 0$. This example demonstrates that in some inequalities derived in this appendix, the constants indeed depend on $C_*(D)$. For special domains used in this paper, we will find these constants by dilation; see Proposition A.5.

Lemma A.3. *Let $D \subset \mathbb{R}^m$ satisfy the cone property. Set $\|\cdot\|_{D,a} = \|\cdot\|_a$. If $0 < a < b, c = \lambda a + (1 - \lambda)b$ and $0 < \lambda < 1$, then $|u|_c \leq C|u|_a^\lambda(|u|_a + |u|_b)^{1-\lambda}$, where C depends only on a, b .*

Proof. The case $|u|_a \geq |u|_c$ is obvious and we may assume that $|u|_a \leq |u|_c$. If $|u|_a = 0$, then u is a polynomial of degree $< a$. Then $|u|_c = 0$ too for $c > a$, and the inequality holds. We may assume that $|u|_a \neq 0$. Without loss of generality, we may assume that $|u|_a = 1 < |u|_c$. If $|u|_b \leq 1$, Lemma A.2 implies that $|u|_c \leq C$ and the inequality holds. Therefore, it suffices to verify

$$|u|_c \leq C|u|_b^{1-\lambda}, \quad \text{if } |u|_a = 1 < |u|_b. \tag{A.4}$$

We first assume the inequality for integer c and verify it for non-integer c . Set $[c] = k$. We have $\lambda = \frac{b-c}{b-a}$ and $1 - \lambda = \frac{c-a}{b-a}$. Depending on whether a, b are in $[k, k + 1]$, we have the following cases.

Case i) $k \leq a < c < b \leq k + 1$. Since $c = \lambda a + (1 - \lambda)b$ then $c - k = \lambda(a - k) + (1 - \lambda)(b - k)$. Consider first the case $a = k$. Then $c - k = (1 - \lambda)(b - k)$. Let $v = \partial^I u$ with $|I| = k$. Hence

$$\begin{aligned} \frac{|v(y) - v(x)|}{|y - x|^{c-k}} &= |v(y) - v(x)|^\lambda \left| \frac{v(y) - v(x)}{|y - x|^{b-k}} \right|^{1-\lambda} \leq 2^\lambda \|v\|_0^\lambda |v|_{b-k}^{1-\lambda} \\ &\leq C'|u|_k^\lambda |\partial^I u|_{b-k}^{1-\lambda} \leq C'|u|_k^\lambda |u|_b^{1-\lambda}, \end{aligned}$$

where the second last inequality is obtained by (A.2) using $k = a > 0$. Therefore, $|u|_c \leq C|u|_a^\lambda |u|_b^{1-\lambda}$. Assume now that $a - k > 0$. Then $a - k, b - k, c - k$ are in $(0, 1]$. Computing the Hölder ratio gives us $|v|_{c-k} \leq |v|_{a-k}^\lambda |v|_{b-k}^{1-\lambda}$, i.e. $|u|_c \leq |u|_a^\lambda |u|_b^{1-\lambda}$. We emphasize that we have proved $|u|_c \leq C|u|_a^\lambda |u|_b^{1-\lambda}$ (and hence (A.4) for case i)) without using any condition on $|u|_a, |u|_b, |u|_c$ other than that on a, b, c, k .

Case ii) $a < k < c < b \leq k + 1$. We get $|u|_c \leq C|u|_k^{(b-c)/(b-k)} |u|_b^{(c-k)/(b-k)}$, by case i). By the assumption for the integer case (applied to triple $a < k < b$) we obtain

$$|u|_k \leq C|u|_b^{(k-a)/(b-a)}. \tag{A.5}$$

Eliminating $|u|_k$ from two inequalities gives us (A.4); indeed

$$\frac{k - a}{b - a} \cdot \frac{b - c}{b - k} + \frac{c - k}{b - k} = \frac{c - a}{b - a}.$$

Case iii) $k \leq a < c < k + 1 < b$. We have $|u|_c \leq C|u|_{k+1}^{(c-a)/(k+1-a)}$, by case i). The assumption on the integer case (applied to triple $a < k + 1 < b$) gives us

$$|u|_{k+1} \leq C|u|_b^{(k+1-a)/(b-a)}. \quad (\text{A.6})$$

Eliminating $|u|_{k+1}$ gives us (A.4).

Case iv) $a < k < c < k + 1 < b$. Then (A.5) and (A.6) are still valid. By i), we have $|u|_c \leq C|u|_k^{k+1-c}|u|_{k+1}^{c-k}$. Using (A.5)-(A.6) and eliminating $|u|_k, |u|_{k+1}$ gives us (A.4).

Finally, we prove (A.4) when c is a positive integer by repeating an argument in [9].

Fix $x_0 \in D$ and let V be a cone in D with vertex x_0 , height and opening $1/C_*$. Since V is convex, for $x \in V$ and $0 < \epsilon < 1$ we can define

$$u_x^\epsilon(y) = u((1 - \epsilon)x + \epsilon y), \quad y \in V.$$

Then $|u_x^\epsilon|_{V,b} \leq \epsilon^b|u|_b$ and $|u_x^\epsilon|_{V,a} \leq \epsilon^a$. Since $|u|_b > 1$, there is an $\epsilon \in (0, 1)$ so that $\epsilon^a = \mu = \epsilon^b|u|_b$. Now, apply Lemma A.2 to the domain V and the function $\mu^{-1}u_x^\epsilon$. For any multiindex I with $|I| = c$, we have

$$\epsilon^c|\partial^I u(x)| = |\partial^I u_x^\epsilon(x)| \leq C_0\mu = C_0(\epsilon^a)^\lambda(\epsilon^b|u|_b)^{1-\lambda}.$$

Canceling ϵ 's shows $|\partial^I u(x)| \leq C_0|u|_b^{1-\lambda}$ for $x \in \bar{V}$. Since C_0 does not depend on $x_0 \in \bar{V}$, we get $\|\partial^c u\|_0 \leq C_0|u|_b^{1-\lambda}$ on D ; by (A.2), $|u|_c \leq C\|\partial^c u\|_0$ and (A.4) follows. \square

Proposition A.4. *Let $D \subset \mathbb{R}^m$ have the cone property and denote $\|\cdot\|_{D,a}$ by $\|\cdot\|_a$. Let a, b, a_j, b_j be nonnegative real numbers.*

- (i) $\|u\|_{\lambda a + (1-\lambda)b} \leq C_{a,b}\|u\|_a^\lambda\|u\|_b^{1-\lambda}$ for $0 < \lambda < 1$.
- (ii) $\|uv\|_a \leq C_a(\|u\|_0\|v\|_a + \|u\|_a\|v\|_0)$.
- (iii) *Suppose that $D_j \subset \mathbb{R}^{n_j}$ has the cone property for $j = 1, \dots, k$. Let a_j, c_j be non-negative real numbers, and let (b_1, \dots, b_k) be in the convex hull of (a_1, \dots, a_k) and (c_1, \dots, c_k) . Then*

$$\prod_{j=1}^k \|u_j\|_{D_j, b_j} \leq C_{k+|a|+|b|+|c|} \left(\prod_{j=1}^k \|u\|_{D_j, a_j} + \prod_{j=1}^k \|u\|_{D_j, c_j} \right).$$

- (iv) *Let f be a map from D into $D' \subset \mathbb{R}^n$. Assume that D' has the cone property. Then*

$$\begin{aligned} \|u \circ f\|_a &\leq C_a(\|u'\|_{D', a-1}\|f'\|_0^a + \|u'\|_{D', 0}\|f'\|_{a-1}) + \|u\|_{D', 0}, \quad a \geq 1, \\ \|u \circ f\|_a &\leq C \min(\|u'\|_{D', 0}\|f\|_a, \|u\|_{D', a}\|f'\|_0^a) + \|u\|_{D', 0}, \quad 0 \leq a \leq 1. \end{aligned}$$

(v) Let $f = I + \tilde{f}$ be a C^a map from D into \mathbb{R}^m . There exists $C_0 > 1$ such that if $|\tilde{f}'| < C_0^{-1}$ then

$$\|f^{-1} - I\|_{f(D),a} \leq C_a \|\tilde{f}\|_a, \quad a \geq 0.$$

(vi) Let $f = I + \tilde{f}$ be a C^1 map from D into $D' \subset \mathbb{R}^m$ with $|f'| < C_0$. Assume that $D \cup D'$ is contained in a **convex** domain D'' of the cone property. Then

$$\|u \circ f - u\|_a \leq C_a (\|u'\|_{D'',a} \|\tilde{f}\|_0 + \|u'\|_{D'',0} \|\tilde{f}\|_a), \quad a \geq 0.$$

Proof. (i)-(iii) are proved in [9]. The proof for (iii) is for D_j being the same domain. In fact, by (i), one has $\log \|u_j\|_{D_j,b_j} \leq \lambda \log \|u_j\|_{D_j,a_j} + (1-\lambda) \log \|u_j\|_{D_j,c_j} + \log C$ for all j . Sum over $j = 1, \dots, k$ and take exponential on both sides. The convexity of e^x yields (iii). For $0 \leq a \leq 1$, two inequalities in (iv) are verified directly. Assume that (iv) holds when a is replaced by $a - 1$. Assume that $a > 1$. Then $(u \circ f)' = u'(f)f'$. For both cases of $0 < a - 1 \leq 1$ and $a - 1 > 1$ we obtain

$$\begin{aligned} \|(u \circ f)'\|_{a-1} &\leq C (\|u'(f)\|_{a-1} \|f'\|_0 + \|u'(f)\|_0 \|f'\|_{a-1}) \\ &\leq C' (\|u'\|_{D',a-1} \|f'\|_0^a + \|u'\|_0 \|f'\|_{a-1}), \end{aligned}$$

where $\|u'\|_{a-1} \equiv \|u'\|_{D',a-1}$. Note that for the second inequality when $a > 2$, we have used

$$\begin{aligned} \|u'\|_1 \|f'\|_{a-2} \|f'\|_0 &\leq C \|u'\|_0^{\frac{a-2}{a-1}} \|u'\|_{a-1}^{\frac{1}{a-1}} \|f'\|_0^{\frac{1}{a-1}} \|f'\|_{a-1}^{\frac{a-2}{a-1}} \|f'\|_0 \\ &\leq C' (\|u'\|_{a-1} \|f'\|_0^a + \|u'\|_0 \|f'\|_{a-1}). \end{aligned}$$

This gives us (iv) as $\|u \circ f\|_a \leq \|u\|_0 + \|(u \circ f)'\|_{a-1}$.

(v). It follows immediately from Lemma A.1 when $0 \leq a \leq 1$. We now prove it for $a > 1$ by using a variant of counting scheme in Section 2. Define

$$\widehat{\partial}^{1+k} v = \sum_{j_1+\dots+j_l \leq k} p(\tilde{f}') \partial^{j_1} v \cdots \partial^{j_l} v, \quad j_i = |J_i| - 1 \geq 0, \quad (\text{A.7})$$

where $p(\tilde{f}')$ is a polynomial in $(I + \tilde{f}')^{-1}$ and \tilde{f}' and it might be different when it reoccurs. Let $g = f^{-1}$ and $\tilde{g} = f^{-1} - I$. Let $\mathbf{1}$ denote the identity matrix. We have $\tilde{g}' = -((\mathbf{1} + \tilde{f}')^{-1} \tilde{f}') \circ g$, i.e. $\partial^I \tilde{g} = (\widehat{\partial}^1 \tilde{f}) \circ g$ for $|I| = 1$. Inductively,

$$\partial^J \tilde{g} = (\widehat{\partial}^{|J|} \tilde{f}) \circ g, \quad |J| \geq 1. \quad (\text{A.8})$$

By Lemma A.1, we have $\frac{1}{2}|y_1 - y_0| \leq |g(y_1) - g(y_0)| \leq 2|y_1 - y_0|$. Let $k = [a] - 1$ and $\alpha = a - k - 1$. Thus $\|g\|_{D',1+k+\alpha} \leq C \|f\|_0 + C \sum_{j \leq k} \|\widehat{\partial}^{1+j} \tilde{f}\|_\alpha$. Now

$$\begin{aligned} \|\widehat{\partial}^{1+k} \tilde{f}\|_\alpha &\leq \sum_{j_1+\dots+j_l \leq k} C' \left\{ \|\tilde{f}\|_{1+\alpha} \|\tilde{f}\|_{1+j_1} \cdots \|\tilde{f}\|_{1+j_l} \right. \\ &\quad \left. + \sum_{1 \leq i \leq l} \|\tilde{f}\|_{1+j_1} \cdots \|\tilde{f}\|_{1+j_i+\alpha} \cdots \|\tilde{f}\|_{1+j_l} \right\}. \end{aligned}$$

By (iii) we obtain $\|\tilde{g}\|_{D',1+k+\alpha} \leq \sum C (\|\tilde{f}\|_1^l + \|\tilde{f}\|_1^{l-1}) \|\tilde{f}\|_{1+k+\alpha} \leq C' \|\tilde{f}\|_a$.

(vi). By convexity of D'' , we have $u(x + \tilde{f}(x)) - u(x) = \int_0^1 u'(x + t\tilde{f}(x)) dt$. Consider case $0 \leq a < 1$. We have

$$\begin{aligned} \left| \int_0^1 \{u'(y + t\tilde{f}(y)) - u'(x + t\tilde{f}(x))\} dt \right| &\leq \max_t \|u'\|_{D'',a} |y - x + t(\tilde{f}(y) - \tilde{f}(x))|^a \\ &\leq C \|u'\|_{D'',a} (1 + \|\tilde{f}\|_{D,1}) |y - x|^\alpha. \end{aligned}$$

By $\|\tilde{f}\|_1 \leq C$, one sees easily that

$$\|u(I + \tilde{f}) - u\|_a \leq C(\|u'\|_{D'',a} \|\tilde{f}\|_0 + \|u'\|_{D'',0} \|\tilde{f}\|_a),$$

which is (vi) for $0 \leq a < 1$. Assume that the above inequality holds when a is replaced by $a - 1$. Assume that $a \geq 1$. We need to estimate the $\|\cdot\|_{a-1}$ norm of

$$\partial_{x_i}(u \circ (I + \tilde{f}) - u) = (\partial_{y_i} u) \circ (I + \tilde{f}) - \partial_{x_i} u + \sum_{1 \leq j \leq m} (\partial_{y_j} u) \circ (I + \tilde{f}) \cdot \partial_{x_i} \tilde{f}^j. \tag{A.9}$$

By the induction assumption, we have

$$\|(\partial_{y_i} u) \circ (I + \tilde{f}) - \partial_{x_i} u\|_{a-1} \leq C(\|\partial^2 u\|_{D'',a-1} \|\tilde{f}\|_0 + \|\partial^2 u\|_{D'',0} \|\tilde{f}\|_{a-1}).$$

We need to put $\|\partial^2 u\|_{D'',l} \|\tilde{f}\|_{a-1-l} \leq \|u'\|_{D'',l+1} \|\tilde{f}\|_{a-1-l}$ into the desired form. By (iii), we get

$$\|u'\|_{D'',l+1} \|\tilde{f}\|_{a-1-l} \leq C(\|\tilde{f}\|_0 \|u'\|_{D'',a} + \|u'\|_{D'',0} \|\tilde{f}\|_a). \tag{A.10}$$

We now treat the term in the sum of (A.9). Set $\|u'\|_b \equiv \|u'\|_{D'',b}$. By (iv) we have $\|(\partial_{y_j} u)(I + \tilde{f})\|_{a-1} \leq C(\|u'\|_{a-1} + \|u'\|_1 \|\tilde{f}\|_{a-1}) + \|u'\|_0$. Thus

$$\begin{aligned} \|(\partial_{y_j} u)(I + \tilde{f}) \cdot \partial_{x_i} \tilde{f}^j\|_{a-1} &\leq C((\|u'\|_{a-1} + \|u'\|_1 \|\tilde{f}\|_{a-1}) \|\tilde{f}\|_1 + \|u'\|_0 \|\tilde{f}\|_a) \\ &\leq C'(\|u'\|_{a-1} \|\tilde{f}\|_1 + \|u'\|_1 \|\tilde{f}\|_{a-1} + \|u'\|_0 \|\tilde{f}\|_a). \end{aligned}$$

We can put the first two terms in the desired form by (A.10). □

Two inequalities in the next proposition are used in estimating P and Q .

Proposition A.5. *Let D be a convex domain in \mathbb{R}^m satisfying*

$$B_{\rho/c_0} \subset D \subset B_{c_0\rho}, \quad 0 < \rho \leq 3.$$

Let $a_ = 0$ for $0 \leq a \leq 1$ and $a_* = a$ for $a > 1$. Let $\|\cdot\|_a = \|\cdot\|_{C^a(D)}$. Then*

$$\begin{aligned} \rho^{a_*} \left\| \prod_{j=1}^m u_j \right\|_a &\leq C_{a,c_0} \sum_{j=1}^m \|u_j\|_a \prod_{i \neq j} \|u_i\|_0, \\ \rho^e \prod_{j=1}^m \|u\|_{d_j+b_j} &\leq C_{a,b,c,c_0} \left(\prod_{j=1}^m \|u_j\|_{d_j+a_j} + \prod_{j=1}^m \|u_j\|_{d_j+c_j} \right), \end{aligned}$$

where $e = (b_1 + d_1 - [d_1]) + \dots + (b_m + d_m - [d_m])$, a_j, c_j, d_j are non-negative real numbers, and (b_1, \dots, b_m) is in the convex hull of (a_1, \dots, a_m) and (c_1, \dots, c_m) .

Proof. The first estimate is trivial if $a \leq 1$. So assume that $a > 1$. We know that D is convex and $B_{\rho/c_0} \subset D \subset B_{c_0\rho}$. Thus D has the cone property if $1 \leq \rho \leq \rho_0$, where the cone of fixed size with vertex at $x \in \overline{D}$ can be found from the convex hull of x and B_{ρ/c_0} . Assume now that $0 < \rho < 1$. Consider the isotropic dilation $S_\rho(x) = \rho x$. Then, $D_* = S_\rho^{-1}D$ has the cone property. Since $0 < \rho < 1$, we have

$$\rho^a \|u\|_a \leq \|u \circ S_\rho\|_{D_*,a} \leq \|u\|_a. \tag{A.11}$$

By Proposition A.4 (ii) we obtain

$$\rho^a \left\| \prod_{j=1}^m u_j \right\|_a \leq \left\| \prod_{j=1}^m u_j \circ S_\rho \right\|_{D_*,a} \leq C \sum_{j=1}^m \|u_j \circ S_\rho\|_{D_*,a} \prod_{i \neq j} \|u_i \circ S_\rho\|_{D_*,0}.$$

Using (A.11), we get the first inequality easily. Let $l_j \leq [d_j]$ be any non-negative integers. By (A.11) and Proposition A.4 (iii), we get

$$\begin{aligned} \rho^e \prod_{j=1}^m \|\partial^{l_j} u_j\|_{b_j+d_j-[d_j]} &\leq \prod_{j=1}^m \|(\partial^{l_j} u_j) \circ S_\rho\|_{D_*,b_j+d_j-[d_j]} \\ &\leq C \left(\prod_{j=1}^m \|(\partial^{l_j} u_j) \circ S_\rho\|_{D_*,a_j+d_j-[d_j]} + \prod_{j=1}^m \|(\partial^{l_j} u_j) \circ S_\rho\|_{D_*,c_j+d_j-[d_j]} \right) \\ &\leq C \left(\prod_{j=1}^m \|u_j\|_{l_j+a_j+d_j-[d_j]} + \prod_{j=1}^m \|u_j\|_{l_j+c_j+d_j-[d_j]} \right) \\ &\leq C \left(\prod_{j=1}^m \|u_j\|_{d_j+a_j} + \prod_{j=1}^m \|u_j\|_{d_j+c_j} \right). \end{aligned}$$

Summing over all non-negative integers $l_j \leq [d_j]$ gives us the second inequality. \square

One can also obtain other inequalities via dilation. The inequalities below are not directly used in this paper.

Proposition A.6. *Let a_* be as in Proposition A.5. Let $\rho, D, \|\cdot\|_a$ be as in Proposition A.5. Let a, b, a_j, b_j be nonnegative real numbers.*

- (i) $\rho^{c_*} \|u\|_{\lambda a+(1-\lambda)b} \leq C_{a,b} \|u\|_a^\lambda \|u\|_b^{1-\lambda}$ for $0 < \lambda < 1$, where $c_* = \lambda a + (1 - \lambda)b$.
- (ii) Let f be a map from D into $D' \subset \mathbb{R}^n$. Assume that D' is convex and $B_{c_0^{-1}\rho} \subset D' \subset B_{c_0\rho}$. Then

$$\begin{aligned} \rho^{a_*} \|u \circ f\|_a &\leq C_{a,c_0} \rho (\|u'\|_{D',a-1} \|f'\|_0^a + \|u'\|_{D',0} \|f'\|_{a-1}) + \|u\|_{D',0}, \quad a \geq 1, \\ \|u \circ f\|_a &\leq C_{c_0} \min(\|u\|_{D',1} \|f\|_a, \|u\|_{D',a} \|f\|_1^a) + \|u\|_{D',0}, \quad 0 \leq a \leq 1. \end{aligned}$$

(iii) Let $a_{**} = 0$ for $0 \leq a \leq 2$ and $a_{**} = a - 1$ for $a > 2$. Let f be a C^a map from D into \mathbb{R}^m . There exists $C_0 > 1$ such that if $\|f' - I\| < 1/C_0$ then

$$\rho^{a_{**}} \|f^{-1} - I\|_{f(D),a} \leq C_a \|f - I\|_a.$$

(iv) Let $f = I + \tilde{f}$ be a C^1 map from D into $D' \subset \mathbb{R}^m$ with $\|f'\| < C_0$. Assume that $D \cup D'$ is contained in a **convex** domain D'' satisfying $B_{\rho/C_0} \subset D'' \subset B_{C_0\rho}$. Then

$$\rho^{a_*} \|u \circ f - u\|_a \leq C_{a,C_0} (\|u'\|_{D'',a} \|\tilde{f}\|_0 + \|u'\|_{D'',0} \|\tilde{f}\|_a).$$

Proof. We may assume that $0 < \rho < 1$. Let $u_*(x) = u(\rho x)$, $f^*(x) = \rho^{-1}f(\rho x) = x + \tilde{f}^*(x)$, and $D_* = \rho^{-1}D$. Then D_* has the cone property.

(i) is immediate, by applying Proposition A.4 to u_* and by using (A.11).

(ii). The case $0 \leq a \leq 1$ is verified directly. Assume that $a > 1$. Applying Proposition A.4 to $u_* \circ f^*$, we get

$$\|u_* \circ f^*\|_{D_*,a} \leq C_a (\|\partial u_*\|_{D'_*,a-1} \|f^{*'}\|_{D_*,0}^a + \|\partial u_*\|_{D'_*,0} \|f^{*'}\|_{D_*,a-1}) + \|u_*\|_{D'_*,0}.$$

Since $f^{*'}(x) = f'(\rho x)$, then $\|f^{*'}\|_{D_*,b} \leq \|f'\|_{D,b}$ for $b \geq 0$. By (A.11), we also have

$$\begin{aligned} \|u_* \circ f^*\|_{D_*,a} &= \|(u \circ f) \circ S_\rho\|_{D_*,a} \geq \rho^a \|u \circ f\|_a, \\ \|\partial u_*\|_{D'_*,a-1} &= \|\rho(\partial u)_*\|_{D'_*,a-1} \leq \rho \|\partial u\|_{a-1}, \quad a \geq 1. \end{aligned}$$

Simplifying gives us (ii).

(iv). Let $\tilde{f} = f - I$. The case $0 \leq a \leq 1$ is verified directly. Assume that $a > 1$. Then

$$\|u_* \circ f^* - u_*\|_{D_*,a} \leq C (\|\tilde{f}^*\|_a \|\partial u_*\|_{D''_*,0} + \|\tilde{f}^*\|_0 \|\partial u_*\|_{D''_*,a}).$$

As in (ii), we can get (iv) by (A.11) and

$$\|\tilde{f}^*\|_{D'_*,a} = \rho^{-1} \|\tilde{f} \circ S_\rho\|_{D'_*,a} \leq \rho^{-1} \|\tilde{f}\|_a.$$

(iii). (A dilation would give us $a_{**} = a$.) Let $g = f^{-1} = I + \tilde{g}$. We have $\tilde{g} = -\tilde{f} \circ g$ and $\tilde{g}' = -\{\tilde{f}'(\mathbf{1} + \tilde{f}')^{-1}\} \circ g$. By Lemma A.1, $C^{-1}|x' - x| \leq |g(x') - g(x)| \leq C|x' - x|$. We get immediately

$$\|\tilde{g}\|_{f(D),a} \leq C \|\tilde{f}\|_a, \quad 0 \leq a \leq 2.$$

Assume that $k \geq 1$. Recall that by (A.7)-(A.8),

$$\partial^K \tilde{g} = \sum_{j_1 + \dots + j_l \leq k} \{p(\tilde{f}') \partial^{j_1} \tilde{f} \dots \partial^{j_l} \tilde{f}\} \circ g, \quad k = |K| - 1 \geq 0, \quad j_i = |J_i| - 1 \geq 0.$$

Set $1 + k = [a]$ and $\alpha = a - k - 1$. Computing the Hölder ratio of (A.8) and applying Proposition A.5 with $d_j = 1$, we obtain (iii) from

$$\|\partial^K \tilde{g}\|_\alpha \leq C \sum_{j_1 + \dots + j_l \leq k} \rho^{-j_1 - \dots - j_l - \alpha} \|\tilde{f}\|_{1+k+\alpha} \leq C' \rho^{-k-\alpha} \|\tilde{f}\|_{1+k+\alpha}. \quad \square$$

Remark A.7. If all norms in Proposition A.6 are replaced by the scalar-invariant norm $\|\cdot\|^*$, defined as

$$\|u\|_{D,a}^* = \|u \circ S_d\|_{S_d^{-1}D,a}$$

with d being the diameter of D , we can take $c_* = a_* = a_{**} = 0$ for the proposition. For the use of scalar-invariant norms to derive estimates on the Bochner-Martinelli-Koppelman formula for balls in \mathbb{C}^n , see [20].

B. The Henkin homotopy formula

Recall notation $z' = (z_1, \dots, z_{n-1})$, $z = (z', z^n)$ and $x = \pi(z) = (\text{Re } z, \text{Im } z')$.

In this appendix, we will derive the following version of Henkin’s homotopy formula.

Theorem B.1. *Let $M \subset \mathbb{C}^n$ be a graph $y^n = |z'|^2 + \hat{r}(x)$ over $D \subset \mathbb{R}^{2n-1}$. Assume that $0 < \rho < \rho_0 \leq 3$ and*

$$\overline{D_{\rho_0}} \subset D, \quad \hat{r}(0) = 0, \quad \partial \hat{r}(0) = 0, \quad \|\hat{r}\|_{\rho_0,2} = \|\hat{r}\|_{D_{\rho_0},2} < 1/C_0 \quad (\text{B.1})$$

with C_0 sufficiently large. Assume that $0 < \rho < \rho_0 \leq 3$. Let φ be a continuous tangential $(0, q)$ -form on $\overline{M_\rho}$. Assume that $\bar{\partial}_M \varphi$ is continuous as currents on M_ρ and admits a continuous extension on $\overline{M_\rho}$. If $0 < q < n - 2$, then on M_ρ and as currents

$$\varphi = \bar{\partial}_M(P_0 + P_1)\varphi + (Q_0 + Q_1)\bar{\partial}_M\varphi \quad \text{mod } \bar{\partial}r,$$

where P_0, P_1, Q_0, Q_1 are defined by (3.5).

Recall that $M_\rho = M \cap \{|x^n|^2 + y^n < \rho^2\}$ and $D_\rho = \{x \in D : |x^n|^2 + y^n < \rho^2\}$. When M is strictly convex, see Henkin [7] for the proof. Our proof will follow [21] via Stokes’ theorem.

Set $r = -y^n + |z'|^2 + \hat{r}(x)$ and define

$$F(\zeta, z) = r_z \cdot (\zeta - z) + \frac{1}{2} \sum_{1 \leq j,k \leq n} r_{z^j z^k} (\zeta^j - z^j)(\zeta^k - z^k), \quad (\text{B.2})$$

$$S_t(z) = \pi\{\zeta \in M : |F(\zeta, z)| = t\}, \quad S'_t(\zeta) = \pi\{z \in M : |F(\zeta, z)| = t\}.$$

Note that $r_{z^j z^k} = \hat{r}_{z^j z^k}$.

Lemma B.2. *Let $n \geq 2$. Let M be as in Theorem B.1. There exists $C_1 > 1$ satisfying the following.*

- (i) *If $0 < t < (\rho_0 - \rho)^2/C_1$ and $z \in M_\rho$, then $S_t(z)$ and $S'_t(z)$ are compact subsets of D_{ρ_0} .*
- (ii) *Assume that $\hat{r} \in \mathcal{C}^3(\overline{D_{\rho_0}})$, $z \in M_\rho$ and $0 < t < C_1^{-1} \min\{(\rho_0 - \rho)^2, (1 + \|\hat{r}\|_{\rho_0,3})^{-1}\}$. Then $S_t(z)$ and $S'_t(z)$ are smooth and of classes \mathcal{C}^3 and \mathcal{C}^1 , respectively.*

Proof. Set $\epsilon = \|\hat{f}\|_{\rho_0,2}$. (i) By Lemma 5.1, D_{ρ_0} is convex and $\text{dist}(\partial D_{\rho_0}, \partial D_{\rho}) \geq (\rho_0 - \rho)/C$. On $M_{\rho_0} \times M_{\rho_0}$, $|r_z \cdot (\zeta - z)| \geq |\zeta - z|^2/C$ by Lemma 6.3 and $|\hat{r}_{z^j z^k}| \leq C\epsilon$. Hence, (i) follows from

$$|F(\zeta, z)| \geq |\zeta - z|^2/(2C). \tag{B.3}$$

(ii) Fix $z \in M_{\rho_0}$. Let $\zeta_*^n = -2ir_z(\zeta - z)$. By Lemma 6.1, $\xi_* = (\text{Re } \zeta_*, \text{Im } \zeta'_*)$ are coordinates of M_{ρ_0} and

$$\xi_*^n = \text{Re}(-2ir_z \cdot (\zeta - z)) = \xi^n - x^n + 2 \text{Im}(\bar{z}' \cdot \zeta') + \text{Im}\{2\hat{r}_z \cdot (\zeta - z)\}, \tag{B.4}$$

$$\eta_*^n = \text{Im}(-2ir_z \cdot (\zeta - z)) = |\zeta'_*|^2 + \sum_{|I|=2} \mathcal{R}_*^2 \hat{f}(\xi, x) \xi_*^I, \tag{B.5}$$

where \mathcal{R}_*^I is defined by (6.5). Write $u = \text{Re}(-2iF(\zeta, z))$ and $v = \text{Im}(-2iF(\zeta, z))$. We have

$$\begin{aligned} -2iF(\zeta, z) &= -2ir_z \cdot (\zeta - z) - i \sum_{1 \leq j, k \leq n} r_{z^j z^k} (\zeta^j - z^j)(\zeta^k - z^k), \\ u(\xi_*) &= \xi_*^n + \tilde{u}(\xi_*), \quad \tilde{u}(\xi_*) = \sum_{|I|=2} a_I \circ \Psi^{-1}(\xi_*, x) \xi_*^I, \end{aligned} \tag{B.6}$$

$$v(\xi_*) = |\zeta'_*|^2 + \tilde{v}(\xi_*), \quad \tilde{v}(\xi_*) = \sum_{|I|=2} b_I \circ \Psi^{-1}(\xi_*, x) \xi_*^I, \tag{B.7}$$

where $|a_I| + |b_I| \leq C\|\hat{f}\|_{\rho_0,2} \leq C'\epsilon$ and $|\partial^1 a_I(\zeta, z)| + |\partial^1 b_I(\zeta, z)| \leq C\|\hat{f}\|_{\rho_0,3}$. Suppose that $|F(\zeta, z)| = t$ and

$$t^{1/2}\|\hat{f}\|_{\rho_0,3} < 1/C_0 \tag{B.8}$$

and C_0 is sufficiently large. To show that $S_t(z)$ is smooth, we need to verify that $d_{\zeta'_*, \xi_*^n}(u^2 + v^2) \neq 0$ when $u^2 + v^2 = t^2$. By (B.3)-(B.4) we know that

$$|\zeta - z|/C \leq |\xi_*| \leq C|\zeta - z| \leq C't^{1/2}.$$

By (B.3) and (B.8), we obtain $|\zeta - z|\|\hat{f}\|_{\rho_0,3} \leq 1/C_0$ and

$$|\partial^1 \tilde{u}| + |\partial^1 \tilde{v}| \leq C(\|\hat{f}\|_{\rho_0,3}|\zeta - z|^2 + \epsilon|\zeta - z|) \leq C'(C_0^{-1} + \epsilon)|\xi_*|.$$

Assume that $d_{\zeta'_*, \xi_*^n}(u^2 + v^2) = 0$. Then

$$(\xi_*^n + \tilde{u})(1 + \tilde{u}_{\xi_*^n}) + v v_{\xi_*^n} = 0, \quad u \tilde{u}_{\zeta'_*} + v(\bar{\zeta}'_* + \tilde{v}_{\zeta'_*}) = 0.$$

From the first identity we get $|\xi_*^n| \leq C(|\tilde{u}| + |v|) \leq C'(|\zeta'_*|^2 + |\xi_*^n|^2)$. Therefore, for $t < 1/C$, $|\xi_*^n| \leq C|\zeta'_*|^2$. Now, $|v| \geq |\zeta'_*|^2 - C\epsilon(|\zeta'_*|^2 + |\xi_*^n|^2) \geq \frac{1}{2}|\zeta'_*|^2$ and $|u| \leq C|\zeta'_*|^2$. By Lemma 5.2, $|\partial_{\xi_*^n}^1 \xi| \leq C$, and by (B.7) and (B.8),

$$\begin{aligned} |\tilde{v}_{\zeta'_*}| &\leq |\zeta'_*|/2, \quad |\bar{\zeta}'_* + \tilde{v}_{\zeta'_*}| \geq |\zeta'_*|/2, \\ |\zeta'_*|^3/4 &\leq |v(\bar{\zeta}'_* + \tilde{v}_{\zeta'_*})| = |u \tilde{u}_{\zeta'_*}| \leq C|\zeta'_*|^2(\epsilon + C_0^{-1})|\zeta'_*|. \end{aligned}$$

Hence $\zeta'_* = \xi^n_* = 0$, when C_0^{-1}, ϵ are sufficiently small. This shows that $t = 0$, a contradiction. It is clear that for a fixed z , $u^2 + v^2$ is a function of class C^3 in (ζ', ξ^n) . This shows that $S_t(z)$ is smooth and of class C^3 .

For a fixed $\zeta \in M_\rho$, (ζ'_*, ξ^n_*) still form a coordinate system of class C^2 for M . We have the same formulae (B.4)-(B.5), only to vary $z \in M_{\rho_0}$; see Remark 6.2. Thus (B.6)-(B.7) are still valid, where ξ is fixed and z varies in M_{ρ_0} . The same argument shows that $S'_t(\zeta)$ is smooth and of class C^1 . □

Let us recall from Section 3

$$\Omega_{0,q}^{+-} = \frac{(r_\zeta - r_z) \cdot d\zeta \wedge r_z \cdot d\zeta \wedge (\bar{\partial}_\zeta \partial_\zeta r)^{n-2-q} \wedge (\bar{\partial}_z r_z \wedge d\zeta)^q}{(r_\zeta \cdot (\zeta - z))^{n-1-q} (r_z \cdot (\zeta - z))^{q+1}}.$$

The following computation is essential in Folland-Stein [3]. See also Romero [17].

Lemma B.3. *Let $n \geq 2$ and $0 \leq q \leq n - 1$. Let M satisfy (B.1) and $\hat{r} \in C^3$. Let $F(\zeta, z)$ be the Levi polynomial of r about $z \in M$. Let φ be a continuous tangential $(0, q)$ -form on M . Then*

$$\lim_{t \rightarrow 0} \int_{|F(\zeta, z)|=t, \zeta \in M} \varphi(\zeta) \wedge \Omega_{0,q}^{+-}(\zeta, z) = \frac{(2\pi i)^n}{2} \varphi(z) \pmod{\bar{\partial}r(z)}, \tag{B.9}$$

where the convergence is uniformly in z on each compact subset of M .

In the lemma, $|F(z, \cdot)| = t$ is oriented as the boundary of the domain $|F(z, \cdot)| < t$ in M on which $dV = d\xi^1 \wedge d\eta^1 \wedge \dots \wedge d\eta^{n-1} \wedge d\xi^n$ is the volume-form.

Here is an outline. Following [3], we will use the Levi polynomial $F(\zeta, z)$ to define new coordinates of M near z and compute each term in the kernel. On $T_z^{1,0}M$ there are two quadratic forms $h = \sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} t^j \bar{t}^k$, $A = \sum_{1 \leq j, k \leq n} r_{z^j z^k} t^j t^k$. We will express the kernel in h and A , with error terms. When we compute the residue, the error terms can be removed via non-isotropic dilation. This gives us a limit kernel (see (B.19) below) on a non-isotropic sphere. Roughly speaking, the limit kernel is expressed in h and A , but not in \bar{A} . For latter purpose we will use $-2iF(\zeta, z)$, instead of its linear part $-2ir_z \cdot (\zeta - z)$, as part of coordinates. Without \bar{A} , we remove A in the limit kernel by averaging. By a linear transformation, we reduce h to the identity, and compute the residue.

CHANGE NOTATION. Let $\zeta'_* = \zeta' - z'$ and

$$\zeta^n_* = -2iF(\zeta, z) = -2ir_z \cdot (\zeta - z) - i \sum_{1 \leq j, k \leq n} \hat{r}_{z^j z^k} (\zeta^j - z^j)(\zeta^k - z^k). \tag{B.10}$$

Then near $z \in M_{\rho_0}$, ζ'_*, ξ^n_* form coordinates of M_{ρ_0} . We will modify some earlier computations where the approximate Heisenberg transformation is used.

Applying the Taylor formula on convex domain $D_{\rho_0} \times \mathbb{R}$ and then letting $\zeta, z \in M_{\rho_0}$, we obtain

$$r(\zeta) - r(z) = 2 \operatorname{Re} F(\zeta, z) + \sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} (\zeta^j - z^j) \overline{(\zeta^k - z^k)} + E(\zeta, z),$$

$$E(\zeta, z) = o(|(\zeta' - z', \xi^n - x^n)|^2), \quad |\partial^1 E(\zeta, z)| \leq C \|\hat{r}\|_{\rho_0, 3} |(\zeta' - z', \xi^n - x^n)|^2.$$

With $z \in M_{\rho_0}$ the equation $r(\zeta) = 0$ becomes

$$-\eta_*^n + \sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} (\zeta^j - z^j) \overline{(\zeta^k - z^k)} + E(\zeta, z) = 0,$$

where $\zeta_n - z_n$, with $\zeta' - z' = \zeta_*$, is a local solution to (B.10). Replace $\zeta - z$ by the solution expressed in ζ_* . Solving for η_*^n from the new equation shows that for $(\zeta'_*, \xi_*^n) \in \pi \psi_z(M_{\rho_0})$

$$\eta_*^n = \sum_{1 \leq \alpha, \beta < n} h_{\alpha\bar{\beta}} \zeta_*^\alpha \bar{\zeta}_*^\beta + o(|\zeta'_*|^2) + O(|\xi_*^n| |\zeta'_*| + |\xi_*^n|^2). \tag{B.11}$$

Write $A(\zeta'_*, \xi_*^n) = o(k)$ if $t^{-k} A(t\zeta'_*, t^2\xi_*^n)$ tends to zero uniformly for $|\zeta'_*| + |\xi_*^n| < 1$ and small $|t|$, where A is a function or differential form. Write $A(\zeta'_*, \xi_*^n) = O(k)$ if $|t^{-k} A(t\zeta'_*, t^2\xi_*^n)| \leq C$. Thus $\zeta' - z' = O(1)$, $\zeta_*^n = O(2)$, $d\xi_*^n = O(1)$ and $d\xi_*^n = O(2)$. By (B.10)-(B.11), $\zeta^n - z^n = -\frac{r_{z'}}{r_{z^n}} \cdot \zeta'_* + O(2)$. The Levi matrix $(h_{\alpha\bar{\beta}})$ at z is determined by

$$\sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} (\zeta^j - z^j) \overline{(\zeta^k - z^k)} = \sum_{1 \leq \alpha, \beta < n} h_{\alpha\bar{\beta}} \zeta_*^\alpha \bar{\zeta}_*^\beta + o(2). \tag{B.12}$$

Explicitly, we have

$$h_{\alpha\bar{\beta}} = r_{z^\alpha \bar{z}^\beta} - r_{z^\alpha \bar{z}^n} \frac{r_{z^\beta}}{r_{z^n}} - \frac{r_{z^\alpha}}{r_{z^n}} r_{z^n \bar{z}^\beta} + r_{z^n \bar{z}^n} \frac{r_{z^\alpha}}{r_{z^n}} \frac{r_{z^\beta}}{r_{z^n}}. \tag{B.13}$$

We also have the holomorphic quadratic form

$$\sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} (\zeta^j - z^j) (\zeta^k - z^k) = \sum_{1 \leq \alpha, \beta < n} A_{\alpha\beta} \zeta_*^\alpha \zeta_*^\beta + o(2). \tag{B.14}$$

Here $A_{\alpha\beta} = A_{\beta\alpha}$. Set $h = (h_{\alpha\bar{\beta}})$ and $A = (A_{\alpha\beta})$. To simplify notation, we define

$$h(\zeta'_*, \bar{\zeta}'_*) = \sum_{1 \leq \alpha, \beta < n} h_{\alpha\bar{\beta}} \zeta_*^\alpha \bar{\zeta}_*^\beta, \quad A(\zeta'_*, \zeta'_*) = \sum_{1 \leq \alpha, \beta < n} A_{\alpha\beta} \zeta_*^\alpha \zeta_*^\beta.$$

We can write

$$\bar{\partial}_{\zeta_*} h(\zeta'_*, \bar{\zeta}'_*) = h(\zeta'_*, d\bar{\zeta}'_*), \quad \partial_{\zeta_*} A(\zeta'_*, \zeta'_*) = 2A(d\zeta'_*, \zeta'_*).$$

We will need to express the kernel in $h = (h_{\alpha\bar{\beta}})$ and $A = (A_{\alpha\beta})$, but not in $(\bar{A}_{\alpha\beta})$. This will play a crucial role, when we eliminate A later via averaging.

For the denominator of the kernel, using (B.10)-(B.11) we get

$$\begin{aligned} -2ir_z \cdot (\zeta - z) &= \zeta_*^n + i \sum_{1 \leq j, k \leq n} r_{z^j z^k} (\zeta^j - z^j) (\zeta^k - z^k) \\ &= \xi_*^n + i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(2). \end{aligned} \quad (\text{B.15})$$

Using $r_\zeta \cdot (\zeta - z) = r_z \cdot (\zeta - z) + (r_\zeta - r_z) \cdot (\zeta - z)$, we get

$$\begin{aligned} -2ir_\zeta \cdot (\zeta - z) &= -2ir_z \cdot (\zeta - z) - 2i \sum_{1 \leq j, k \leq n} r_{z^j z^k} (\zeta^j - z^j) (\zeta^k - z^k) \\ &\quad - 2i \sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} (\zeta^j - z^j) (\bar{\zeta}^k - \bar{z}^k) + o(2) \\ &= -2ir_z \cdot (\zeta - z) - 2i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(2) \quad (\text{by (B.12), (B.14)}) \\ &= \xi_*^n - i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(2). \quad (\text{by (B.15)}) \end{aligned}$$

We arrive at the basic relations

$$-2ir_\zeta \cdot (\zeta - z) = \xi_*^n - i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(2), \quad (\text{B.16})$$

$$-2ir_z \cdot (\zeta - z) = \xi_*^n + i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(2). \quad (\text{B.17})$$

We now compute the numerator of the kernel. Using the first-order expansion of r_ζ about z , we get

$$\begin{aligned} (r_\zeta - r_z) \cdot d\zeta &= \sum_{1 \leq j, k \leq n} \{r_{z^j z^k} (\zeta^k - z^k) + r_{z^j \bar{z}^k} (\bar{\zeta}^k - \bar{z}^k) + O(2)\} d\zeta^j \\ &= \partial_\zeta \sum_{1 \leq j, k \leq n} \left\{ \frac{1}{2} r_{z^j z^k} (\zeta^j - z^j) (\zeta^k - z^k) + r_{z^j \bar{z}^k} (\zeta^j - z^j) (\bar{\zeta}^k - \bar{z}^k) \right\} \\ &\quad + o(2). \end{aligned}$$

Note that for fixed z , $\zeta \rightarrow \zeta_*$ is holomorphic. So we can switch the above ∂_ζ to ∂_{ζ_*} and then restrict it to M , which gives us

$$(r_\zeta - r_z) \cdot d\zeta = h(d\zeta'_*, \bar{\zeta}'_*) + A(d\zeta'_*, \zeta'_*) + o(2). \quad (\text{B.18})$$

Recall that $\zeta_*^n = -2ir_z \cdot (\zeta - z) - i \sum_{1 \leq j, k \leq n} r_{z^j z^k} (\zeta^j - z^j) (\zeta^k - z^k)$. Applying ∂_ζ gives us

$$\begin{aligned} -2ir_z \cdot d\zeta &= \partial_\zeta \left\{ \zeta_*^n + i \sum_{1 \leq j, k \leq n} r_{z^j z^k} (\zeta^j - z^j) (\zeta^k - z^k) \right\} \\ &= \partial_\zeta \left\{ \zeta_*^n + i \sum_{1 \leq \alpha, \beta < n} A_{\alpha\beta} \zeta_*^\alpha \zeta_*^\beta + o(2) \right\} \quad (\text{by (B.14)}) \\ &= d\zeta_*^n + 2iA(d\zeta'_*, \zeta'_*) + o(2). \end{aligned}$$

Recall that z is fixed. So on M we have

$$\begin{aligned}\bar{\partial}_\zeta \partial_{\zeta'} r &= \bar{\partial} \partial r(\zeta) = \bar{\partial} \partial \left\{ \sum_{1 \leq j, k \leq n} r_{z^j \bar{z}^k} (\zeta^j - z^j) \overline{(\zeta^k - z^k)} + o(2) \right\} \\ &= \bar{\partial} \partial \left\{ \sum_{1 \leq \alpha, \beta < n} h_{\alpha\beta} \bar{\zeta}_*^\alpha \bar{\zeta}_*^{-\beta} \right\} + o(2) \quad \text{by (B.12)} \\ &= -h(d\zeta'_*, d\bar{\zeta}'_*) + o(2).\end{aligned}$$

On M , $\bar{\partial} r_{z^j} = \bar{\partial}_M r_{z^j} \bmod \bar{\partial} r(z)$ and $\bar{\partial} r_{z^j} = \sum_{\beta=1}^{n-1} (r_{z^j \bar{z}^\beta} - \frac{r_{z^\beta}}{r_{z^n}} r_{z^j \bar{z}^n}) d\bar{z}^\beta \bmod \bar{\partial} r(z)$. Recall that $\zeta^n - z^n = -r_{z^n}^{-1} r_{z'} \cdot \zeta'_* + O(2)$. Thus $d\zeta^n = -\frac{r_{z'}}{r_{z^n}} \cdot d\zeta'_* + O(2)$. Therefore,

$$\begin{aligned}(\bar{\partial} r_z) \wedge d\zeta &= \sum_{1 \leq \alpha, \beta < n} (r_{z^\alpha \bar{z}^\beta} - \frac{r_{z^\beta}}{r_{z^n}} r_{z^\alpha \bar{z}^n}) d\bar{z}^\beta \wedge d\zeta_*^\alpha + O(2) \\ &\quad - \sum_{1 \leq \alpha, \beta < n} (r_{z^n \bar{z}^\beta} - \frac{r_{z^\beta}}{r_{z^n}} r_{z^n \bar{z}^n}) d\bar{z}^\beta \wedge \frac{r_{z^\alpha}}{r_{z^n}} d\zeta_*^\alpha \quad \bmod \bar{\partial} r(z).\end{aligned}$$

Looking at (B.13), we see that

$$(\partial r_z) \wedge d\zeta = -h(d\zeta'_*, d\bar{z}') + O(2) \quad \bmod \bar{\partial} r(z).$$

We apply the non-isotropy dilation $T_t(\zeta'_*, \xi_*^n) = (t\zeta'_*, t^2\xi_*^n)$ and summarize the above as

$$\begin{aligned}t^{-2} T_t^* \{-2ir_\zeta \cdot (\zeta - z)\} &= \xi_*^n - i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(t^0), \\ t^{-2} T_t^* \{-2ir_z \cdot (\zeta - z)\} &= \xi_*^n + i(h(\zeta'_*, \bar{\zeta}'_*) + A(\zeta'_*, \zeta'_*)) + o(t^0), \\ t^{-2} T_t^* \{(r_\zeta - r_z) \cdot d\zeta\} &= h(d\zeta'_*, \bar{\zeta}'_*) + A(d\zeta'_*, \zeta'_*) + o(t^0), \\ t^{-2} T_t^* \{-2ir_z \cdot d\zeta\} &= d\xi_*^n + 2iA(d\zeta'_*, \zeta'_*) + o(t^0), \\ t^{-2} T_t^* \{\bar{\partial}_\zeta r_\zeta \wedge d\zeta\} &= -h(d\zeta'_*, d\bar{\zeta}'_*) + o(t^0), \\ t^{-1} T_t^* \{\bar{\partial}_z r_z \wedge d\zeta\} &= -h(d\zeta'_*, d\bar{z}') + O(t) \quad \bmod \bar{\partial} r(z).\end{aligned}$$

By (B.10) and (B.15), the subset of M defined by $2|F(\zeta, z)| = t^2$ has the form

$$S_t(z) = \left\{ (\zeta'_*, \xi_*^n) : |\xi_*^n + ih(\zeta'_*, \bar{\zeta}'_*) + o(2)| = t^2 \right\}.$$

Then we have a non-isotropic sphere

$$\lim_{t \rightarrow 0} T_t^{-1}(S_t) = \left\{ (\zeta'_*, \xi_*^n) : |\xi_*^n + ih(\zeta'_*, \bar{\zeta}'_*)| = 1 \right\} \stackrel{\text{def}}{=} S.$$

Set $N_*(\zeta'_*, \xi_*^n) = \xi_*^n + ih(\zeta'_*, \bar{\zeta}'_*)$. As t tends to 0, $t^q T_t^* \Omega_{(0,q)}^{+-} \bmod \bar{\partial}r(z)$ converges uniformly in a neighborhood of S to $\tilde{\Omega}$. Here $\tilde{\Omega}$ is

$$\frac{(h(d\zeta'_*, \bar{\zeta}'_*) + A(d\zeta'_*, \zeta'_*)) \wedge (d\zeta_*^n + 2iA(d\zeta'_*, \zeta'_*)) \wedge h(d\zeta'_*, d\bar{\zeta}'_*)^{n-q-2} \wedge h(d\zeta'_*, d\bar{z}'_*)^q}{-(2i)^{1-n} \bar{N}_*^{n-q-1} N_*^{q+1} (1 - i\bar{N}_*^{-1} \zeta'_* A \zeta_*'^T)^{n-q-1} (1 + iN_*^{-1} \zeta'_* A \zeta_*'^T)^{q+1}}. \quad (\text{B.19})$$

Here $N_* = N_*(\zeta'_*, \xi_*^n)$. Note that except at the origin, the above form is smooth in the (ζ'_*, ξ_*^n) -space. Therefore, for $\varphi(\zeta) = \sum_{|I|=q} \varphi_{\bar{T}}^I(\zeta) d\bar{\zeta}'^I$

$$\begin{aligned} \int_{2|F(\zeta,z)|=t^2} \varphi(\zeta) \wedge \Omega_{0,q}^{+-}(\zeta, z) &= \int_{(\zeta'_*, \xi_*^n) \in S_t(z)} \varphi(\zeta(\zeta'_*, \xi_*^n)) \wedge \Omega_{0,q}^{+-}(\zeta(\zeta'_*, \xi_*^n), z) \\ &= \int_{(\zeta'_*, \xi_*^n) \in T_t^{-1} S_t(z)} T_t^* \left\{ \varphi(\zeta(\zeta'_*, \xi_*^n)) \wedge \Omega_{0,q}^{+-}(\zeta(\zeta'_*, \xi_*^n), z) \right\} \\ &= \sum_{|I|=q} \varphi_{\bar{T}}^I(z) \int_{(\zeta'_*, \xi_*^n) \in T_t^{-1} S_t(z)} T_t^* \left\{ d\bar{\zeta}'^I \wedge \Omega_{0,q}^{+-}(\zeta(\zeta'_*, \xi_*^n), z) \right\} \bmod \bar{\partial}r(z). \end{aligned}$$

We obtain

$$\lim_{t \rightarrow 0} \int_{2|F(\zeta,z)|=t^2} \varphi(\zeta) \wedge \Omega_{0,q}^{+-}(\zeta, z) = \sum_{|I|=q} \varphi_{\bar{T}}^I(z) \int_S d\bar{\zeta}'^I \wedge \tilde{\Omega}(\zeta'_*, \xi_*^n, z) \bmod \bar{\partial}r(z).$$

Return to $\tilde{\Omega}$, defined by (B.19). We are ready to remove A via an averaging. By (B.1), $|A|$ is small. Express $(1 - i\bar{N}_*^{-1} \zeta'_* A \zeta_*'^T)^{-(n-q-1)} (1 + iN_*^{-1} \zeta'_* A \zeta_*'^T)^{-(q+1)}$ as a convergent power series in $\bar{N}_*^{-1} \zeta'_* A \zeta_*'^T$ and $N_*^{-1} \zeta'_* A \zeta_*'^T$. Note that $N_*(\xi_*^n)$ is invariant under the rotation $e_\theta: (\zeta'_*, \xi_*^n) \rightarrow (e^{i\theta} \zeta'_*, \xi_*^n)$. Using $|I| = q$, we can verify that

$$\begin{aligned} \int_{(\zeta'_*, \xi_*^n) \in S} d\bar{\zeta}'^I \wedge \tilde{\Omega}(\zeta'_*, \xi_*^n, z) &= \frac{1}{2\pi} \int_0^{2\pi} d\theta \int_{(\zeta'_*, \xi_*^n) \in e_\theta S} d\bar{\zeta}'^I \wedge \tilde{\Omega}(\zeta'_*, \xi_*^n, z) \\ &= \frac{1}{2\pi} \int_0^{2\pi} d\theta \int_{(\zeta'_*, \xi_*^n) \in S} e_\theta^* \{ d\bar{\zeta}'^I \wedge \tilde{\Omega} \} \\ &= \frac{1}{2\pi} \int_{(\zeta'_*, \xi_*^n) \in S} \int_{\theta=0}^{2\pi} e_\theta^* \{ d\bar{\zeta}'^I \wedge \tilde{\Omega} \} d\theta. \end{aligned}$$

Therefore $\int_{(\zeta'_*, \xi_*^n) \in S} d\bar{\zeta}'^I \wedge \tilde{\Omega}(\zeta'_*, \xi_*^n, z) = \int_{(\zeta'_*, \xi_*^n) \in S} d\bar{\zeta}'^I \wedge \Omega'(\zeta'_*, \xi_*^n, z)$, where

$$\Omega'(\zeta'_*, \xi_*^n, z) = \frac{d\bar{\zeta}'^I h \bar{\zeta}'^T \wedge d\zeta_*^n \wedge (d\zeta'_* \wedge h d\bar{\zeta}'^T)^{n-q-2} \wedge (d\zeta'_* \wedge h d\bar{z}'^T)^q}{-(2i)^{1-n} \bar{N}_*^{n-q-1} (\xi_*^n) N_*^{q+1} (\xi_*^n)}.$$

In eliminating A , we have used the fact that there are no terms $\overline{\zeta_*^\alpha} \overline{\zeta_*^\beta}$ in (B.16)-(B.17).

Take a linear transformation $\hat{\zeta}' = U(\zeta'_*)$, $\hat{\xi}^n = \xi_*^n$ such that

$$h(\zeta'_*, \overline{\zeta_*}) = |\hat{\zeta}'|^2, \quad \hat{\zeta}^n = \hat{\xi}^n + i|\hat{\zeta}'|^2.$$

Let $\hat{z}' = U(z')$. Under the new coordinates and on $|\hat{\zeta}^n| = 1$, $\Omega'(\zeta'_*, \xi_*^n, z)$ becomes

$$\begin{aligned} \widehat{\Omega}(\hat{\zeta}', \hat{\xi}^n, \hat{z}') &= \frac{(\overline{\hat{\zeta}'} \cdot d\hat{\zeta}') \wedge d\hat{\zeta}^n \wedge (d\hat{\zeta}' \wedge d\overline{\hat{\zeta}'})^{n-q-2} \wedge (d\hat{\zeta}' \wedge d\overline{\hat{z}'})^q}{-(2i)^{1-n} (\hat{\zeta}^n)^{n-q-1} (\hat{\zeta}^n)^{q+1}} \\ &= (\overline{\hat{\zeta}'} \cdot d\hat{\zeta}') \wedge d\mu(\hat{\zeta}^n) \wedge (d\hat{\zeta}' \wedge d\overline{\hat{\zeta}'})^{n-q-2} \wedge (d\hat{\zeta}' \wedge d\overline{\hat{z}'})^q. \end{aligned}$$

Here

$$\mu(\hat{\zeta}^n) = \begin{cases} -(2i)^{n-1} (n-2q-1)^{-1} (\hat{\zeta}^n)^{n-2q-1}, & n-2q \neq 1, \\ -(2i)^{n-1} \log \hat{\zeta}^n, & n-2q = 1. \end{cases}$$

Note that $\text{Im} \hat{\zeta}^n \geq 0$ and hence $\mu(\hat{\zeta}^n)$ is smooth and single-valued on $|\hat{\zeta}^n| = 1$. To remove the differential on μ , we apply Stokes' theorem and get

$$\begin{aligned} c_q d\overline{\hat{z}'^I} &\stackrel{\text{def}}{=} \int_{|\hat{\zeta}^n|=1} d\overline{\hat{\zeta}'^I} \wedge (\overline{\hat{\zeta}'} \cdot d\hat{\zeta}') \wedge d\mu(\hat{\zeta}^n) \wedge (d\hat{\zeta}' \wedge d\overline{\hat{\zeta}'})^{n-q-2} \wedge (d\hat{\zeta}' \wedge d\overline{\hat{z}'})^q \\ &= - \int_{|\hat{\zeta}^n|=1} \mu(\hat{\zeta}^n) (d\hat{\zeta}' \wedge d\overline{\hat{\zeta}'})^{n-q-1} \wedge d\overline{\hat{\zeta}'^I} \wedge (d\hat{\zeta}' \wedge d\overline{\hat{z}'})^q. \end{aligned}$$

Now, the differential form is a multiple of the volume-form on \mathbb{C}^{n-1} (the $\hat{\zeta}'$ -space). The projection from $|\hat{\zeta}^n| = 1$ to the ball $|\hat{\zeta}'| \leq 1$ is two-to-one, branched over $|\hat{\zeta}'| = 1$. Recall that $dV(\zeta) = d\xi^1 \wedge d\eta^1 \wedge \cdots \wedge d\eta^{n-1} \wedge d\xi^n$ defines the orientation of M and $(\zeta', \xi^n) \rightarrow (\zeta'_*, \xi_*^n)$ preserves the orientation. Since $S: |\hat{\zeta}^n| = 1$ is obtained via the dilation of the boundary of $|F(z, \cdot)| < t$, S must be oriented as the boundary of $|\hat{\zeta}^n| < 1$ on which $dV(\hat{\zeta})$ is the volume-form. After considering the orientation, we conclude that

$$c_q d\overline{\hat{z}'^I} = \int_{|\hat{\zeta}'| \leq 1} (\mu(\hat{\zeta}_-^n) - \mu(\hat{\zeta}_+^n)) (d\hat{\zeta}' \wedge d\overline{\hat{\zeta}'})^{n-q-1} \wedge d\overline{\hat{\zeta}'^I} \wedge (d\hat{\zeta}' \wedge d\overline{\hat{z}'})^q.$$

Here $dV(\hat{\zeta}') = d\hat{\xi}^1 \wedge d\hat{\eta}^1 \wedge \cdots \wedge d\hat{\eta}^{n-1}$ is the volume-form on the $\hat{\zeta}'$ -space, and

$$\hat{\zeta}_+^n = (1 - |\hat{\zeta}'|^4)^{1/2} + i|\hat{\zeta}'|^2, \quad \hat{\zeta}_-^n = -(1 - |\hat{\zeta}'|^4)^{1/2} + i|\hat{\zeta}'|^2.$$

To rewrite the integrand, introduce variables $\xi = (\xi_1, \dots, \xi_q)$, η, x, y so that

$$d\overline{\hat{\zeta}'^I} = d\overline{\xi^I}, \quad d\hat{\zeta}' \wedge d\overline{\hat{\zeta}'} = d\xi \wedge d\overline{\xi} + d\eta \wedge d\overline{\eta}, \quad d\hat{\zeta}' \wedge d\overline{\hat{z}'} = d\xi \wedge d\overline{x} + d\eta \wedge d\overline{y}.$$

Then $(d\xi \wedge d\bar{\xi} + d\eta \wedge d\bar{\eta})^{n-q-1} \wedge d\bar{\xi}^l \wedge (d\xi \wedge d\bar{x} + d\eta \wedge d\bar{y})^q$ equals

$$\begin{aligned} & (d\eta \wedge d\bar{\eta})^{n-q-1} \wedge d\bar{\xi}^l \wedge (d\xi \wedge d\bar{x} + d\eta \wedge d\bar{y})^q \\ &= (d\eta \wedge d\bar{\eta})^{n-q-1} \wedge d\bar{\xi}^l \wedge (d\xi \wedge d\bar{x})^q \\ &= (-1)^q (d\eta \wedge d\bar{\eta})^{n-q-1} \wedge (d\xi \wedge d\bar{\xi})^q \wedge d\bar{x}^l. \end{aligned}$$

The last term equals $(-1)^q \binom{n-1}{q}^{-1} (d\xi \wedge d\bar{\xi} + d\eta \wedge d\bar{\eta})^{n-1} \wedge d\bar{x}^l$. Therefore

$$c_q d\bar{z}^l = (-1)^q \binom{n-1}{q}^{-1} \int_{|\hat{\xi}'| < 1} (\mu(\hat{\xi}_-^n) - \mu(\hat{\xi}_+^n)) (d\hat{\xi}' \wedge d\bar{\xi}')^{n-1} \wedge d\bar{z}^l.$$

We now compute c_q . Using the polar coordinates, we get

$$\begin{aligned} c_q &= (-1)^q \binom{n-1}{q}^{-1} (n-1)! (-2i)^{n-1} \int_{|\hat{\xi}'| < 1} (\mu(\hat{\xi}_-^n) - \mu(\hat{\xi}_+^n)) dV(\hat{\xi}') \\ &= \tilde{c}_q \int_0^1 (\mu((1-r^4)^{1/2} + ir^2) - \mu(-(1-r^4)^{1/2} + ir^2)) r^{2n-3} dr \\ &= \frac{\tilde{c}_q}{2(n-1)} \int_{r=0}^1 r^{2n-2} d\{\mu((1-r^4)^{1/2} + ir^2) - \mu(-(1-r^4)^{1/2} + ir^2)\}. \end{aligned}$$

Note that the last identity is obtained by integration by parts. Here

$$\tilde{c}_q = (-1)^{q+1} \binom{n-1}{q}^{-1} (n-1)! (-2i)^{n-1} \sigma_{n-1}, \quad \sigma_{n-1} = \frac{2\pi^{n-1}}{(n-2)!}.$$

Letting $r^2 = s$ and $z = (1-s^2)^{1/2} + is$, we get

$$\begin{aligned} c_q &= \frac{\tilde{c}_q}{2(n-1)} \int_{s=-1}^1 s^{n-1} d\{\mu((1-s^2)^{1/2} + is)\} \\ &= \frac{\tilde{c}_q}{4(n-1)} \int_{|z|=1} (\operatorname{Im} z)^{n-1} d\mu(z) = \frac{-(2i)^{n-1} \tilde{c}_q}{4(n-1)} \int_{|z|=1} (\operatorname{Im} z)^{n-1} z^{n-2q-2} dz. \end{aligned}$$

A simple residue computation yields $c_q = \frac{1}{2} (2\pi i)^n$ for $0 \leq q \leq n-1$. The proof of Lemma B.3 is complete.

We need the following lemma from [7], where $|F(\zeta, z)| = t$ is replaced by $|\zeta - z| = t$ and only $r \in \mathbb{C}^2$ is needed.

Lemma B.4. *Let $1 \leq q \leq n-1$. Let M, ρ_0 be as in Lemma B.3. Assume that r is of class \mathcal{C}^3 . Let φ be a continuous $(0, q)$ -form on M_{ρ_0} . If $0 < \rho < \rho_0$ and $t > 0$ is sufficiently small, then on M_ρ , in the sense of currents and modulo $\bar{\partial}r(z)$,*

$$\int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \bar{\partial}_z \Omega_{(0, q-1)}^{+-}(\zeta, z) = (-1)^{q-1} \bar{\partial}_b \int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z).$$

Proof. For the convenience of the reader, we reproduce the proof in [7]. Since $r \in \mathcal{C}^3$, for t sufficiently small, both integrals are continuous on M_ρ . To verify the identity in the sense of currents, let ψ be a smooth $(n, n-1-q)$ -form with compact support in M_ρ . Interchanging the order of integration, we have

$$\begin{aligned} \mathcal{I}(z) &= \int_{z \in M} \int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \bar{\partial}_z \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \psi(z) \\ &= - \int_{\zeta \in M} \int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \bar{\partial}_z \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \psi(z) \quad (\text{by Fubini}) \\ &= \int_{\zeta \in M} \int_{F(\zeta, z) > t} \psi(z) \wedge \bar{\partial}_z \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \varphi(\zeta). \end{aligned}$$

Applying Stokes' theorem yields

$$\begin{aligned} \mathcal{I}(z) &= (-1)^q \int_{\zeta \in M} \int_{F(\zeta, z) > t} \bar{\partial}_z \psi(z) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \varphi(\zeta) \\ &\quad + (-1)^{q+1} \int_{\zeta \in M} \int_{F(\zeta, z) = t} \psi(z) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \varphi(\zeta) \\ &= \int_{\zeta \in M} \int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \bar{\partial}_z \psi(z) \\ &\quad + (-1)^{q+1} \int_{\zeta \in M} \int_{F(\zeta, z) = t} \varphi(\zeta) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \psi(z). \end{aligned}$$

Interchanging the order of integration in both terms yields

$$\begin{aligned} \mathcal{I}(z) &= - \int_{z \in M} \int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \bar{\partial}_z \psi(z) \\ &\quad + (-1)^{q+1} \int_{z \in M} \int_{F(\zeta, z) = t} \varphi(\zeta) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \psi(z) \\ &= - \int_{z \in M} \int_{F(\zeta, z) > t} \varphi(\zeta) \wedge \Omega_{(0, q-1)}^{+-}(\zeta, z) \wedge \bar{\partial}_z \psi(z). \end{aligned}$$

Here the second last term vanishes by counting total degree in ζ , which equals $q + 2n - q - 1 > 2n - 2$. \square

As in [21], the above two lemmas can be used to derive the Henkin homotopy formula for $(0, q)$ -forms on \overline{M}_ρ via Stokes' theorem, when $r \in \mathcal{C}^3$ satisfies (B.1) and $\varphi \in \mathcal{C}^1(\overline{M}_{\rho_0})$. Here are the details. Let $M^t(z) = M_{\rho_0} \cap \{\zeta : |F(\zeta, z)| > t\}$. We will use $\bar{\partial}_\zeta \Omega_{(0, q)}^{+-} + \bar{\partial}_z \Omega_{(0, q)}^{+-} = 0$ for $\zeta \neq z$ and $1 \leq q \leq n-2$; see [21]. For

$z \in M_\rho$ and t sufficiently small,

$$\begin{aligned} & \int_{\partial M^t(z)} \varphi(\zeta) \wedge \Omega_{(0,q)}^{+-}(\zeta, z) \\ &= \int_{M^t(z)} \bar{\partial}_\zeta \varphi \wedge \Omega_{(0,q)}^{+-} + (-1)^q \int_{M^t(z)} \varphi(\zeta) \wedge \bar{\partial}_\zeta \Omega_{(0,q)}^{+-} \\ &= \int_{M^t(z)} \bar{\partial}_\zeta \varphi \wedge \Omega_{(0,q)}^{+-}(\zeta, z) + (-1)^{q-1} \int_{M^t(z)} \varphi(\zeta) \wedge \bar{\partial}_z \Omega_{(0,q-1)}^{+-}(\zeta, z). \end{aligned}$$

Using Lemmas B.3-B.4 and letting $t \rightarrow 0$, we obtain, modulo $\bar{\partial}r(z)$,

$$\int_{\partial M_{\rho_0}} \varphi(\zeta) \wedge \Omega_{(0,q)}^{+-}(\zeta, z) = c_0 \varphi(z) + \int_{M_{\rho_0}} \bar{\partial}_\zeta \varphi \wedge \Omega_{(0,q)}^{+-} + \bar{\partial}_b \int_{M_{\rho_0}} \varphi(\zeta) \wedge \Omega_{(0,q-1)}^{+-}.$$

Now assume that $0 < q < n-2$. Then $-\Omega_{(0,q)}^{+-} = \bar{\partial}_\zeta \Omega_{(0,q)}^{0+-} + \bar{\partial}_z \Omega_{(0,q-1)}^{0+-}$; see [21].

We get that modulo $\bar{\partial}r(z)$

$$\begin{aligned} & \int_{\partial M_{\rho_0}} \varphi(\zeta) \wedge \Omega_{(0,q)}^{+-}(\zeta, z) \\ &= - \int_{\partial M_{\rho_0}} \varphi(\zeta) \wedge \bar{\partial}_\zeta \Omega_{(0,q)}^{0+-} - \int_{\partial M_{\rho_0}} \varphi(\zeta) \wedge \bar{\partial}_z \Omega_{(0,q-1)}^{0+-} \\ &= (-1)^q \int_{\partial M_{\rho_0}} \bar{\partial}_b \varphi(\zeta) \wedge \Omega_{(0,q)}^{0+-} - (-1)^q \bar{\partial}_b \int_{\partial M_{\rho_0}} \varphi(\zeta) \wedge \Omega_{(0,q-1)}^{0+-}. \end{aligned}$$

Therefore, modulo $\bar{\partial}r(z)$ and as currents,

$$\begin{aligned} c_0 \varphi(z) &= \bar{\partial}_b \left\{ - \int_{M_{\rho_0}} \varphi(\zeta) \wedge \Omega_{(0,q-1)}^{+-}(\zeta, z) - (-1)^q \int_{\partial M_{\rho_0}} \varphi(\zeta) \wedge \Omega_{(0,q-1)}^{0+-}(\zeta, z) \right\} \\ &\quad + \left\{ - \int_{M_{\rho_0}} \bar{\partial}_b \varphi(\zeta) \wedge \Omega_{(0,q)}^{+-}(\zeta, z) + (-1)^q \int_{\partial M_{\rho_0}} \bar{\partial}_b \varphi(\zeta) \wedge \Omega_{(0,q)}^{0+-}(\zeta, z) \right\} \\ &= -\bar{\partial}_b \left\{ \int_{M_{\rho_0}} \Omega_{(0,q-1)}^{+-}(\zeta, z) \wedge \varphi(\zeta) + \int_{\partial M_{\rho_0}} \Omega_{(0,q-1)}^{0+-}(\zeta, z) \wedge \varphi(\zeta) \right\} \\ &\quad - \left\{ \int_{M_{\rho_0}} \Omega_{(0,q)}^{+-}(\zeta, z) \wedge \bar{\partial}_b \varphi(\zeta) + \int_{\partial M_{\rho_0}} \Omega_{(0,q)}^{0+-}(\zeta, z) \wedge \bar{\partial}_b \varphi(\zeta) \right\}. \end{aligned}$$

We now derive the homotopy formula by reducing the regularity condition. Notice that when M is \mathcal{C}^2 and strictly convex it is proved by Henkin [21].

Fix a tangential $(0, q)$ -form φ on M_{ρ_0} . Let $0 < q < n - 2$.

Case a) $r \in \mathcal{C}^2$ and $\varphi \in \mathcal{C}^1(\overline{M}_{\rho_0})$. Write $x = (\operatorname{Re} z, \operatorname{Im} z')$ and $\xi = (\operatorname{Re} \zeta, \operatorname{Im} \zeta')$ with $\zeta, z \in M_{\rho_0}$. We first consider the case when φ has compact support, say in M_{ρ_1} with $\rho_1 < \rho_0$. Then $P'_{M_{\rho_0}} \varphi$ has no boundary term. Recall that $\Psi(\xi, x) = (\tilde{\psi}_x(\xi), x) = (\xi_*, x)$ is defined by relations $\zeta'_* = \zeta' - z'$, $\zeta_*^n = -2ir_z(\zeta - z)$ and $\zeta, z \in M_{\rho_0}$. By (8.2)-(8.3) and (8.6), on D_{ρ_0}

$$P'_{0, M_{\rho_0}} \varphi(x) = \sum_{|I|=q-1} \mathcal{I}_{\overline{I}}(x) d\bar{z}'^I,$$

$$\mathcal{I}_{\overline{I}}(x) = \sum_{|K|=1} \sum_{|J|=q} \int_{B_{\rho_0}} \left((\varphi_{\overline{J}} \tilde{A}_{\overline{I}K}^{\overline{J}}) \circ \Psi^{-1} \cdot \hat{T}_1^{-a} \cdot \hat{T}_2^{-b} \right) (\xi_*, x) \cdot \hat{k}_{ab}^K(\xi_*) dV(\xi_*),$$

$$\varphi_{\overline{J}}(\xi, x) = \varphi_{\overline{J}}(\xi), \quad \hat{T}_1(\xi_*, x) = 1 + \sum_{|L|=2} C_L \circ \Psi^{-1}(\xi_*, x) \xi_*^L N_*^{-1}(\xi_*).$$

Here $\tilde{A}_{\overline{I}K}^{\overline{J}}$, C_L are functions of the form $\partial_*^2 r$, $\partial_*^2 \hat{r}$, respectively, defined in Section 8. An analogous formula holds for \hat{T}_2 . These functions depend only on derivatives of \hat{r} of order at most two. We take a sequence of \mathcal{C}^∞ functions \hat{r}^m converging to \hat{r} in \mathcal{C}^2 -norm on \overline{D}_{ρ_0} . Subtracting \hat{r}^m by its Taylor polynomial of order 1 about the origin, we may assume that $\hat{r}^m(0) = 0$ and $\partial \hat{r}^m(0) = 0$. In what follows we use the letter m to indicate dependence on \hat{r}^m . Let M^m be the graph $y^n = |z'|^2 + \hat{r}^m(x)$ over D_{ρ_0} . Let $\rho \in (\rho_1, \rho_0)$. There exists m_* such that for $m > m_*$

$$D_{\rho_1} \subset \overline{D}_\rho^m \subset D_{\rho_0}, \quad D_\rho^m \stackrel{\text{def}}{=} \{x \in D_{\rho_0} : |x|^2 + \hat{r}^m(x) < \rho^2\}.$$

Assume that $m > m_*$. Then by $\|\hat{r}^m\|_{D_{\rho_0}, 2} < 1/C_0$, D_ρ^m is strictly convex. Note that φ is still a tangential $(0, q)$ -form of M^m on D_{ρ_0} . By the homotopy formula for the smooth M^m ,

$$\varphi = \bar{\partial}_{M^m} P'_{M_\rho^m} \varphi + Q'_{M_\rho^m} \bar{\partial}_{M^m} \varphi. \quad (\text{B.20})$$

By the formula of \hat{T}_j , it is clear that $|\hat{T}_{j,m}| \geq 1/2$ for m sufficiently large and $0 \neq \xi_* \in B_{\rho} \cap \tilde{\psi}_x^m(D_\rho)$. Since $\varphi_{\overline{J}}$ has compact support, we have

$$| \{ (\varphi_{\overline{J}} \tilde{A}_{\overline{I}K}^{\overline{J}m}) \circ \Psi_m^{-1} \cdot \hat{T}_{1,m}^{-a} \cdot \hat{T}_{2,m}^{-b} \} (\xi_*, x) | \leq C, \quad \xi_* \neq 0.$$

Since $\hat{k}_{ab}^I \in L^1_{\text{loc}}$, by the dominated convergence theorem, $\mathcal{I}_{\overline{I}}^m$ converges to $\mathcal{I}_{\overline{I}}$ pointwise on \overline{D}_{ρ_1} as $m \rightarrow \infty$. Thus, $P'_{M_\rho^m} \varphi$ converges to $P'_{M_\rho} \varphi$ pointwise on \overline{D}_{ρ_1} .

By the \mathcal{C}^0 -estimate on P'_0 (see (9.3)), we know that $|P'_{M^m_\rho}\varphi| < C$ on $\overline{D}_{\rho_1} \subset \cap_m D^m_\rho$. Since φ is of class \mathcal{C}^1 , $\overline{\partial}_{M^m}\varphi$ converges to $\overline{\partial}_M\varphi$ uniformly on \overline{D}_{ρ_0} . Next, we fix an $(n, n - q - 1)$ -form $\psi = \sum_{|J|=n-q-1} \psi_J dz^1 \wedge \cdots \wedge dz^{n-1} \wedge \partial r(z) \wedge d\overline{z}^J$ of class \mathcal{C}^1 and of compact support in D_{ρ_1} . Then $\psi^m = \sum_{|J|=n-q-1} \psi_J dz^1 \wedge \cdots \wedge dz^{n-1} \wedge \partial r^m(z) \wedge d\overline{z}^J$ is an $(n, n - 1 - q)$ -form of M^m on D^m_ρ . Obviously, ψ^m converges to ψ in \mathcal{C}^1 -norm uniformly on \mathbb{R}^{2n-1} . Now one can see that $\int_{D_{\rho_1}} P'_{0, M^m_\rho} \varphi \wedge \overline{\partial}_{M^m} \psi^m$ converges to $\int_{D_{\rho_1}} P'_{0, M_\rho} \varphi \wedge \overline{\partial}_M \psi$. A similar argument shows that $Q'_{0, M^m_\rho} \overline{\partial}_{M^m} \varphi$ are uniformly bounded and converge to $Q'_{0, M_\rho} \overline{\partial}_M \varphi$ pointwise on D_{ρ_1} . Thus $\int_{D_{\rho_1}} (Q'_{0, M^m_\rho} \overline{\partial}_{M^m} \varphi) \wedge \psi^m$ converges to $\int_{D_{\rho_1}} (Q'_{0, M_\rho} \overline{\partial}_M \varphi) \wedge \psi$. By the homotopy formula (B.20),

$$\begin{aligned} \int_{D_{\rho_1}} \varphi \wedge \psi^m &= (-1)^q \int_{D_{\rho_1}} (P'_{0, M^m_\rho} \varphi) \wedge \overline{\partial}_{M^m} \psi^m \\ &\quad + \int_{D_{\rho_1}} (Q'_{0, M^m_\rho} \overline{\partial}_{M^m} \varphi) \wedge \psi^m. \end{aligned} \tag{B.21}$$

Taking limits, we see that as currents, $\varphi = \overline{\partial}_M P'_{0, M_\rho} \varphi + Q'_{0, M_\rho} \overline{\partial}_M \varphi$ holds on D_{ρ_1} . Since φ has compact support in $D_{\rho_1} \subset D_\rho$, we replace the domain of integration M_ρ by M_{ρ_0} and add boundary integrals. We get $\varphi = \overline{\partial}_M P'_{M_{\rho_0}} \varphi + Q'_{M_{\rho_0}} \overline{\partial}_M \varphi$ on D_{ρ_1} as currents, and hence on D_{ρ_0} whenever φ has compact support in D_{ρ_0} .

Return to the general case. Let M^m be as before. Take any ρ_1, ρ_2, ρ such that $0 < \rho_1 < \rho_2 < \rho < \rho_0$. Take a \mathcal{C}^∞ function χ which is 1 on D_{ρ_2} and has compact support in D_{ρ_0} . Let $\varphi_0 = \chi\varphi$ and $\varphi_1 = (1 - \chi)\varphi$. We have proved that $\varphi_0 = \overline{\partial}_M P'_{M_{\rho_0}} \varphi_0 + Q'_{M_{\rho_0}} \overline{\partial}_M \varphi_0$ as currents on D_{ρ_0} . Let ψ, ψ^m be as before, which are supported in D_{ρ_1} . By an analogy of (B.21), for $m > m_*$ we have

$$\int_{D_{\rho_1}} \varphi_1 \wedge \psi^m = (-1)^q \int_{D_{\rho_1}} (P'_{M^m_\rho} \varphi_1) \wedge \overline{\partial}_{M^m} \psi^m + \int_{D_{\rho_1}} (Q'_{M^m_\rho} \overline{\partial}_{M^m} \varphi_1) \wedge \psi^m.$$

Since $\text{supp } \psi^m \subset D_{\rho_1}$ and $\text{supp } \varphi_1 \subset \overline{D}_{\rho_0} \setminus D_{\rho_2}$, the dominated convergence theorem implies the convergence of $\int_{D_{\rho_1}} (P'_{0, M^m_\rho} \varphi_1) \wedge \overline{\partial}_{M^m} \psi^m$ to $\int_{D_{\rho_1}} (P'_{0, M_\rho} \varphi_1) \wedge \overline{\partial}_M \psi$. The integrands for the boundary integrals $(P'_{1, M^m_\rho} \varphi_1)(x)$ are of class \mathcal{C}^1 in a neighborhood of ∂D_ρ , when $x \in D_{\rho_2}$. Note that the map d_ρ sending $x \in \partial D_\rho$ to $r_\rho(x)x \in \partial D^m_\rho$ with $r_\rho > 0$ converges to the identity map in the \mathcal{C}^2 norm as $m \rightarrow \infty$. Thus we can verify that $P'_{1, M^m_\rho} \varphi_1$ converges uniformly to $P'_{1, M_\rho} \varphi_1$ on D_{ρ_1} . Combining with $\text{supp } \psi^m \subset D_{\rho_1}$, we obtain the convergence of $\int_{D_{\rho_1}} (P'_{1, M^m_\rho} \varphi_1) \wedge \overline{\partial}_{M^m} \psi^m$ to $\int_{D_{\rho_1}} (P'_{1, M_\rho} \varphi_1) \wedge \overline{\partial}_M \psi$. One can verify that $\int_{D_{\rho_1}} (Q'_{1, M^m_\rho} \overline{\partial}_{M^m} \varphi_1) \wedge \psi^m$ converges to $\int_{D_{\rho_1}} (Q'_{1, M_\rho} \overline{\partial}_M \varphi_1) \wedge \psi$. Hence $\varphi_1 = \overline{\partial}_M P_{M_\rho} \varphi_1 + Q_{M_\rho} \overline{\partial}_M \varphi_1$ holds

as currents on D_{ρ_1} . The definition of φ_1 is independent of $\rho > \rho_2$. Letting $\rho \rightarrow \rho_0$, we get $\varphi_1 = \bar{\partial}_M P_{M_{\rho_0}} \varphi_1 + Q_{M_{\rho_0}} \bar{\partial}_M \varphi_1$ on D_{ρ_1} . Add to φ_0 . We get $\varphi = \bar{\partial}_M P_{M_{\rho_0}} \varphi + Q_{M_{\rho_0}} \bar{\partial}_M \varphi$ as currents on D_{ρ_1} , and hence on D_{ρ_0} .

Case b) $r \in \mathcal{C}^2(D)$ and $\varphi, \bar{\partial}_M \varphi \in \mathcal{C}^0(\bar{M}_{\rho_0})$. We verify the homotopy formula by the Friedrichs approximation theorem, for which we need the commutator of a smoothing operator S_t and $\bar{\partial}_M$, applied to tangential $(0, q)$ -forms.

Recall that on M

$$\theta(\xi) = -2i\partial r(\zeta) = a d\xi^n \pmod{(d\zeta^\alpha, d\bar{\zeta}^\beta)}, \quad a = 1 + \hat{r}_{\xi^n}^2. \tag{B.22}$$

Let $\varphi = \sum_{|I|=q} \varphi_{\bar{I}}(x) d\bar{z}^{\bar{I}}$ be a continuous tangential $(0, q)$ form on M . Let χ be a smooth function of compact support in \mathbb{R}^{2n-1} such that $\int \chi dV = 1$. Let $\chi_t(x) = t^{1-2n} \chi(t^{-1}x)$ and define $S_t \varphi = \sum_{|I|=q} \varphi_{\bar{I}} * \chi_t d\bar{z}^{\bar{I}}$. Recall that $\bar{X}_\alpha = \partial_{\bar{z}^\alpha} + b_\alpha \partial_{x^n}$ with $b_\alpha = -r_{\bar{z}^\alpha} / (2r_{z^n})$.

Lemma B.5. *Let $0 \leq q < n - 1$. Let $M \subset \mathbb{C}^n$ be a graph of class \mathcal{C}^2 over $D \subset \mathbb{C}^{n-1} \times \mathbb{R}$. Assume that φ is a continuous tangential $(0, q)$ -form on M_ρ such that $\bar{\partial}_M \varphi$ is continuous on M_ρ . Let $0 < \rho' < \rho$. Then for t sufficiently small and on $M_{\rho'}$,*

$$\begin{aligned} [\bar{\partial}_M, S_t] \varphi(x) = & \sum_{|I|=q, 1 \leq \alpha < n} \int \left\{ \varphi_{\bar{I}}(x - t\xi) t^{-1} (b_{\bar{\alpha}}(x) - b_{\bar{\alpha}}(x - t\xi)) (\partial_{\xi^n} \chi)(\xi) \right. \\ & \left. - (\varphi_{\bar{I}} a^{-1} \bar{X}_\alpha a)(x - t\xi) \chi(\xi) \right\} dV(\xi) \wedge d\bar{z}^{\bar{\alpha}} \wedge d\bar{z}^{\bar{I}}. \end{aligned}$$

Proof. We follow a computation in [4]; see also [10, page 121]. Let $\varphi = \sum_{|I|=q} \varphi_{\bar{I}} d\bar{z}^{\bar{I}}$.

By the assumption, we have $\bar{\partial}_M \varphi = \sum_{|J|=q+1} \psi_{\bar{J}} d\bar{z}^{\bar{J}}$ with $\psi_{\bar{J}} \in \mathcal{C}^0$. Now

$$\begin{aligned} \bar{\partial}_M S_t \varphi(x) &= \sum_{|I|=q} \sum_{1 \leq \alpha < n} \int \varphi_{\bar{I}}(\xi) \bar{X}_\alpha^{(x)} \chi_t(x - \xi) dV(\xi) \wedge d\bar{z}^{\bar{\alpha}} \wedge d\bar{z}^{\bar{I}}, \\ S_t \bar{\partial}_M \varphi(x) &= \sum_{|J|=q+1} \int \psi_{\bar{J}}(\xi) \chi_t(x - \xi) dV(\xi) \wedge d\bar{z}^{\bar{J}}. \end{aligned}$$

Set $v = d\zeta^1 \wedge \dots \wedge d\zeta^{n-1} \wedge d\xi^n$. We may assume that $dV = d\bar{\zeta}^1 \wedge \dots \wedge d\bar{\zeta}^{n-1} \wedge v$. For each $J = (j_1, \dots, j_{q+1})$, there exist increasing indices $J^* = (j_1^*, \dots, j_{n-2-q}^*)$ and $\epsilon^J = \pm 1$ such that $d\bar{\zeta}^1 \wedge \dots \wedge d\bar{\zeta}^{n-1} = \epsilon^J d\bar{\zeta}^{\bar{J}^*} \wedge d\bar{\zeta}^{\bar{J}}$. Assume that $\psi_{\bar{J}}$ are

skew-symmetric. Thus

$$\begin{aligned} S_t \bar{\partial}_M \varphi(x) &= \sum_{|J|=q+1} \epsilon^J \int \psi_{\bar{J}}(\xi) \chi_t(x - \xi) d\bar{\zeta}'^J \wedge d\bar{\zeta}'^{J*} \wedge \nu \wedge d\bar{z}'^J \\ &= \sum_{|J|=q+1} \frac{\epsilon^J}{(q+1)!} \int \bar{\partial}_M^{(\xi)} \varphi \wedge \chi_t(x - \xi) d\bar{\zeta}'^{J*} \wedge \nu \wedge d\bar{z}'^J. \end{aligned}$$

By Stokes' formula, (3.1) and (B.22), $S_t \bar{\partial}_M \varphi(x)$ equals

$$\begin{aligned} &\sum_{|J|=q+1} \frac{(-1)^{q+1} \epsilon^J}{(q+1)!} \int \varphi(\xi) \wedge a(\xi) \bar{\partial}_M^{(\xi)} (a^{-1}(\xi) \chi_t(x - \xi)) \wedge d\bar{\zeta}'^{J*} \wedge \nu \wedge d\bar{z}'^J \\ &= \sum_{|I|=q} \sum_{1 \leq \alpha < n} \sum_{|J|=q+1} \frac{-\epsilon^J}{(q+1)!} \int \varphi_{\bar{I}}(\xi) a(\xi) \bar{X}_\alpha^{(\xi)} \frac{\chi_t(x - \xi)}{a(\xi)} d\bar{\zeta}'^\alpha \wedge d\bar{\zeta}'^I \wedge d\bar{\zeta}'^{J*} \wedge \nu \wedge d\bar{z}'^J \\ &= - \sum_{|I|=q} \sum_{1 \leq \alpha < n} \int \varphi_{\bar{I}}(\xi) a(\xi) \bar{X}_\alpha^{(\xi)} (a^{-1}(\xi) \chi_t(x - \xi)) dV(\xi) \wedge d\bar{z}'^\alpha \wedge d\bar{z}'^I. \end{aligned}$$

Here the last identity is seen as follows. Set $d\bar{z}'^\alpha \wedge d\bar{z}'^I = \epsilon_{\alpha I}^J d\bar{z}'^J$, if the identity can hold; otherwise, set $\epsilon_{\alpha I}^J = 0$. In the above summation $\sum_{|I|=q} \sum_{|J|=q+1}$, we may restrict to terms with $\epsilon_{\alpha I}^J \neq 0$. For $\epsilon_{\alpha I}^J \neq 0$, we have

$$\begin{aligned} \epsilon^J d\bar{\zeta}'^\alpha \wedge d\bar{\zeta}'^I \wedge d\bar{\zeta}'^{J*} \wedge d\bar{z}'^J &= \epsilon^J \epsilon_{\alpha I}^J d\bar{\zeta}'^J \wedge d\bar{\zeta}'^{J*} \wedge d\bar{z}'^J \\ &= \epsilon^J d\bar{\zeta}'^J \wedge d\bar{\zeta}'^{J*} \wedge d\bar{z}'^\alpha \wedge d\bar{z}'^I. \end{aligned}$$

Note that the last term is independent of J . We have $(q+1)!$ such terms for a fixed (αI) . Set $E_\alpha(\xi, x) = a(\xi) \bar{X}_\alpha^{(\xi)} (a^{-1}(\xi) \chi_t(x - \xi)) + \bar{X}_\alpha^{(x)} (\chi_t(x - \xi))$. We get

$$[\bar{\partial}_M, S_t] \varphi(x) = \sum_{|I|=q} \sum_{1 \leq \alpha < n} \int \varphi_{\bar{I}}(\xi) E_\alpha(\xi, x) dV(\xi) \wedge d\bar{z}'^\alpha \wedge d\bar{z}'^I,$$

which can be put into the form in the lemma. \square

We now derive the homotopy formula for the case b) via smoothing. We have

$$\bar{\partial}_M S_t \varphi - \bar{\partial}_M \varphi = S_t \bar{\partial}_M \varphi - \bar{\partial}_M \varphi + [\bar{\partial}_M, S_t](\varphi - S_t' \varphi) + [\bar{\partial}_M, S_t] S_t' \varphi.$$

Fix $\rho < \rho_1 < \rho_0$ and $\epsilon > 0$. Fix $t' > 0$ sufficiently small such that $|S_t' \varphi - \varphi| < \epsilon$ on M_{ρ_1} . By Lemma B.5, for all small t

$$\sup_{M_\rho} |[\bar{\partial}_M, S_t](\varphi - S_t' \varphi)| \leq C \sup_{M_{\rho_1}} |\varphi - S_t' \varphi| \leq C\epsilon. \quad (\text{B.23})$$

Let $S_t' \varphi = \sum_{|I|=q} \psi_{\bar{T}} d\bar{z}'^I$. Then $[\bar{\partial}_M, S_t] = \sum [b_{\bar{\alpha}} \partial_{x^n}, S_t]$ and

$$([\bar{\partial}_M, S_t] \psi)(x) = \sum_{1 \leq \alpha < n} \sum_{|I|=q} \int \chi_t(y) (b_{\bar{\alpha}}(x) - b_{\bar{\alpha}}(x-y)) \partial_{x^n} \psi_{\bar{T}}(x-y) dV(y) \wedge d\bar{z}'^I.$$

This shows that $\lim_{t \rightarrow 0} [\bar{\partial}_M, S_t] S_t' \varphi = 0$ on M_ρ . We also have $\lim_{t \rightarrow 0} S_t \bar{\partial}_M \varphi = \bar{\partial}_M \varphi$ on M_ρ . Therefore, (B.23) implies that $\lim_{t \rightarrow 0} \bar{\partial}_M S_t \varphi = \bar{\partial}_M \varphi$ on M_ρ . Now by case a), $\varphi_t = \bar{\partial}_M P'_{M_\rho} \varphi_t + Q'_{M_\rho} \bar{\partial}_M \varphi_t$ holds on M_ρ as currents. Letting $t \rightarrow 0$ and then $\rho \rightarrow \rho_0$, we obtain $\varphi = \bar{\partial}_M P'_{M_0} \varphi + Q'_{M_0} \bar{\partial}_M \varphi$ on M_{ρ_0} in the sense of currents.

References

- [1] J. BRUNA and J. M. BURGÚÉS, *Holomorphic approximation and estimates for the $\bar{\partial}$ -equation on strictly pseudoconvex nonsmooth domains*, Duke Math. J. **55** (1987), 539–596.
- [2] S.-C. Chen and M.-C. Shaw, “Partial Differential Equations in Several Complex Variables”, AMS/IP Studies in Advanced Mathematics, Vol. 19, Amer. Math. Soc., Providence, RI, International Press, Boston, MA, 2001.
- [3] G. B. FOLLAND and E. M. STEIN, *Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group*, Comm. Pure Appl. Math. **27** (1974), 429–522.
- [4] K. O. FRIEDRICH, *The identity of weak and strong extensions of differential operators*, Trans. Amer. Math. Soc. **55** (1944), 132–151.
- [5] X. GONG and S. M. WEBSTER, *Regularity for the CR vector bundle problem I*, Pure and Appl. Math. Quarterly **6** (2010), 983–998.
- [6] X. GONG and S. M. WEBSTER, *Regularity in the local CR embedding problem*, J. Geom. Anal. DOI: 10.1007/s12220 - 010 - 9192 - 6.
- [7] G. M. HENKIN, *The Lewy equation and analysis on pseudoconvex domains*, Russian Math. Surveys **32** (1977), 59–130.
- [8] G. M. HENKIN and A. V. ROMANOV, *Exact Hölder estimates of the solutions of the $\bar{\partial}$ -equation*, Izv. Akad. Nauk SSSR Ser. Mat. **35** (1971), 1171–1183.
- [9] L. HÖRMANDER, *The boundary problems of physical geodesy*, Arch. Rational Mech. Anal. **62** (1976), 1–52.
- [10] L. HÖRMANDER, “An Introduction to Complex Analysis in Several Variables”, North Holland, Amsterdam, 3rd edition, 1990.
- [11] N. KERZMAN, *Hölder and L^p estimates for solutions of $\bar{\partial}u = f$ in strongly pseudoconvex domains*, Comm. Pure Appl. Math. **24** (1971), 301–379.
- [12] I. LIEB and R. M. RANGE, *Lösungsoperatoren für den Cauchy-Riemann-Komplex mit C^k -Abschätzungen*, Math. Ann. **253** (1980), 145–164.
- [13] L. MA and J. MICHEL, *Local regularity for the tangential Cauchy-Riemann complex*, J. Reine Angew. Math. **442** (1993), 63–90.
- [14] L. MA and J. MICHEL, *Regularity of local embeddings of strictly pseudoconvex CR structures*, J. Reine Angew. Math. **447** (1994), 147–164.
- [15] J. MICHEL and L. MA, *On the regularity of CR structures for almost CR vector bundles*, Math. Z. **218** (1995), 135–142.
- [16] A. NAGEL and J.-P. ROSAY, *Nonexistence of homotopy formula for $(0, 1)$ forms on hypersurfaces in C^3* , Duke Math. J. **58** (1989), 823–827.
- [17] C. ROMERO, “Potential Theory for the Kohn Laplacian on the Heisenberg Group”, Thesis, University of Minnesota, 1991.

- [18] M.-C. SHAW, *L^p estimates for local solutions of $\bar{\partial}_b$ on strongly pseudo-convex CR manifolds*, Math. Ann. **288** (1990), 35–62.
- [19] Y. T. SIU, *The $\bar{\partial}$ problem with uniform bounds on derivatives*, Math. Ann. **207** (1974), 163–176.
- [20] S. M. WEBSTER, *A new proof of the Newlander-Nirenberg theorem*, Math. Z. **201** (1989), 303–316.
- [21] S. M. WEBSTER, *On the local solution of the tangential Cauchy-Riemann equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **6** (1989), 167–182.
- [22] S. M. WEBSTER, *On the proof of Kuranishi's embedding theorem*, Ann. Inst. H. Poincaré Anal. Non Linéaire **6** (1989), 183–207.
- [23] S. M. WEBSTER, *The integrability problem for CR vector bundles*, In: “Several Complex Variables and Complex Geometry”, Part 3 (Santa Cruz, CA, 1989), Proc. Sympos. Pure Math., 52, Part 3, Amer. Math. Soc., Providence, RI, 1991, 355–368.

Department of Mathematics
University of Wisconsin
Madison, WI 53706, USA
gong@math.wisc.edu

Department of Mathematics
University of Chicago
Chicago, IL 60637, USA
webster@math.uchicago.edu