



ANNALES

DE

L'INSTITUT FOURIER

Hirokazu NASU

Obstructions to deforming curves on a 3-fold, II: Deformations of degenerate curves on a del Pezzo 3-fold

Tome 60, n° 4 (2010), p. 1289-1316.

http://aif.cedram.org/item?id=AIF_2010__60_4_1289_0

© Association des Annales de l'institut Fourier, 2010, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

OBSTRUCTIONS TO DEFORMING CURVES ON A 3-FOLD, II: DEFORMATIONS OF DEGENERATE CURVES ON A DEL PEZZO 3-FOLD

by Hirokazu NASU (*)

ABSTRACT. — We study the Hilbert scheme $\text{Hilb}^{sc}V$ of smooth connected curves on a smooth del Pezzo 3-fold V . We prove that any degenerate curve C , i.e. any curve C contained in a smooth hyperplane section S of V , does not deform to a non-degenerate curve if the following two conditions are satisfied: (i) $\chi(V, \mathcal{I}_C(S)) \geq 1$ and (ii) for every line ℓ on S such that $\ell \cap C = \emptyset$, the normal bundle $N_{\ell/V}$ is trivial (i.e. $N_{\ell/V} \simeq \mathcal{O}_{\mathbb{P}^1}^{\oplus 2}$). As a consequence, we prove an analogue (for $\text{Hilb}^{sc}V$) of a conjecture of J. O. Kleppe, which is concerned with non-reduced components of the Hilbert scheme $\text{Hilb}^{sc}\mathbb{P}^3$ of curves in the projective 3-space \mathbb{P}^3 .

RÉSUMÉ. — Nous étudions le schéma de Hilbert $\text{Hilb}^{sc}V$ des courbes lisses connexes sur une variété de del Pezzo lisse V de dimension 3. Nous montrons qu'aucune courbe C dégénérée, c'est-à-dire, aucune courbe C contenue dans une section hyperplane S de V , se déforme en une courbe non-dégénérée, si les deux conditions suivantes sont satisfaites : (i) $\chi(V, \mathcal{I}_C(S)) \geq 1$ et (ii) pour chaque droite ℓ sur S telle que $\ell \cap C = \emptyset$, le fibré normal $N_{\ell/V}$ de ℓ dans V est trivial. Par conséquent, nous prouvons un analogue (pour $\text{Hilb}^{sc}V$) d'une conjecture de J. O. Kleppe, qui concerne les composantes non-réduites du schéma de Hilbert $\text{Hilb}^{sc}\mathbb{P}^3$ des courbes dans l'espace projectif \mathbb{P}^3 de dimension 3.

1. Introduction

This paper is a sequel to a joint work [13] with Shigeru Mukai. In [13] the embedded deformations of smooth curves C on a smooth projective 3-fold V have been studied under the presence of a smooth surface S such that $C \subset S \subset V$, especially when V is a uniruled 3-fold. In this paper, the same subject is studied in detail especially when V is a del Pezzo 3-fold.

Keywords: Hilbert scheme, infinitesimal deformation, del Pezzo variety.

Math. classification: 14C05, 14H10, 14D15.

(*) Supported in part by the 21-st century COE program “Formation of an International Center of Excellence in the Frontiers of Mathematics and Fostering of Researchers in Future Generations”.

It is known that even if the deformations of C in S and the deformations of S in V behave well, those of C in V behave badly in general. For example, even if $\text{Hilb } V$ and $\text{Hilb } S$ are nonsingular of expected dimension $\chi(N_{S/V})$ and $\chi(N_{C/S})$ at $[S]$ and $[C]$ respectively, $\text{Hilb } V$ can be generically non-reduced along some component passing through $[C]$ (cf. Mumford's example in [14]). Such non-reduced components of the Hilbert scheme $\text{Hilb}^{sc} V$ of smooth connected curves on V have been constructed for many uniruled 3-folds V in [13]. The non-reducedness is originated from the non-surjectivity of the restriction map

$$(1.1) \quad H^0(S, N_{S/V}) \xrightarrow{|_C} H^0(C, N_{S/V}|_C).$$

We say that C is *stably degenerate* if every small global deformation of C in V is contained in a deformation S' of S in V (cf. Definition 4.1). If (1.1) is surjective, then C is stably degenerate (cf. Proposition 4.3). However if it is not surjective, then there exists a first order deformation \tilde{C} of C in V which is not contained in any first order deformation \tilde{S} of S . In this paper, we consider the following problem raised by Mukai:

PROBLEM 1.1. — *Suppose that (1.1) is not surjective and $\chi(V, \mathcal{I}_C(S)) > 0$. Then (1) Is C stably degenerate? (2) Is $\text{Hilb}^{sc} V$ singular at $[C]$?*

Here \mathcal{I}_C denotes the ideal sheaf of C in V and $\mathcal{I}_C(S) := \mathcal{I}_C \otimes \mathcal{O}_V(S)$. J. O. Kleppe [8] and Ph. Ellia [2] considered Problem 1.1 for the case where V is the projective 3-space \mathbb{P}^3 , S is a smooth cubic surface in \mathbb{P}^3 and C is a smooth connected curve of degree d lying on S . Kleppe gave a conjecture (cf. Conjectures 5.1), which can be reformulated as follows:

CONJECTURE 1.2. — *Let $C \subset S \subset \mathbb{P}^3$ be as above and assume that $\chi(\mathbb{P}^3, \mathcal{I}_C(3)) \geq 1$. Then:*

- (1) *If C is linearly normal, then every small global deformation C' of C in \mathbb{P}^3 is contained in a cubic surface $S' \subset \mathbb{P}^3$, i.e., C is stably degenerate, and*
- (2) *Suppose that C is general and $d > 9$. Then $\text{Hilb}^{sc} \mathbb{P}^3$ is nonsingular at $[C]$ if and only if $H^1(\mathbb{P}^3, \mathcal{I}_C(3)) = 0$.*

As a testing ground of his conjecture, we consider Problem 1.1 for the case where V is a smooth del Pezzo 3-fold (cf. §2.2), S is a smooth member of the class $|H|$ of the polarization H of V , i.e., a smooth del Pezzo surface in V , and C is a smooth connected curve on S . The following theorem is an analogue of Kleppe's conjecture.

THEOREM 1.3. — *Let $C \subset S \subset V$ be as above and assume that $\chi(V, \mathcal{I}_C(S)) \geq 1$. If every line ℓ on S such that $C \cap \ell = \emptyset$ is a good line on V (i.e., the normal bundle $N_{\ell/V}$ of ℓ in V is trivial), then:*

- (1) C is stably degenerate, and
- (2) $\text{Hilb}^{sc} V$ is nonsingular at $[C]$ if and only if $H^1(V, \mathcal{I}_C(S)) = 0$.

If $\chi(V, \mathcal{I}_C(S)) < 1$, then it follows from a dimension count that C is not stably degenerate (Proposition 4.7). If some ℓ is a *bad line* on V (i.e., $N_{\ell/V} \not\cong \mathcal{O}_{\mathbb{P}^1}^{\oplus 2}$) then C is not necessarily stably degenerate (Proposition 5.4). As a corollary to Theorem 1.3, we give a sufficient condition for a maximal family W of degenerate curves on V to become an irreducible component of the Hilbert scheme $\text{Hilb}^{sc} V$ and determine whether $\text{Hilb}^{sc} V$ is generically non-reduced along W or not (Theorem 4.14).

One of the main tools used in this paper is the infinitesimal analysis of the Hilbert scheme developed in [13]. As is well known, every infinitesimal deformation \tilde{C} of C in V of the first order (i.e., over $\text{Spec } k[t]/(t^2)$) determines a global section $\alpha \in H^0(N_{C/V})$ and a cohomology class $\text{ob}(\alpha) \in H^1(N_{C/V})$ (called the *obstruction*) such that \tilde{C} lifts to a deformation over $\text{Spec } k[t]/(t^3)$ if and only if $\text{ob}(\alpha) = 0$ (cf. §2.3). Let $\pi_{C/S} : N_{C/V} \rightarrow N_{S/V}|_C$ be the natural projection. In [13] Mukai and Nasu studied the exterior component of α and $\text{ob}(\alpha)$, i.e., the images of α and $\text{ob}(\alpha)$ by the induced maps $H^i(\pi_{C/S}) : H^i(N_{C/V}) \rightarrow H^i(N_{S/V}|_C)$ ($i = 0, 1$), respectively. They proved that if there exists a curve E on S such that $(E^2)_S < 0$ (e.g. $(-1)\text{-}\mathbb{P}^1$ on S) and the exterior component of α lifts to a global section $v \in H^0(N_{S/V}(E)) \setminus H^0(N_{S/V})$, then the exterior component of $\text{ob}(\alpha)$ is nonzero provided that certain additional conditions on E , C and v hold (see [13, Theorem 2.2]). Such a rational section v of $N_{S/V}$ admitting a pole along E is called an *infinitesimal deformation with a pole*. In §3 we see that an infinitesimal deformation with a pole along E induces an obstructed infinitesimal deformation of the open surface $S^\circ := S \setminus E$ in the open 3-fold $V^\circ := V \setminus E$ (Theorem 3.1). By using this fact, we prove Theorem 1.3 in §4. In §5 we give some examples of generically non-reduced components of the Hilbert scheme of curves on a del Pezzo 3-fold as an application.

Acknowledgements. I should like to express my sincere gratitude to Professor Shigeru Mukai. He showed me the example of non-reduced components of the Hilbert scheme of canonical curves in §5.2 as a simplification of Mumford’s example of a non-reduced component of $\text{Hilb}^{sc} \mathbb{P}^3$. This motivated me to research the topic of this paper. Throughout this research, he made many suggestions which are useful for obtaining and improving the

proofs. According to his suggestion, I studied the deformation theory of an open surface in an open 3-fold and organized §3. I am grateful to Professor Jan Oddvar Kleppe for giving me useful comments on Hilbert-flag schemes and for finding a gap in the proof of Lemma 4.8 in a earlier version. I should like to thank the referee, who showed me a straight proof of Proposition 4.3 and led me to a simplification of §4.1 and §4.2. According to his/her recommendation, I give the classes of non-reduced components of the Hilbert scheme explicitly in Proposition 5.5.

Notation and Conventions. We work over an algebraically closed field k of characteristic 0. Let V be a scheme over k and let X be a closed subscheme of V . Then \mathcal{I}_X denotes the ideal sheaf of X in V and $N_{X/V}$ denotes the normal sheaf $(\mathcal{I}_X/\mathcal{I}_X^2)^\vee$ of X in V . For a sheaf \mathcal{F} on V , we denote the restriction map $H^i(V, \mathcal{F}) \rightarrow H^i(X, \mathcal{F}|_X)$ by $|_X$. We denote the Euler-Poincaré characteristic of \mathcal{F} by $\chi(V, \mathcal{F})$ or $\chi(\mathcal{F})$. $\text{Hilb}^{sc} V$ denotes the open subscheme of the Hilbert scheme $\text{Hilb} V$ whose point corresponds to a smooth connected curve on V .

2. Preliminaries

The results in this section will be used in § 4. Proposition 2.4 and Lemma 2.5 are important to our proof of Proposition 4.9 and 4.10, respectively.

2.1. Del Pezzo surfaces

A *del Pezzo surface* is a smooth surface S whose anti-canonical divisor $-K_S$ is ample. Every del Pezzo surface is isomorphic to \mathbb{P}^2 blown up at fewer than 9 points or $\mathbb{P}^1 \times \mathbb{P}^1$. We denote the blow-up of \mathbb{P}^2 at $(9-n)$ -points by S_n . A curve $\ell \simeq \mathbb{P}^1$ on S_n is called a *line*⁽¹⁾ if $\ell \cdot (-K_S) = 1$. Every (-1) - \mathbb{P}^1 on S_n is a line and every line on S_n is a (-1) - \mathbb{P}^1 . A curve q on S_n is called a *conic* if $q \cdot (-K_S) = 2$ and $q^2 = 0$.

LEMMA 2.1. — *Let D be a divisor on a del Pezzo surface S . If D is nef and $\chi(S, -D) \geq 0$, then $H^1(S, -D) = 0$.*

⁽¹⁾ There exists no line on \mathbb{P}^2 and on $\mathbb{P}^1 \times \mathbb{P}^1$.

Proof. — If $D^2 > 0$ then the assertion follows the Kawamata-Viehweg vanishing. Since D is a nef divisor on a del Pezzo surface, we have $D^2 \geq 0$. Now we assume that $D^2 = 0$. If $S = S_n$, then D is linearly equivalent to a multiple mq ($m \geq 0$) of a conic q on S . By the Riemann-Roch theorem, we have

$$\begin{aligned} \chi(S, -D) &= \frac{1}{2}(-mq) \cdot (-mq - K_S) + \chi(\mathcal{O}_S) \\ &= -m + 1. \end{aligned}$$

Thus we have $m = 0$ or 1 by assumption. This implies that $H^1(-mq) = 0$. If $S = \mathbb{P}^1 \times \mathbb{P}^1$, then D is of bidegree $(m, 0)$ or $(0, m)$ with $m \geq 0$. Again by the Riemann-Roch theorem, we have $\chi(-D) = -m + 1 \geq 0$. Thus $H^1(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(-D)) = 0$. □

LEMMA 2.2. — *Let D be an effective divisor on a del Pezzo surface S . Then the lines ℓ such that $D \cdot \ell < 0$ are mutually disjoint. The fixed part⁽²⁾ $\text{Bs } |D|$ of the linear system $|D|$ on S is equal to*

$$- \sum_{D \cdot \ell < 0} (D \cdot \ell)\ell.$$

Proof. — We prove the two assertions at the same time. It is clear that any line ℓ satisfying $D \cdot \ell < 0$ is contained in $\text{Bs } |D|$. On the other hand, except for lines on S every irreducible curve C on S can move on S by the linearly equivalence since $\chi(C) \geq 2$ and $H^2(C) = 0$. Hence $|D|$ is decomposed into the sum

$$|D| = |D'| + \sum_{i=1}^k m_i \ell_i,$$

of a linear system $|D'|$ on S such that $\text{Bs } |D'| = \emptyset$ and some lines ℓ_i on S with coefficients $m_i \in \mathbb{Z}_{>0}$ ($1 \leq i \leq k$). If $\ell_i \cap \ell_j \neq \emptyset$ for some $i \neq j$, then $\ell_i + \ell_j$ is a (reducible) conic on S and can move on S by $\chi(\ell_i + \ell_j) = 2$. Thus ℓ_i 's are mutually disjoint. Now we prove that $D \cdot \ell_i < 0$ for any i . Since $m_i = (D' - D) \cdot \ell_i > 0$, it suffices to show that $D' \cdot \ell_i = 0$. Since D' is nef, we have $(D')^2 \geq 0$. Since $-K_S$ is ample, so is $D' - K_S$. Hence we have $H^1(D') = H^1((D' - K_S) + K_S) = 0$ by the Kodaira vanishing. If $D' \cdot \ell_i \geq 1$, then it follows from the exact sequence

$$0 \longrightarrow \mathcal{O}_S(D') \longrightarrow \mathcal{O}_S(D' + \ell_i) \longrightarrow \mathcal{O}_S(D' + \ell_i)|_{\ell_i} \longrightarrow 0$$

that $h^0(D' + \ell_i) > h^0(D')$. Thus we have $D' \cdot \ell_i = 0$. □

(2) the base locus of dimension one

LEMMA 2.3. — Let E be a disjoint union of m lines ($m \geq 0$) on a del Pezzo surface S and let $\varepsilon : S \rightarrow F$ be the blow-down of E from S . If a divisor D on F satisfies $h^0(F, D) \geq m$, then we have the following:

- (1) $h^0(S, \varepsilon^*D - E) = h^0(F, D) - m$, and
- (2) If $H^1(S, \varepsilon^*D) = 0$, then $H^1(S, \varepsilon^*D - E) = 0$.

Proof. — (1) Let ℓ_i ($1 \leq i \leq m$) be the disjoint lines on S and let $E := \sum_{i=1}^m \ell_i$. We put $D_j := \varepsilon^*D - \sum_{1 \leq i \leq j} \ell_i$. Since the image of ℓ_i on F is a point, we have $h^0(D_j) \geq h^0(D) - j$ for every $1 \leq j \leq m$. Moreover since $D_{j-1} \cdot \ell_j = 0$, Lemma 2.2 shows that ℓ_j is not contained in $\text{Bs } |D_{j-1}|$. Hence $\dim |D_j|$ decreases one by one as j increases. Therefore we have $h^0(\varepsilon^*D - E) = h^0(D_m) = h^0(D) - m$.

(2) An exact sequence $0 \rightarrow \mathcal{O}_S(\varepsilon^*D - E) \rightarrow \mathcal{O}_S(\varepsilon^*D) \rightarrow \mathcal{O}_E \rightarrow 0$ on S induces an exact sequence

$$H^0(S, \varepsilon^*D) \xrightarrow{\rho} H^0(E, \mathcal{O}_E) \longrightarrow H^1(S, \varepsilon^*D - E) \longrightarrow H^1(S, \varepsilon^*D)$$

of cohomology groups. Then ρ is surjective by (1) and $H^1(S, \varepsilon^*D) = 0$ by assumption. Hence we have $H^1(S, \varepsilon^*D - E) = 0$. □

Let C be a smooth connected curve on a del Pezzo surface S . We consider the restriction to C of the anti-canonical linear system $| -K_S |$ on S . The restriction map $H^0(-K_S) \rightarrow H^0(-K_S|_C)$ is not surjective in general. Let ℓ_i ($1 \leq i \leq m$) be the lines on S disjoint to C . Let us define an effective divisor E on S by the sum

$$E := \sum_{i=1}^m \ell_i$$

and we put $E := 0$ if there exists no such ℓ_i . If C is neither a line nor a conic, then the ℓ_i 's are mutually disjoint: indeed if $\ell_i \cap \ell_j \neq \emptyset$ for some $i \neq j$, then $q := \ell_i + \ell_j$ is a conic on S and hence C intersects q by $C \cdot q > 0$.

PROPOSITION 2.4. — Assume that C is irrational and $\chi(S, -K_S - C) \geq 0$. Then we have $H^1(S, -K_S + E - C) = 0$ and the restriction map

$$(2.1) \quad H^0(S, -K_S + E) \xrightarrow{|_C} H^0(C, -K_S|_C)$$

is surjective. If C is not elliptic either, then the map (2.1) is an isomorphism.

Proof. — It suffices to show that $H^1(-K_S + E - C) = 0$ by the exact sequence

$$(2.2) \quad 0 \longrightarrow \mathcal{O}_S(-K_S + E - C) \longrightarrow \mathcal{O}_S(-K_S + E) \longrightarrow \mathcal{O}_S(-K_S)|_C \longrightarrow 0.$$

Claim. Put $D_1 := C + K_S - E$. Then D_1 is nef.

Since S is regular (i.e., $H^1(K_S) = 0$), the restriction map $|_C : H^0(C + K_S) \rightarrow H^0(K_C)$ is surjective. Since $C \not\cong \mathbb{P}^1$, the linear system $|C + K_S|$ on S is non-empty. Let ℓ be a line on S . Since C is not a line, we have $C \cdot \ell \geq 0$ and hence $(C + K_S) \cdot \ell \geq -1$. By Lemma 2.2, ℓ is contained in $\text{Bs}|C + K_S|$ if and only if $C \cap \ell = \emptyset$. Thus we have $E = \text{Bs}|C + K_S|$ and $|D_1|$ does not have base components. In particular, D_1 is nef.

It follows from the exact sequence

$$(2.3) \quad 0 \rightarrow \mathcal{O}_S(-K_S - C) \rightarrow \mathcal{O}_S(-K_S + E - C) \rightarrow \underbrace{\mathcal{O}_S(-K_S + E)|_E}_{\simeq \mathcal{O}_E} \rightarrow 0$$

that $\chi(-D_1) = \chi(-K_S - C) + \chi(\mathcal{O}_E) \geq 0$. Hence we have $H^1(-D_1) = 0$ by Lemma 2.1.

Now we assume that C is not elliptic. Then $K_C \not\sim 0$ and hence $C + K_S \not\sim E$ by adjunction. Thus $D_1 \not\sim 0$ and $H^0(-D_1) = 0$. Therefore (2.1) is injective. □

LEMMA 2.5. — *If C is not rational nor elliptic and $\chi(S, -K_S - C) \geq 0$, then the map*

$$H^1(S, -K_S + 3E) \xrightarrow{|_C} H^1(C, -K_S|_C)$$

induced by (2.2) $\otimes \mathcal{O}_S(2E)$ is injective.

Proof. — It suffices to show that $H^1(-K_S + 3E - C) = 0$. Let $\varepsilon : S \rightarrow F$ be the blow-down of E from S . Then there exists a divisor D_2 on F such that $\varepsilon^*D_2 \sim C + 2K_S - 2E$. By the Serre duality, it suffices to show that $H^1(\varepsilon^*D_2 - E) = 0$.

Claim. $H^i(S, \varepsilon^*D_2) = 0$ for $i = 1, 2$.

By (2.3) $\otimes \mathcal{O}_S(E)$, there exists an exact sequence

$$H^1(S, -K_S + E - C) \rightarrow H^1(S, -K_S + 2E - C) \rightarrow H^1(E, (-K_S + 2E)|_E).$$

Since $H^1((-K_S + 2E)|_E) \simeq H^1(\mathcal{O}_E(E)) = 0$ and $H^1(-K_S + E - C) = 0$ by Proposition 2.4, we have $H^1(-K_S + 2E - C) = 0$. By the Serre duality, we have $H^1(\varepsilon^*D_2) = 0$. Similarly by the Serre duality, we have $H^2(\varepsilon^*D_2) \simeq H^0(K_S - \varepsilon^*D_2)^\vee$. Since C is not rational nor elliptic, we have $(K_S - \varepsilon^*D_2) \cdot C = (-K_S - C) \cdot C = -\text{deg } K_C < 0$. Hence we have $H^2(\varepsilon^*D_2) = 0$ because C is nef. Thus the claim has been proved.

By this claim, we have $h^0(F, D_2) = h^0(S, \varepsilon^*D_2) = \chi(S, \varepsilon^*D_2)$. Then an easy calculation shows that $\chi(\varepsilon^*D_2) = \chi(-K_S - C) + \chi(\mathcal{O}_E)$. Since $\chi(-K_S - C) \geq 0$, we have $h^0(F, D_2) = \chi(S, \varepsilon^*D_2) \geq m$, where m is the number of components of E . Since $H^1(\varepsilon^*D_2) = 0$, Lemma 2.3 (2) shows that $H^1(\varepsilon^*D_2 - E) = 0$. □

Let S be a smooth projective surface and let L be a line bundle on S .

LEMMA 2.6. — *Let E be a disjoint union of irreducible curves E_i ($i = 1, \dots, m$) on S such that $E_i^2 < 0$ and let $\iota : S^\circ := S \setminus E \hookrightarrow S$ be the open immersion. If $\deg(L|_{E_i}) \leq 0$ for every i , then the map*

$$H^1(S, L) \rightarrow H^1(S^\circ, L|_{S^\circ})$$

induced by the sheaf inclusion $L \hookrightarrow L \otimes \iota_ \mathcal{O}_{S^\circ}$ is injective.*

The proof is similar to that of [13, Lemma 2.5] and we omit it here. Lemma 2.6 allows us to identify $H^1(S, L(nE))$ ($n \geq 0$) with their images in $H^1(S^\circ, L|_{S^\circ})$. As a result, under the identification we obtain a natural filtration

$$H^1(S, L) \subset H^1(S, L(E)) \subset H^1(S, L(2E)) \subset \dots \subset H^1(S^\circ, L|_{S^\circ})$$

on $H^1(S^\circ, L|_{S^\circ})$.

2.2. Del Pezzo threefolds

A *del Pezzo threefold* is a pair (V, H) consisting of a (smooth) irreducible projective variety V of dimension 3 and an ample Cartier divisor H on V such that $-K_V = 2H$. Here H is called the *polarization* of V and sometimes omitted. The self-intersection number $n := H^3$ is called the *degree* of V . It is known that the linear system $|H|$ on V determines a double cover $\varphi_{|H|} : V \rightarrow \mathbb{P}^3$ if $n = 2$, and an embedding $\varphi_{|H|} : V \hookrightarrow \mathbb{P}^{n+1}$ if $n \geq 3$. If S is a smooth member of $|H|$, then the pair $(S, H|_S)$ is a del Pezzo surface of degree n . Every smooth del Pezzo 3-fold is one of V_n ($1 \leq n \leq 8$) or V'_6 in Table 2.1, in which $\mathbb{L}^{(i)}$ denotes a linear subspace of dimension i , and n and ρ respectively denote the degree and the Picard number of V_n (and of V'_6) (cf. [4],[5],[6]). It is known that a smooth 3-fold $V \subset \mathbb{P}^{n+1}$ ($n \geq 3$) is a del Pezzo 3-fold of degree n if a linear section $[V \subset \mathbb{P}^{n+1}] \cap H_1 \cap H_2$ with two general hyperplanes $H_1, H_2 \subset \mathbb{P}^{n+1}$ is an elliptic normal curve in \mathbb{P}^{n-1} .

We briefly review the basics of the Hilbert scheme of lines on a del Pezzo 3-fold. We refer to Iskovskih ([6],[7]) for the details. Let (V, H) be a smooth del Pezzo 3-fold of degree n . By a *line* on (V, H) , we mean a reduced irreducible curve ℓ on V such that $(\ell \cdot H)_V = 1$ and $\ell \simeq \mathbb{P}^1$. If $n \leq 7$ then V contains a line ℓ . Then there are only the following possibilities for the

Table 2.1. Del Pezzo 3-folds

del Pezzo 3-folds	n	ρ	
$V_1 = (6) \subset \mathbb{P}(3, 2, 1, 1, 1)$	1	1	a weighted hypersurface of degree 6
$V_2 = (4) \subset \mathbb{P}(2, 1, 1, 1, 1)$	2	1	a weighted hypersurface of degree 4 ^(a)
$V_3 = (3) \subset \mathbb{P}^4$	3	1	a cubic hypersurface
$V_4 = (2) \cap (2) \subset \mathbb{P}^5$	4	1	a complete intersection of two quadrics
$V_5 = [\text{Gr}(2, 5) \xrightarrow{\text{Plücker}} \mathbb{P}^9] \cap \mathbb{L}^{(6)}$	5	1	a linear section of Grassmannian
$V_6 = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{\text{Segre}} \mathbb{P}^7$	6	3	
$V'_6 = [\mathbb{P}^2 \times \mathbb{P}^2 \xrightarrow{\text{Segre}} \mathbb{P}^8] \cap \mathbb{L}^{(7)}$	6	2	
$V_7 = \text{Bl}_{\text{pt}} \mathbb{P}^3 \subset \mathbb{P}^8$	7	2	the blow-up of \mathbb{P}^3 at a point ^(b)
$V_8 = \mathbb{P}^3 \xrightarrow{\text{Veronese}} \mathbb{P}^9$	8	1	the Veronese image of \mathbb{P}^3

^(a) Another realization of V_2 is a double cover of \mathbb{P}^3 branched along a quartic surface.
^(b) V_7 is realized as the projection of $V_8 \subset \mathbb{P}^9$ from one of its points.

normal bundle $N_{\ell/V}$ of ℓ in V :

$$\begin{aligned}
 (0,0): \quad N_{\ell/V} &\simeq \mathcal{O}_{\mathbb{P}^1}^{\oplus 2} && \text{(i.e., trivial),} \\
 (1,-1): \quad N_{\ell/V} &\simeq \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(1), \\
 (2,-2): \quad N_{\ell/V} &\simeq \mathcal{O}_{\mathbb{P}^1}(-2) \oplus \mathcal{O}_{\mathbb{P}^1}(2) && \text{(only if } n = 1 \text{ or } 2), \\
 (3,-3): \quad N_{\ell/V} &\simeq \mathcal{O}_{\mathbb{P}^1}(-3) \oplus \mathcal{O}_{\mathbb{P}^1}(3) && \text{(only if } n = 1).
 \end{aligned}$$

In this paper, ℓ is called a *good line* if $N_{\ell/V}$ is trivial, and called a *bad line* otherwise. If $n \geq 3$, then every line on V is of type $(0, 0)$ or $(1, -1)$. The Hilbert scheme Γ of lines on V is called the *Fano surface* of V , and in fact every irreducible (non-embedded) component of Γ is of dimension two. Let $\Gamma_i \subset \Gamma$ be an irreducible component and let S_i be the universal family of lines on V over Γ_i . Then there exists a natural diagram

$$\begin{array}{ccc}
 S_i & \xrightarrow{p} & V \\
 \pi \downarrow & & \\
 \Gamma_i & &
 \end{array}$$

By [7, Chap. III, Proposition 1.3 (iv)], if $n \geq 3$ then we have either

- (a) p is surjective; in this case a general line in Γ_i is a good line; or
- (b) $p(S_i) \simeq \mathbb{P}^2$ is a plane on $V \subset \mathbb{P}^{n+1}$; in this case every line in Γ_i is a bad line.

We have either (a) or (b) also when $n \leq 2$. (See the proof⁽³⁾ in [7], which works for $n \leq 2$.) If $n \neq 7$ then every irreducible component of Γ is of type (a). If $n = 7$ then Γ consists of two irreducible components $\Gamma_i \simeq \mathbb{P}^2 (i = 0, 1)$, one of which is of type (a), while the other is of type (b). Consequently, we have

⁽³⁾ In the proof, the assumption that $\text{char } k = 0$ is used.

LEMMA 2.7 (Iskovskih). — *Every smooth del Pezzo 3-fold of degree $n \neq 8$ contains a good line.*

LEMMA 2.8. — *Let (V, H) be a smooth del Pezzo 3-fold of degree n and let S be a general member of $|H|$. If $n \neq 7$ then every line on S is good. If $n = 7$ then there exist three lines ℓ_0, ℓ_1, ℓ_2 on S forming the configuration in Figure 2.1. Then ℓ_0 is bad, while ℓ_1 and ℓ_2 are good.*




Figure 2.1. $(-1)\text{-}\mathbb{P}^1$'s on S_7

Proof. — There exists no line on V_8 . If $n \neq 7$, then the locus \mathfrak{B} of bad lines in the Fano surface Γ is of dimension one. Let p_i denote the projection of

$$\{(\ell, S) \mid \ell \subset S\} \subset \Gamma \times |H|$$

to the i -th factor. Since the fiber of p_1 is of dimension $n - 1$, $p_2(p_1^{-1}(\mathfrak{B}))$ is a proper closed subset of $|H| \simeq \mathbb{P}^{n+1}$. Hence every line on a general member S of $|H|$ is a good line.

Suppose that $V = V_7$, i.e., the blow-up of \mathbb{P}^3 at a point. Then S is a del Pezzo surface S_7 , i.e., a blow-up of \mathbb{P}^2 at two distinct points, and hence there exist three lines ℓ_0, ℓ_1, ℓ_2 on S as in Figure 2.1. Here ℓ_0 is distinguished by the fact that it intersects both of the other lines. Let P be the exceptional divisor of the blow-up $V_7 \rightarrow \mathbb{P}^3$. Then $P \simeq \mathbb{P}^2$ is a unique plane on V_7 and ℓ_0 is the intersection of S with P (cf. [7, Chap. II, §1.4]). Since $N_{\ell_0/P} \simeq \mathcal{O}_{\mathbb{P}^1}(1)$, ℓ_0 is a bad line on V_7 . On the other hand, ℓ_1 and ℓ_2 are good lines on V_7 since S is general. □

2.3. Infinitesimal deformations and obstructions

Let V be a smooth variety and let X be a smooth closed subvariety of V . An (embedded) first order infinitesimal deformation of X in V is a closed subscheme $\tilde{X} \subset V \times \text{Spec } k[t]/(t^2)$ which is flat over $\text{Spec } k[t]/(t^2)$ and whose central fiber is X . It is well known that there exists a one to one correspondence between the group of homomorphisms $\alpha : \mathcal{I}_X \rightarrow \mathcal{O}_X$ and

the first order infinitesimal deformations \tilde{X} of X in V . In what follows, we identify \tilde{X} with α and abuse the notation. The standard exact sequence

$$(2.4) \quad 0 \longrightarrow \mathcal{I}_X \longrightarrow \mathcal{O}_V \longrightarrow \mathcal{O}_X \longrightarrow 0$$

induces $\delta : \text{Hom}(\mathcal{I}_X, \mathcal{O}_X) \rightarrow \text{Ext}^1(\mathcal{I}_X, \mathcal{I}_X)$ as a coboundary map. Then $\alpha \in \text{Hom}(\mathcal{I}_X, \mathcal{O}_X)$ (i.e., \tilde{X}) lifts to a deformation over $\text{Spec } k[t]/(t^3)$ if and only if

$$\text{ob}(\alpha) := \delta(\alpha) \cup \alpha \in \text{Ext}^1(\mathcal{I}_X, \mathcal{O}_X)$$

is zero, where \cup is the cup product map

$$\text{Ext}^1(\mathcal{I}_X, \mathcal{I}_X) \times \text{Hom}(\mathcal{I}_X, \mathcal{O}_X) \xrightarrow{\cup} \text{Ext}^1(\mathcal{I}_X, \mathcal{O}_X).$$

(We refer to [11, Chap. I §2]. See also [15], [1], [3] and [10].) Then $\text{ob}(\alpha)$ is called the *obstruction* of α (i.e., \tilde{X}). Since both X and V are smooth, $\text{ob}(\alpha)$ is contained in $H^1(X, N_{X/V}) \subset \text{Ext}^1(\mathcal{I}_X, \mathcal{O}_X)$ (cf. [11, Chap. I, Prop. 2.14]). Since $\text{Hom}(\mathcal{I}_X, \mathcal{O}_X) \simeq H^0(N_{X/V})$, we regard α as a global section of $N_{X/V}$ from now on.

If X is a hypersurface of V , i.e., of codimension one in V , then $\text{ob}(\alpha)$ becomes a simple cup product. Let $\delta_1 : H^0(X, N_{X/V}) \rightarrow H^1(V, \mathcal{O}_V)$ be the coboundary map of the exact sequence $0 \rightarrow \mathcal{O}_V \rightarrow \mathcal{O}_V(X) \rightarrow N_{X/V} \rightarrow 0$. Let us define a map⁽⁴⁾

$$(2.5) \quad d_X : H^0(X, N_{X/V}) \longrightarrow H^1(X, \mathcal{O}_X)$$

by the composition of δ_1 and the restriction map $H^1(\mathcal{O}_V) \xrightarrow{|_X} H^1(\mathcal{O}_X)$. Then we have

LEMMA 2.9. — *Let X be a smooth hypersurface of V . Then $\text{ob}(\alpha)$ for $\alpha \in H^0(N_{X/V})$ is equal to the cup product $d_X(\alpha) \cup \alpha$, where \cup is the cup product map*

$$H^1(X, \mathcal{O}_X) \times H^0(X, N_{X/V}) \xrightarrow{\cup} H^1(X, N_{X/V}).$$

Proof. — Since $\mathcal{I}_X \simeq \mathcal{O}_V(-X)$ is a line bundle on V , we have $\text{Ext}^i(\mathcal{I}_X, \mathcal{O}_X) \simeq H^i(N_{X/V})$ ($i = 0, 1$) and $\text{Ext}^1(\mathcal{I}_X, \mathcal{I}_X) \simeq H^1(\mathcal{O}_V)$. Hence the coboundary map δ appearing in the definition of $\text{ob}(\alpha)$ is nothing but the coboundary map δ_1 of (2.4) $\otimes \mathcal{O}_V(X)$. Since α is a cohomology class on X , the cup product map $H^1(\mathcal{O}_V) \rightarrow H^1(N_{X/V})$ with α factors through the restriction map $|_X$. □

⁽⁴⁾ The map d_X is equal to the map $d_{X, \mathcal{O}_V(X)}$ defined in [13, §2.1 (2.3)].

We recall the definition of exterior component introduced in [13]. Let X be a smooth closed subvariety of V and let Y be a smooth hypersurface of V containing X . Then the natural projection $\pi_{X/Y} : N_{X/V} \rightarrow N_{Y/V}|_X \simeq \mathcal{O}_X(Y)$ of normal bundles induces the maps $H^i(\pi_{X/Y}) : H^i(N_{X/V}) \rightarrow H^i(N_{Y/V}|_X)$, where $i = 0, 1$, of their cohomology groups. Let α be a global section of $N_{X/V}$.

DEFINITION 2.10. — $\pi_{X/Y}(\alpha)$ and $\text{ob}_Y(\alpha)$ denote the images of α and $\text{ob}(\alpha)$ by the maps $H^0(\pi_{X/Y})$ and $H^1(\pi_{X/Y})$, respectively. They are called the exterior components of α and $\text{ob}(\alpha)$, respectively.

Roughly speaking, $\pi_{X/Y}(\alpha)$ is the projection of the normal vector α of X in V onto the normal directions to Y in V . Then $\text{ob}_Y(\alpha)$ represents the obstruction to deforming X into this direction. We recall a basic fact on exterior components.

LEMMA 2.11 ([13, Lemma 2.4]). — Let $\pi_{X/Y}(\alpha)$ and $\text{ob}_Y(\alpha)$ be the exterior components of α and $\text{ob}(\alpha)$, respectively. If there exists a global section v of $N_{Y/V}$ whose restriction $v|_X$ to X coincides with $\pi_{X/Y}(\alpha)$, then we have

$$\text{ob}_Y(\alpha) = \text{ob}(v)|_X$$

where $\text{ob}(v)|_X \in H^1(X, N_{Y/V}|_X)$ is the restriction of $\text{ob}(v) \in H^1(Y, N_{Y/V})$ to X .

Lemma 2.11 together with Lemma 2.9 shows that $\text{ob}_Y(\alpha) = d_Y(v)|_X \cup \pi_{X/Y}(\alpha)$, where d_Y is the map (2.5) for Y and \cup is the cup product map

$$(2.6) \quad H^1(X, \mathcal{O}_X) \times H^0(X, N_{Y/V}|_X) \xrightarrow{\cup} H^1(X, N_{Y/V}|_X).$$

Let E be an effective divisor of Y disjoint to X (i.e., $X \cap E = \emptyset$). Let Y° and V° denote the two complements of E in Y and V , respectively. Every rational section v of $N_{Y/V} \simeq \mathcal{O}_Y(Y)$ having poles only along E determines a global section v° of the normal bundle N_{Y°/V° of Y° in V° and hence the obstruction $\text{ob}(v^\circ) \in H^1(N_{Y^\circ/V^\circ})$ to deforming Y° in V° . Let ι denote the open immersion of $Y^\circ \hookrightarrow Y$. Then a natural homomorphism $\iota_* N_{Y^\circ/V^\circ} \rightarrow N_{Y/V}|_X (= [\iota_* \mathcal{O}_{Y^\circ} \rightarrow \mathcal{O}_X] \otimes N_{Y/V})$ of sheaves on Y induces a map $H^1(N_{Y^\circ/V^\circ}) \xrightarrow{|\iota} H^1(N_{Y/V}|_X)$. Since $\text{ob}(\alpha)$ is (and hence $\text{ob}_Y(\alpha)$ is) determined by a neighborhood of X , we have the following variant of Lemma 2.11.

LEMMA 2.12. — Let α be a global section of $N_{X/V}$. If there exists a rational section v of $N_{Y/V}$ whose only poles are along E and whose restriction

to X coincides with $\pi_{X/Y}(\alpha)$, then we have

$$\text{ob}_Y(\alpha) = \text{ob}(v^\circ)|_X,$$

where $\text{ob}(v^\circ)|_X$ is the image of $\text{ob}(v^\circ)$ by the map $H^1(Y^\circ, N_{Y^\circ/V^\circ}) \xrightarrow{|\chi} H^1(X, N_{Y/V}|_X)$.

3. Infinitesimal deformations with a pole

Let V be a smooth projective 3-fold, S a smooth surface in V , E a smooth connected curve on S . We put $V^\circ := V \setminus E$ and $S^\circ := S \setminus E$, the complementary open subvarieties. In this section, we study the first order infinitesimal deformations of S° in V° , when the self-intersection number of E on S is negative. We are interested in a rational section v of $N_{S/V}$ having a pole only along E and of order one, that is, $v \in H^0(N_{S/V}(E)) \setminus H^0(N_{S/V})$. Let $\iota : S^\circ \hookrightarrow S$ be the open immersion. Then $\iota_*\mathcal{O}_{S^\circ}$ contains $\mathcal{O}_S(nE)$ as a subsheaf for any $n \geq 0$. Hence the natural sheaf injection $N_{S/V}(nE) \hookrightarrow \iota_*N_{S^\circ/V^\circ}$ induces $H^0(S, N_{S/V}(nE)) \hookrightarrow H^0(S^\circ, N_{S^\circ/V^\circ})$ for each n . Therefore v determines a first order infinitesimal deformation of S° in V° . The main theorem of this section is the following.

THEOREM 3.1. — *Let v be as above and assume that $E^2 < 0$ and $\det N_{E/V} := \bigwedge^2 N_{E/V}$ is trivial. If the exact sequence*

$$(3.1) \quad 0 \longrightarrow N_{E/S} \longrightarrow N_{E/V} \longrightarrow N_{S/V}|_E \longrightarrow 0$$

does not split, then the first order infinitesimal deformation of $S^\circ \subset V^\circ$ determined by v does not lift to a deformation over $\text{Spec } k[t]/(t^3)$.

Let n be a non-negative integer. In what follows, we identify $H^0(N_{S/V}(nE))$ with its image in $H^0(N_{S^\circ/V^\circ})$. We shall prove that the obstruction $\text{ob}(v)$ is nonzero in $H^1(N_{S^\circ/V^\circ})$. Let d_{S° denote the map (2.5) for $X = S^\circ$. Then by Lemma 2.9, $\text{ob}(v)$ is equal to the cup product $d_{S^\circ}(v) \cup v$, where \cup is the cup product map

$$H^1(S^\circ, \mathcal{O}_{S^\circ}) \times H^0(S^\circ, N_{S^\circ/V^\circ}) \xrightarrow{\cup} H^1(S^\circ, N_{S^\circ/V^\circ}).$$

The inclusion $\mathcal{O}_S(nE) \hookrightarrow \iota_*\mathcal{O}_{S^\circ}$ of sheaves induces a map $H^1(S, \mathcal{O}_S(nE)) \rightarrow H^1(S^\circ, \mathcal{O}_{S^\circ})$ of cohomology groups. Suppose that $E^2 < 0$. Then this map is injective by Lemma 2.6. Hence we identify $H^1(\mathcal{O}_S(nE))$ with its image in $H^1(S^\circ, \mathcal{O}_{S^\circ})$. Under this identification, there exists a natural filtration

$$H^1(S, \mathcal{O}_S) \subset H^1(S, \mathcal{O}_S(E)) \subset H^1(S, \mathcal{O}_S(2E)) \subset \dots \subset H^1(S^\circ, \mathcal{O}_{S^\circ})$$

on $H^1(S^\circ, \mathcal{O}_{S^\circ})$. Suppose now that $\det N_{E/V}$ is trivial. Then under similar identifications, there exists a natural filtration

$$H^1(S, N_{S/V}(E)) \subset H^1(S, N_{S/V}(2E)) \subset \cdots \subset H^1(S^\circ, N_{S^\circ/V^\circ})$$

on $H^1(S^\circ, N_{S^\circ/V^\circ})$, because we have $\deg N_{S/V}(nE)|_E = \deg(\det N_{E/V}) + (n - 1)E^2 = (n - 1)E^2 \leq 0$ for $n \geq 1$. Then it follows from [13, Proposition 2.6 (1)] that the image of d_{S° over $H^0(N_{S/V}(E))$ is contained in $H^1(\mathcal{O}_S(2E))$. By the commutative diagram

$$\begin{array}{ccc} H^1(\mathcal{O}_{S^\circ}) & \times & H^0(N_{S^\circ/V^\circ}) & \xrightarrow{\cup} & H^1(N_{S^\circ/V^\circ}) \\ \cup & & \cup & & \cup \\ H^1(\mathcal{O}_S(2E)) & \times & H^0(N_{S/V}(E)) & \xrightarrow{\cup} & H^1(N_{S/V}(3E)), \end{array}$$

the image of the obstruction map ob over $H^0(N_{S/V}(E))$ is contained in $H^1(N_{S/V}(3E))$. The following lemma is essential to the proof of Theorem 3.1. Let d_S denote the restriction of the map d_{S° to $H^0(S, N_{S/V}(E))$.

LEMMA 3.2 ([13, Proposition 2.6 (2)]). — *Let $\partial : H^0(N_{S/V}(E)|_E) \rightarrow H^1(\mathcal{O}_E(2E)) \simeq H^1(N_{E/S}(E))$ be the coboundary map of the exact sequence (3.1) $\otimes \mathcal{O}_S(E)$. Then the diagram*

$$\begin{array}{ccc} H^0(S, N_{S/V}(E)) & \xrightarrow{d_S} & H^1(S, \mathcal{O}_S(2E)) \\ \downarrow |_E & & \downarrow |_E \\ H^0(E, N_{S/V}(E)|_E) & \xrightarrow{\partial} & H^1(E, \mathcal{O}_E(2E)) \end{array}$$

is commutative.

Proof of Theorem 3.1. It suffices to show that the image $\text{ob}(v)|_E \in H^1(N_{S/V}(3E)|_E)$ of $\text{ob}(v) \in H^1(N_{S/V}(3E))$ is nonzero. By the definition of v , we have $v|_E \neq 0$ in $H^0(N_{S/V}(E)|_E)$. Then the line bundle $N_{S/V}(E)|_E \simeq \det N_{E/V}$ on E is trivial. Since (3.1) does not split by assumption, we have $\partial(v|_E) \neq 0$. Hence by Lemma 3.2, we conclude that

$$\text{ob}(v)|_E = d_{S^\circ}(v)|_E \cup v|_E = \partial(v|_E) \cup v|_E \neq 0. \quad \square$$

If E is a $(-1)\text{-}\mathbb{P}^1$ on S with $\det N_{E/V} \simeq \mathcal{O}_{\mathbb{P}^1}$, then the exact sequence (3.1) does not split if and only if $N_{E/V}$ is trivial.

Example 3.3. — Let V_n be a smooth del Pezzo 3-fold of degree $n \neq 8$ and let E be a good line on V_n , i.e., N_{E/V_n} is trivial (cf. §2.2). If S_n is a smooth hyperplane section of V_n containing E , then there exists an obstructed infinitesimal deformation of $S_n^\circ := S_n \setminus E$ in $V_n^\circ := V_n \setminus E$. Indeed, let $\varepsilon : S_n \rightarrow S_{n+1}$ be the blow-down of E from S_n . Since $N_{S_n/V_n} \simeq -K_{S_n}$, $N_{S_n/V_n}(E) \simeq \varepsilon^*(-K_{S_{n+1}})$, and $h^0(-K_{S_{n+1}}) > h^0(-K_{S_n})$, there exists a global section $v \in H^0(N_{S_n/V_n}(E)) \setminus H^0(N_{S_n/V_n})$. Then by Theorem 3.1,

the first order infinitesimal deformation of S_n° in V_n° determined by v is obstructed.

In the rest of this section, we discuss a generalization of Theorem 3.1, which will be used in the proof of Theorem 1.3. Let E be a disjoint union of smooth connected curves E_i ($i = 1, \dots, m$) on S such that $E_i^2 < 0$ and $\det N_{E_i/V}$ is trivial. By the same symbol E we also denote the divisor $\sum_{i=1}^m E_i$ on S . Let us define V° and S° as above and identify $H^0(N_{S/V}(E))$ with its image in $H^0(N_{S^\circ/V^\circ})$. We compute the restriction to $H^0(N_{S/V}(E))$ of the obstruction map $\text{ob} : H^0(N_{S^\circ/V^\circ}) \rightarrow H^1(N_{S^\circ/V^\circ})$. Lemma 2.6 allows us to regard $H^1(\mathcal{O}_S(2E))$ and $H^1(N_{S/V}(3E))$ as subgroups of $H^1(\mathcal{O}_{S^\circ})$ and $H^1(N_{S^\circ/V^\circ})$, respectively. Then an argument similar to [13, Proposition 2.6 (1)] shows that the image of $H^0(N_{S/V}(E))$ by d_{S° is contained in $H^1(\mathcal{O}_S(2E))$. Therefore we conclude that

LEMMA 3.4. — *The image of $H^0(N_{S/V}(E))$ by ob is contained in $H^1(N_{S/V}(3E)) \subset H^1(N_{S^\circ/V^\circ})$.*

Let v and v' be any global sections of $N_{S/V}(E)$ and $N_{S/V}$, respectively. Then we have $\text{ob}(v+v')|_E = \text{ob}(v)|_E$ in $H^1(N_{S/V}(3E))|_E$. Indeed it follows from the definition of d_{S° (cf. (2.5)) that $d_{S^\circ}(v')$ is contained in $H^1(\mathcal{O}_S)$ and hence

$$\begin{aligned} \text{ob}(v+v') &= (d_{S^\circ}(v) + d_{S^\circ}(v')) \cup (v+v') \\ &= \text{ob}(v) + \underbrace{d_{S^\circ}(v) \cup v' + d_{S^\circ}(v') \cup v + d_{S^\circ}(v') \cup v'}_{\text{contained in } H^1(N_{S/V}(2E))}. \end{aligned}$$

Therefore the obstruction map ob induces a map

$$(3.2) \quad \overline{\text{ob}} : H^0(N_{S/V}(E))/H^0(N_{S/V}) \longrightarrow H^1(N_{S/V}(3E))|_E.$$

PROPOSITION 3.5. — *If $H^1(N_{S/V}) = 0$ and the exact sequence*

$$(3.3) \quad 0 \longrightarrow N_{E_i/S} \longrightarrow N_{E_i/V} \longrightarrow N_{S/V}|_{E_i} \longrightarrow 0$$

does not split for every i , then $\overline{\text{ob}}$ is injective.

This is an immediate consequence of the next lemma.

LEMMA 3.6. — *Under the assumption of Proposition 3.5, $\overline{\text{ob}}$ is equivalent to the quadratic map*

$$k^m \longrightarrow k^n, \quad (a_1, \dots, a_m) \longmapsto (a_1^2, \dots, a_m^2, 0, \dots, 0)$$

of diagonal type, where $n = \dim H^1(N_{S/V}(3E))|_E$.

Proof. — Since $H^1(N_{S/V}) = 0$, the source of the map $\overline{\text{ob}}$ is isomorphic to $H^0(N_{S/V}(E)|_E)$. Moreover there exist global sections v_i of $N_{S/V}(E_i)$ such that $v_i|_E \neq 0$ in $H^0(N_{S/V}(E_i)|_{E_i})$ for all i . Since E_i 's are mutually disjoint, we have $N_{S/V}(E)|_E \simeq \bigoplus_{i=1}^m N_{S/V}(E_i)|_{E_i} \simeq \bigoplus_{i=1}^m \mathcal{O}_{E_i}$. Then there exists a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & H^0(N_{S/V}) & \rightarrow & H^0(N_{S/V}(E)) & \rightarrow & H^0(N_{S/V}(E)|_E) \rightarrow 0 \\ & & \uparrow a_1 & & \uparrow a_2 & & \uparrow a_3 \\ 0 & \rightarrow & \bigoplus_i H^0(N_{S/V}) & \rightarrow & \bigoplus_i H^0(N_{S/V}(E_i)) & \rightarrow & \bigoplus_i H^0(N_{S/V}(E_i)|_{E_i}) \rightarrow 0, \end{array}$$

where the two horizontal sequences are exact and a_i ($1 \leq i \leq 3$) are defined by addition. Since a_1 and a_3 are surjective, so is a_2 . Hence every element $v \in H^0(N_{S/V}(E))$ is written as a k -linear combination $\sum_{i=1}^m c_i v_i$ of $v_i \in H^0(N_{S/V}(E_i))$ and the expression is unique modulo $H^0(N_{S/V})$. By the commutative diagram

$$\begin{array}{ccccc} H^1(\mathcal{O}_S(2E)) & \times & H^0(N_{S/V}(E)) & \xrightarrow{\cup} & H^1(N_{S/V}(3E)) \\ \downarrow |_E & & \downarrow |_E & & \downarrow |_E \\ \bigoplus_i H^1(\mathcal{O}_{E_i}(2E_i)) & \times & \bigoplus_i H^0(N_{S/V}(E_i)|_{E_i}) & \xrightarrow{\cup} & \bigoplus_i H^1(N_{S/V}(3E_i)|_{E_i}), \end{array}$$

we have

$$\text{ob}(v)|_E = (d_{S^\circ}(v) \cup v)|_E = d_{S^\circ}(v)|_E \cup v|_E = \sum_i c_i^2 d_{S^\circ}(v_i)|_{E_i} \cup v_i|_{E_i}.$$

By Lemma 3.2, $d_{S^\circ}(v_i)|_{E_i}$ is equal to $\partial_i(v|_{E_i})$ in $H^1(\mathcal{O}_{E_i}(2E_i))$, where ∂_i is the coboundary map of (3.3). Since (3.3) does not split by assumption, we have $\partial_i(v|_{E_i}) \neq 0$ and hence $d_{S^\circ}(v_i)|_{E_i} \neq 0$ for any i . As a result, $d_{S^\circ}(v_i)|_{E_i} \cup v_i|_{E_i}$ ($1 \leq i \leq m$) form a sub-basis of $H^1(N_{S/V}(3E)|_E)$. \square

Suppose now that E_i is a (-1) - \mathbb{P}^1 on S and $N_{E_i/V}$ is trivial for every $1 \leq i \leq m$. Then Proposition 3.5 shows that

COROLLARY 3.7. — *Let $v \in H^0(N_{S/V}(E)) \setminus H^0(N_{S/V})$ be a global section. If $H^1(N_{S/V}) = 0$, then we have $\text{ob}(v) \neq 0$ in $H^1(N_{S/V}(3E))$ ($\subset H^1(N_{S^\circ/V^\circ})$).*

4. Obstructions to deforming curves

Let V be a smooth projective 3-fold. In this section we study the deformation of smooth curves C on V under the presence of smooth surface S such that $C \subset S \subset V$.

4.1. S -normal curves and S -maximal families

In what follows, we assume that $\text{Hilb } V$ is nonsingular at $[S]$. Then there exists a unique irreducible component U_S of $\text{Hilb } V$ passing through $[S]$. We use the following convention.

DEFINITION 4.1.

- (1) C is said to be stably degenerate if there exists an (Zariski) open neighborhood $U \subset \text{Hilb } V$ of $[C]$ such that for any member $[C'] \in U$, there exists a deformation S' of S in V such that $C' \subset S'$ and $[S'] \in U_S$.
- (2) C is said to be S -normal if the restriction map (1.1) is surjective.

Let

$$V \times U_S \supset \mathcal{S} \xrightarrow{p_2} U_S$$

be the universal family of U_S . Let us denote the Hilbert scheme of smooth connected curves in \mathcal{S} by $\text{Hilb}^{sc} \mathcal{S}$, which is the relative Hilbert scheme of \mathcal{S}/U_S . $\text{Hilb}^{sc} \mathcal{S}$ is regarded as an open subscheme of the Hilbert-flag scheme of V (see [8] for the definition), which parametrizes all flat families of pairs (C, S) of a curve C and a surface S in V such that $C \subset S$. The projection $p_1 : \mathcal{S} \rightarrow V$ induces a natural morphism

$$(4.1) \quad pr_1 : \text{Hilb}^{sc} \mathcal{S} \longrightarrow \text{Hilb}^{sc} V,$$

which is the forgetful morphism $(C, S) \mapsto C$. Then by definition C is stably degenerate if and only if pr_1 is surjective in a neighborhood of $[C] \in \text{Hilb}^{sc} V$.

The next lemma plays an important role in our proof of Theorem 1.3 later (cf. § 4.3).

LEMMA 4.2. — Assume that $H^1(C, N_{C/S}) = 0$. Then:

- (1) The kernel and the cokernel of the tangential map

$$(4.2) \quad \kappa_{C,S} : H^0(C, N_{C/S}) \longrightarrow H^0(C, N_{C/V}).$$

of pr_1 at (C, S) are isomorphic to those of the restriction map (1.1), respectively.

- (2) $\text{Hilb}^{sc} \mathcal{S}$ is nonsingular at (C, S) .

For the proof we refer to [13, Lemma 3.1] for (1) and [9, Lemma 1.10] for (2). We can also prove (1) by using the “fundamental exact sequence relating $A^i(C \subset S)$ and $H^{i-1}(N_{C/V})$ ” in [9].

In what follows, we assume that $H^1(C, N_{C/S}) = 0$. If C is S -normal, then $\kappa_{C,S}$ is surjective by Lemma 4.2 (1). Then by (2) of the same lemma,

$\text{Hilb}^{sc} V$ is nonsingular at $[C]$ and furthermore pr_1 is surjective in a neighborhood of $[C]$. In fact, if C is S -normal, then the morphism pr_1 is smooth at (C, S) (cf. [8, Lemma A10]). Thus we conclude that

PROPOSITION 4.3 (cf. [8],[9]). — *If C is S -normal, then C is stably degenerate and $\text{Hilb}^{sc} V$ is nonsingular at $[C]$.*

We recall the S -maximal family introduced in [13, §3.2]. By the smoothness of $\text{Hilb}^{sc} \mathcal{S}$, there exists a unique irreducible component $W_{S,C}$ of $\text{Hilb}^{sc} \mathcal{S}$ containing (C, S) .

DEFINITION 4.4. — *We define the S -maximal family of curves containing C to be the image of $W_{S,C}$ in $\text{Hilb}^{sc} V$ and denote it by $W_{S,C}$.*

By Proposition 4.3, if C is S -normal then $W_{S,C}$ is an irreducible component of $\text{Hilb}^{sc} V$ and $\text{Hilb}^{sc} V$ is generically smooth along $W_{S,C}$.

4.2. Deformation of curves on a del Pezzo 3-fold

Let V be a smooth del Pezzo 3-fold with the polarization H , S a smooth member of $|H|$, and C a smooth connected curve on S . Let n denote the degree of V and let d and g denote the degree $(:= (C \cdot H)_V)$ and the genus of C , respectively.

By adjunction we have $N_{S/V} \simeq -K_V|_S + K_S$ and $N_{C/S} \simeq -K_S|_C + K_C$. Since $-K_V$ and $-K_S$ are ample, we have $H^1(N_{S/V}) = H^1(N_{C/S}) = 0$. Hence $\text{Hilb} V$ and $\text{Hilb} S$ are nonsingular at $[S]$ and $[C]$, respectively. Thus if C is S -normal, then by Proposition 4.3, C is stably degenerate and $\text{Hilb}^{sc} V$ is nonsingular at $[C]$. Because $H^1(N_{S/V}) = 0$, it follows from the exact sequence

$$(4.3) \quad 0 \longrightarrow N_{S/V}(-C) \longrightarrow N_{S/V} \longrightarrow N_{S/V}|_C \longrightarrow 0$$

that C is S -normal if and only if $H^1(N_{S/V}(-C)) = 0$. There exists a natural exact sequence

$$(4.4) \quad 0 \longrightarrow N_{C/S} \longrightarrow N_{C/V} \xrightarrow{\pi_{C/S}} N_{S/V}|_C \longrightarrow 0.$$

Since $H^1(N_{C/S}) = 0$, we have $H^1(N_{C/V}) \simeq H^1(N_{S/V}|_C)$. Thus every obstruction to deforming C is contained in the cohomology group $H^1(N_{S/V}|_C)$. Since $\chi(N_{C/V}) = (-K_V \cdot C)_V = 2d$, we also have

LEMMA 4.5. — *If $H^1(N_{S/V}|_C) = 0$, then $\text{Hilb}^{sc} V$ is nonsingular of expected dimension $2d$ at $[C]$.*

In particular, if C is rational ($g = 0$) or elliptic ($g = 1$), then the $\text{Hilb}^{sc} V$ is nonsingular at $[C]$ because $H^1(N_{S/V}|_C) \simeq H^1(-K_S|_C) = 0$.

Let $W_{S,C}$ be the S -maximal family $W_{S,C}$ of curves containing C . We compute the dimension of $W_{S,C}$. Let $pr_1 : \text{Hilb}^{sc} \mathcal{S} \rightarrow \text{Hilb}^{sc} V$ be the morphism (4.1).

LEMMA 4.6.

- (1) $\text{Hilb}^{sc} \mathcal{S}$ is nonsingular of dimension $d + g + n$ at (C, S) .
- (2) If $g \geq 2$ or $d \geq n + 1$, then pr_1 is a closed embedding in a neighborhood of (C, S) and $\dim W_{S,C} = d + g + n$.

Proof. — (1) Let $\mathcal{W}_{S,C}$ be the irreducible component of $\text{Hilb}^{sc} \mathcal{S}$ containing (C, S) . By the Riemann-Roch theorem on S , we have $\dim |\mathcal{O}_S(C)| = d + g - 1$. Then $\mathcal{W}_{S,C}$ is birationally equivalent to \mathbb{P}^{d+g-1} -bundle over an open subset of $|H| \simeq \mathbb{P}^{n+1}$. Hence we have $\dim \mathcal{W}_{S,C} = d + g + n$.

(2) By assumption, we have $(-K_S - C) \cdot C = 2 - 2g < 0$ or $(-K_S - C) \cdot (-K_S) = n - d < 0$. Since both C and $-K_S$ are nef, we have $H^0(N_{S/V}(-C)) \simeq H^0(-K_S - C) = 0$. By Lemma 4.2 (1), pr_1 is a closed embedding near (C, S) . Hence we have $\dim W_{S,C} = \dim \mathcal{W}_{S,C}$. □

We denote by $\text{Hilb}_{d,g}^{sc} V$ the open and closed subscheme of $\text{Hilb}^{sc} V$ of curves of degree d and genus g . It is known that the dimension of every irreducible component of $\text{Hilb}_{d,g}^{sc} V$ is greater than or equal to the expected dimension $\chi(N_{C/V}) = 2d$ (cf. [11, Chap. I, Theorem 2.8]).

PROPOSITION 4.7. — *If $\chi(V, \mathcal{I}_C(S)) < 1$, then C is not stably degenerate, i.e., there exists a deformation C' of C in V which is not contained in any deformation S' of S in V .*

Proof. — There exists an exact sequence $[0 \rightarrow \mathcal{I}_C \rightarrow \mathcal{O}_V \rightarrow \mathcal{O}_C \rightarrow 0] \otimes \mathcal{O}_V(S)$ on V . We see that $\chi(\mathcal{O}_C(S)) = d + 1 - g$ and $\chi(\mathcal{O}_V(S)) = n + 2$. Hence $\chi(\mathcal{I}_C(S)) < 1$ is equivalent to $g < d - n$. Then we have $\dim W_{S,C} \leq \dim \mathcal{W}_{S,C} = d + g + n < 2d$. Hence there exists an irreducible component $W' \supset W_{S,C}$ of $\text{Hilb}^{sc} V$ such that $\dim W' > \dim W_{S,C}$. A general member C' of $W' \setminus W_{S,C}$ is such a deformation of C in V . □

4.3. Stably degenerate curves

We devote this subsection to the proof of Theorem 1.3. Notation is the same as in the previous subsection. The following are equivalent: (i) $\chi(V, \mathcal{I}_C(S)) \geq 1$, (ii) $\chi(S, N_{S/V}(-C)) \geq 0$ and (iii) $g \geq d - n$. Indeed

we have already seen in the proof of Proposition 4.7 that (i) and (iii) are equivalent. Also (i) and (ii) are equivalent because we have $\chi(N_{S/V}(-C)) = \chi(\mathcal{I}_C(S)) - 1$ by the exact sequence

$$(4.5) \quad [0 \longrightarrow \mathcal{I}_S \longrightarrow \mathcal{I}_C \longrightarrow \mathcal{O}_S(-C) \longrightarrow 0] \otimes \mathcal{O}_V(S).$$

Throughout this subsection, we assume one of (i),(ii) and (iii) (and hence all).

LEMMA 4.8. — *If $H^1(C, N_{S/V}|_C) = 0$ then C is S -normal.*

Proof. — It suffices to show that $H^1(N_{S/V}(-C)) = 0$. Since $H^2(N_{S/V}) \simeq H^2(-K_S) = 0$ and $H^1(N_{S/V}|_C) = 0$, we obtain $H^2(N_{S/V}(-C)) = 0$ by (4.3). Then by assumption, we have $0 \leq \chi(N_{S/V}(-C)) = h^0(N_{S/V}(-C)) - h^1(N_{S/V}(-C))$. Therefore if $H^0(N_{S/V}(-C)) = 0$, then we have $H^1(N_{S/V}(-C)) = 0$. Suppose that $H^0(N_{S/V}(-C)) \neq 0$. There exists an effective divisor D on S such that $N_{S/V}(-C) \simeq \mathcal{O}_S(D)$. If $D = 0$, then $H^1(N_{S/V}(-C)) = 0$. Suppose that $D \neq 0$. Let h be a general member of $| -K_S |$. Then h is a smooth elliptic curve on S . Since $-K_S$ is ample, we have $\text{deg } \mathcal{O}_S(D)|_h = D \cdot (-K_S) > 0$ and hence $H^1(\mathcal{O}_S(D)|_h) = 0$. Since C is connected, we obtain $H^1(D - h) \simeq H^1(-C) = 0$ from the exact sequence $0 \rightarrow \mathcal{O}_S(-C) \rightarrow \mathcal{O}_S \rightarrow \mathcal{O}_C \rightarrow 0$. Therefore it follows from the exact sequence

$$0 \longrightarrow \mathcal{O}_S(D - h) \longrightarrow \mathcal{O}_S(D) \longrightarrow \mathcal{O}_S(D)|_h \longrightarrow 0$$

that $H^1(N_{S/V}(-C)) \simeq H^1(D) = 0$. □

Let E_1, \dots, E_m be lines on S disjoint to C . We define an effective divisor E on S by $E := \sum_{i=1}^m E_i$. If C is not S -normal, then E is responsible for the abnormality.

PROPOSITION 4.9. — *Suppose that C is not rational nor elliptic.*

- (1) *The restriction map $H^0(S, N_{S/V}(E)) \xrightarrow{|_C} H^0(C, N_{S/V}|_C)$ is an isomorphism.*
- (2) *C is S -normal if and only if there exists no line ℓ such that $C \cap \ell = \emptyset$ (i.e., $E = 0$).*

Proof. — (1) Since $N_{S/V} \simeq -K_S$, we have the assertion by Proposition 2.4.

(2) The “if” part follows from (1). We prove the “only if” part. Suppose that there exist such lines on S . Let $\varepsilon : S \rightarrow F$ be the blow-down of E from S . Then F is also a del Pezzo surface and $\varepsilon^*(-K_F) = -K_S + E$. Since $\text{deg } F > \text{deg } S$, we have $h^0(-K_F) > h^0(-K_S)$. Hence it follows from $N_{S/V} \simeq -K_S$ that $N_{S/V}(E)$ has more global sections than $N_{S/V}$. Hence

we have $h^0(N_{S/V}|_C) = h^0(N_{S/V}(E)) > H^0(N_{S/V})$ by (1). Therefore C is not S -normal. □

Let $\kappa_{C,S} : H^0(N_{C/S}) \rightarrow H^0(N_{C/V})$ denote the tangential map (4.2).

PROPOSITION 4.10. — *Suppose that C is not S -normal. If every E_i is a good line on V , then the obstruction $\text{ob}(\alpha)$ is nonzero for any $\alpha \in H^0(C, N_{C/V}) \setminus \text{im } \kappa_{C,S}$.*

Proof. — Let $\pi_{C/S}(\alpha) \in H^0(N_{S/V}|_C)$ and $\text{ob}_S(\alpha) \in H^1(N_{S/V}|_C)$ be the exterior component of α and $\text{ob}(\alpha)$, respectively (cf. Definition 2.10). We compute $\text{ob}_S(\alpha)$ instead of $\text{ob}(\alpha)$ itself. Since C is not S -normal, by Lemma 4.8, we have $H^1(N_{S/V}|_C) \neq 0$. In particular, C is not rational nor elliptic. By Proposition 4.9 (1), there exists a global section v of $N_{S/V}(E)$ whose restriction $v|_C \in H^0(N_{S/V}|_C)$ to C coincides with $\pi_{C/S}(\alpha)$. Since α is not contained in the image of $\kappa_{C,S}$, $\pi_{C/S}(\alpha)$ is not contained in the image of (1.1) by Lemma 4.2 (1). Hence v is not a global section of $N_{S/V}$, in other words, an infinitesimal deformation with a pole (cf. §3).

Let S° and V° respectively denote the two complements $S \setminus E$ and $V \setminus E$ of E . There exists a natural injection $H^0(S, N_{S/V}(E)) \hookrightarrow H^0(S^\circ, N_{S^\circ/V^\circ})$ of cohomology groups. In what follows, we identify $v \in H^0(S, N_{S/V}(E))$ with its image $v^\circ \in H^0(S^\circ, N_{S^\circ/V^\circ})$, i.e., a first order infinitesimal deformation of S° in V° . Now we prove that the obstruction $\text{ob}(v) \in H^1(S^\circ, N_{S^\circ/V^\circ})$ is nonzero. By Lemma 3.4, $\text{ob}(v)$ is contained in the subgroup $H^1(S, N_{S/V}(3E))$ of $H^1(S^\circ, N_{S^\circ/V^\circ})$. Every component E_i of E is a $(-1)\text{-}\mathbb{P}^1$ on S and its normal bundle $N_{E_i/V}$ in V is trivial by assumption. Therefore by virtue of Corollary 3.7, we have $\text{ob}(v) \neq 0$ in $H^1(S, N_{S/V}(3E))$.

Finally we show that $\text{ob}_S(\alpha) \neq 0$ in $H^1(C, N_{S/V}|_C)$. There exists an exact sequence

$$0 \longrightarrow N_{S/V}(3E - C) \longrightarrow N_{S/V}(3E) \xrightarrow{|_C} N_{S/V}|_C \longrightarrow 0.$$

Since $N_{S/V} \simeq -K_S$, the restriction map $H^1(S, N_{S/V}(3E)) \rightarrow H^1(C, N_{S/V}|_C)$ is injective by Lemma 2.5. Therefore we have $\text{ob}_S(\alpha) = \text{ob}(v)|_C \neq 0$ by Lemma 2.12. □

Now we prove Theorem 1.3. Let C be as in the theorem. Then we have

LEMMA 4.11. — *Every small global deformation of C in V is contained in the S -maximal family $W_{S,C}$ of curves containing C .*

Proof. — Let $C_T \subset V \times T$ be a small global deformation of C , i.e., a flat family C_T over a small open variety T , having a point $0 \in T$ with $C_0 = C$. Given an element of the Zariski tangent space of T at 0 , we obtain

a morphism $\text{Spec } k[t]/(t^2) \rightarrow T$ and a first order infinitesimal deformation $\tilde{C} \rightarrow \text{Spec } k[t]/(t^2)$ of C by base extension. Then by Proposition 4.10 \tilde{C} is contained in the image of the map $\kappa_{(C,S)}$. Hence there exists a first order infinitesimal deformation \tilde{S} of S such that $\tilde{S} \supset \tilde{C}$. Since $\text{Hilb}^{sc} S$ is nonsingular at (C, S) , the first order infinitesimal deformation (\tilde{C}, \tilde{S}) of (C, S) lifts to a global deformation (C_T, S_T) over T . \square

Therefore C is stably degenerate. The rest of the proof is as follows. If C is S -normal, then $\text{Hilb}^{sc} V$ is nonsingular at $[C]$ by Proposition 4.3. Otherwise, there exists a first order infinitesimal deformation \tilde{C} of C not contained in the image of $\kappa_{(C,S)}$. Then $\text{Hilb}^{sc} V$ is singular at $[C]$ by Proposition 4.10. We have an isomorphism $H^1(S, N_{S/V}(-C)) \simeq H^1(V, \mathcal{I}_C(S))$ by the exact sequence (4.5) together with that $H^i(V, \mathcal{I}_S(S)) = H^i(V, \mathcal{O}_V) = 0$ for $i = 1, 2$. Hence C is S -normal if and only if $H^1(V, \mathcal{I}_C(S)) = 0$. The proof of Theorem 1.3 has been completed.

Remark 4.12. — We give two remarks on Theorem 1.3.

- (1) Suppose that V is not isomorphic to a blow-up V_7 of \mathbb{P}^3 at a point. If $S \in |H|$ is general, then by Lemma 2.8, every line on S is a good line on V . Hence every curve C on S is stably degenerate by the theorem. Meanwhile there exists a non-stably degenerate curve C on V_7 which is contained in a general member S of $|H|$ (cf. Proposition 5.4).
- (2) There exists no line on a del Pezzo 3-fold $V_8 \simeq \mathbb{P}^3$. Hence if $V = V_8$, then the assumption of the theorem concerning lines ℓ on S such that $C \cap \ell = \emptyset$ is empty. In fact, if $g \geq d - 8$ then every curve C on V_8 is S -normal and hence stably degenerate. This coincides with the previous result [15, Appendix, Proposition 4.11], which proved that every curve of degree e and genus $p \geq 2e - 8$ in \mathbb{P}^3 lying on a smooth quadric surface $Q_2 \simeq \mathbb{P}^1 \times \mathbb{P}^1$ is stably degenerate.

The following proposition is more practical than Proposition 4.10 in showing that $\text{Hilb}^{sc} V$ is singular at $[C]$.

PROPOSITION 4.13. — *Suppose that C is not rational nor elliptic. If there exists a good line ℓ on V such that $\ell \subset S$ and $C \cap \ell = \emptyset$, then $\text{Hilb}^{sc} V$ is singular at $[C]$.*

The proofs of Proposition 4.10 and Proposition 4.13 are very similar. Take a global section $v \in H^0(N_{S/V}(\ell)) \setminus H^0(N_{S/V})$ and put $\alpha \in H^0(N_{C/V})$ as a lift of $v|_C \in H^0(N_{S/V})$ by the surjective map $\pi_{C/S} : H^0(N_{C/V}) \rightarrow H^0(N_{S/V}|_C)$. Then it is enough to show that $\text{ob}_S(\alpha) \neq 0$ in $H^1(N_{S/V}|_C)$

by reducing it to $\text{ob}(v)|_\ell \neq 0$ as in the proof of Proposition 4.10. We omit the details.

The following is an analogue of Conjecture 5.1 due to Kleppe and Ellia.

THEOREM 4.14. — *Let C be the curve in Theorem 1.3. Then:*

- (1) *The S -maximal family $W_{S,C} \subset \text{Hilb}^{sc} V$ of curves containing C is an irreducible component of $(\text{Hilb}^{sc} V)_{\text{red}}$.*
- (2) *$\text{Hilb}^{sc} V$ is generically smooth along $W_{S,C}$ if $H^1(V, \mathcal{I}_C(S)) = 0$, and generically non-reduced along $W_{S,C}$ otherwise.*

Proof. — (1) By definition $W_{S,C}$ is an irreducible closed subset of $\text{Hilb}^{sc} V$. By Lemma 4.11, $W_{S,C}$ is maximal among all such subsets.

(2) Let C' be a general member of $W_{S,C}$. Then C' is contained in a smooth surface $S' \sim S$ in V . Since C' is general, so is S' in $|S|$. Suppose that $H^1(\mathcal{I}_C(S)) = 0$. Then since (C', S') is a generalization of (C, S) , we have $H^1(\mathcal{I}_{C'}(S')) = H^1(\mathcal{I}_C(S)) = 0$ by the upper semicontinuity. Hence $\text{Hilb}^{sc} V$ is nonsingular at $[C']$ and hence generically smooth along $W_{S,C}$. Suppose that $H^1(\mathcal{I}_C(S)) \neq 0$, i.e., C is not S -normal. Then Lemma 4.8 shows that $H^1(N_{S/V}|_C) \neq 0$ and hence $g \geq 2$. By Proposition 4.9 (2), there exists a line ℓ on S such that $C \cap \ell = \emptyset$. Since $H^1(\mathcal{O}_S) = 0$, the Picard group of S does not change under the smooth deformation of S and hence $\text{Pic } S \simeq \text{Pic } S'$. Since $H^1(\mathcal{O}_S(\ell)) = 0$, the line ℓ is deformed to a line ℓ' on S' . Then we have $C' \cap \ell' = \emptyset$. Moreover since ℓ is a good line, so is ℓ' . Hence $\text{Hilb}^{sc} V$ is singular at $[C']$ by Proposition 4.13. Since C' is a general member of $W_{S,C}$, $\text{Hilb}^{sc} V$ is everywhere singular along $W_{S,C}$ and hence generically non-reduced along $W_{S,C}$. □

5. Original motivation and examples

5.1. Kleppe’s conjecture

The original motivation of the present work was to show the following conjecture due to Kleppe. We denote by $\text{Hilb}_{d,g}^{sc} \mathbb{P}^3$ the open and closed subscheme of $\text{Hilb}^{sc} \mathbb{P}^3$ consisting of curves of degree d and genus g .

CONJECTURE 5.1 (Kleppe, Ellia). — *Let W be a maximal irreducible closed subset of $\text{Hilb}_{d,g}^{sc} \mathbb{P}^3$ whose general member C is contained in a smooth cubic surface. If*

$d \geq 14, \quad g \geq 3d - 18, \quad H^1(\mathbb{P}^3, \mathcal{I}_C(3)) \neq 0 \quad \text{and} \quad H^1(\mathbb{P}^3, \mathcal{I}_C(1)) = 0,$

then W is a component of $(\text{Hilb}^{sc} \mathbb{P}^3)_{\text{red}}$ and $\text{Hilb}^{sc} \mathbb{P}^3$ is generically non-reduced along W .

In the original conjecture [8, Conjecture 4] of Kleppe, the assumption of the linear normality of C (i.e., $H^1(\mathbb{P}^3, \mathcal{I}_C(1)) = 0$) was missing. However Ellia [2] pointed out that the conjecture does not hold for linearly non-normal curves C by a counterexample, and suggested restricting the conjecture to linearly normal ones. The most crucial part to prove this conjecture is the proof of the maximality of W in $(\text{Hilb}^{sc} \mathbb{P}^3)_{\text{red}}$. Once we prove that W is a component of $(\text{Hilb}^{sc} \mathbb{P}^3)_{\text{red}}$, then the non-reducedness of $\text{Hilb}^{sc} \mathbb{P}^3$ along W naturally follows. Therefore Conjecture 5.1 follows from Conjecture 1.2 (1), where the condition $\chi(\mathbb{P}^3, \mathcal{I}_C(3)) \geq 1$ is equivalent to $g \geq 3d - 18$. Recently it has been proved in [15] that Conjecture 5.1 is true if $h^1(\mathbb{P}^3, \mathcal{I}_C(3)) = 1$. Kleppe and Ellia gave a proof for the conjecture under some other conditions, however the whole conjecture is still open.

5.2. Hilbert scheme of canonical curves

Let (V, H) be a polarized variety. We say that a curve $C \subset V$ is *canonical* if $f^*H = K_C$, where $f : C \hookrightarrow V$ is the embedding, or equivalently C is embedded into V by a linear subsystem of $|K_C|$. We apply Theorem 4.14 to prove the following:

PROPOSITION 5.2 (cf. [13]). — *Let V be a smooth del Pezzo 3-fold of degree n . If $n \leq 7$ then the Hilbert scheme $\text{Hilb}^{sc} V$ of smooth connected curves on V has a generically non-reduced component W , whose general member is a canonical curve on V .*

Proof. — Since $n \leq 7$, there exists a good line ℓ on V by Lemma 2.7. Let S_n be a smooth member of $|H|$ containing ℓ . We consider the complete linear system $\Lambda := |-2K_{S_n} + 2\ell|$ on S_n . Let S_{n+1} be the blow-down of ℓ from S_n , which is a del Pezzo surface of degree $n + 1$. Then Λ is the pull-back of $|-2K_{S_{n+1}}| \simeq \mathbb{P}^{3n+3}$ on S_{n+1} . Since Λ is base point free, every general member C of Λ is a smooth connected curve of degree $d = 2n + 2$ and genus $g = n + 2$. Therefore we have $g = d - n$ and hence $\chi(V, \mathcal{I}_C(S)) = 1$. Then ℓ does not intersect C by $(-2K_{S_n} + 2\ell) \cdot \ell = 2 - 2 = 0$. Moreover ℓ is the only such line on S_n . By Theorem 4.14 (1), $W_{S_n, C}$ is an irreducible component of $(\text{Hilb}^{sc} V)_{\text{red}}$. Since $C \cap \ell = \emptyset$, C is not S_n -normal by Proposition 4.9 (2). Therefore $\text{Hilb}^{sc} V$ is generically non-reduced along $W_{S_n, C}$ by Theorem 4.14 (2). By construction, C is the image of a canonical curve $C' \sim -2K_{S_{n+1}}$ on S_{n+1} by the projection $S_{n+1} \cdots \rightarrow S_n$ from a point $p \in S_{n+1} \setminus C'$. \square

Remark 5.3. — The dimension of the irreducible component $W_{S_n, C}$ is equal to $d + g + n = 4n + 4$ by Lemma 4.6 (2). The tangential dimension of $\text{Hilb}^{sc} V$ at a general point $[C]$ of $W_{S_n, C}$ is equal to $h^0(N_{C/V}) = 4n + 5$. Indeed the exact sequence (4.4) is

$$0 \longrightarrow \mathcal{O}_C(2K_C) \longrightarrow N_{C/V} \longrightarrow \mathcal{O}_C(K_C) \longrightarrow 0,$$

since $N_{S/V}|_C \simeq -K_S|_C \simeq K_C$. Hence we have

$$h^0(N_{C/V}) = h^0(2K_C) + h^0(K_C) = (3n + 3) + (n + 2) = 4n + 5.$$

The next example shows that the curve C in Theorem 1.3 is not necessarily stably degenerate if there exists a bad line ℓ on S such that $C \cap \ell = \emptyset$.

Let $V_7 \subset \mathbb{P}^8$ be a smooth del Pezzo 3-fold of degree 7, S_7 a smooth hyperplane section of V_7 . Let ℓ_0, ℓ_1, ℓ_2 be the three lines on S_7 explained in Lemma 2.8, i.e., ℓ_0 is bad and ℓ_1 and ℓ_2 are good. Consider a general member C of $\Lambda := |-2K_{S_7} + 2\ell_0|$. Then C is a smooth connected curve of degree 16 and genus $9 = 16 - 7$ and not S_7 -normal by $C \cap \ell_0 = \emptyset$.

PROPOSITION 5.4. — *Let C be as above. Then there exists a smooth deformation $C' \subset V_7$ of C not contained in any hyperplane section. In other words, C is not stably degenerate.*

Proof. — Recall that V_7 is isomorphic to the blow-up of \mathbb{P}^3 at a point p . It is realized as the projection of the Veronese image $V_8 \subset \mathbb{P}^9$ of \mathbb{P}^3 from $p \in V_8$ (cf. §2.2). Then S_7 is the image by the projection of a hyperplane section $Q_2 \simeq \mathbb{P}^1 \times \mathbb{P}^1$ of V_8 containing p . Hence we have a diagram

$$(5.1) \quad \begin{array}{ccccc} S_7 \simeq \text{Bl}_{2\text{pts}} \mathbb{P}^2 & \subset & V_7 \simeq \text{Bl}_p \mathbb{P}^3 & \subset & \mathbb{P}^8 \\ \downarrow \uparrow & & \pi_p \downarrow \uparrow \Pi_p & & \uparrow \\ Q_2 \simeq \mathbb{P}^1 \times \mathbb{P}^1 & \subset & V_8 \simeq \mathbb{P}^3 & \subset & \mathbb{P}^9, \end{array}$$

where the down arrows (resp. the up arrows) are the blow-up morphisms at (resp. the projections from) $p \in Q_2 \subset V_8 \subset \mathbb{P}^9$. Let $P \simeq \mathbb{P}^2$ denote the exceptional divisor of π_p . Then its intersection with S_7 is equal to the bad line ℓ_0 .

Since $C \cap \ell_0 = \emptyset$ and $C \cdot \ell_i = 4$ for each $i = 1, 2$, π_p maps C isomorphically onto a curve of bidegree $(4, 4)$ on Q_2 . Let Q'_2 be a general hyperplane section of V_8 . Then $Q'_2 \simeq \mathbb{P}^1 \times \mathbb{P}^1$ is mapped isomorphically onto a surface Q''_2 on V_7 by Π_p . Here Q''_2 is linearly equivalent to $S_7 + P$ as a divisor of V_7 and contains a smooth deformation C' of C . Then there exists no hyperplane section of V_7 containing C' . Suppose that there exists such a hyperplane section S'_7 . Then the image $\pi_p(C')$ is contained in the intersection of two hyperplane sections $\pi_p(S'_7)$ and Q'_2 of V_8 . Hence the pull-back of $\pi_p(C')$

in \mathbb{P}^3 by the Veronese embedding is contained in a complete intersection of two quadrics. This is impossible since the degree of the inverse image is equal to $8 > 4$. □

5.3. Non-reduced components of the Hilbert scheme

In this subsection, we give the classes of irreducible components of the Hilbert scheme of del Pezzo 3-folds which are non-reduced by Theorem 4.14 more precisely (cf. Proposition 5.5).

Let V_n be a smooth del Pezzo 3-fold of degree $n \leq 7$ and let $S \subset V_n$ be a smooth member of the class $|H|$ of the polarization of V_n , i.e., a del Pezzo surface. Put $r := 9 - n (\geq 2)$. Then S is isomorphic to a \mathbb{P}^2 blown up at r points in general position, i.e., no three are on a line, no six are on a conic and any cubic containing eight points is smooth at each of them. The class of the pullback l of a line in \mathbb{P}^2 and the r exceptional curves e_i ($1 \leq i \leq r$) form a free \mathbb{Z} -basis of the Picard group $\text{Pic } S \simeq \mathbb{Z}^{r+1}$ of S . Thus given a divisor D on S , we obtain a $(r + 1)$ -tuple $(a; b_1, \dots, b_r)$ of integers as the coefficients of the divisor class $D = al - \sum_{i=1}^r b_i e_i$. On the other hand, for each $r \geq 2$ there exists a Weyl group $W_r \subset \text{Aut}(\text{Pic } S)$. Here W_r is the subgroup generated by the permutations of e_i ($1 \leq i \leq r$) and (for $r \geq 3$) by the additional Cremona element σ given by $\sigma(l) = 2l - e_1 - e_2 - e_3$, $\sigma(e_1) = l - e_2 - e_3$, $\sigma(e_2) = l - e_1 - e_3$, $\sigma(e_3) = l - e_1 - e_2$ and $\sigma(e_i) = e_i$ for $i \notin \{1, 2, 3\}$. The root systems corresponding to the Weyl group W_r ($r = 2, 3, 4, 5, 6, 7, 8$) are $A_1, A_1 \times A_2, A_4, D_5, E_6, E_7, E_8$, respectively (See [12] for the details). Every element of W_r induces a base change of $\text{Pic } S$. By virtue of the Weyl groups and this base change, given a divisor D on S there exists a suitable blow-up $S \rightarrow \mathbb{P}^2$ (in other words, a suitable choice of r exceptional curves on S) such that we have

$$(5.2) \quad b_1 \geq \dots \geq b_r \quad \text{and} \quad a \geq b_1 + b_2 + b_3 \quad (\text{only for } r \geq 3).$$

When (5.2) holds, we say the basis $\{l, e_1, \dots, e_r\}$ of $\text{Pic } S$ is *standard* for D . For the standard basis of $\text{Pic } S$ for D , the linear system $|D|$ on S contains a smooth connected curve C of degree > 2 if and only if $a > b_1$ and $b_r \geq 0$ (and $a \geq b_1 + b_2$ for $r = 2$). The degree d and genus g of C is computed as

$$(5.3) \quad d = 3a - \sum_{i=1}^r b_i \quad \text{and} \quad g = \binom{a-1}{2} - \sum_{i=1}^r \binom{b_i}{2}.$$

Let (d, g) be a pair of integers with $d > 2$ and let $(a; b_1, \dots, b_r)$ be a $(r + 1)$ -tuple of integers satisfying (5.2), (5.3), $a > b_r$ and $b_r \geq 0$ (and

$a \geq b_1 + b_2$ as well for $r = 2$). Then the linear system $|al - \sum_{i=1}^r b_i e_i|$ on S contains a smooth connected member C of degree d and genus g . Then we denote by $W_{(a;b_1, \dots, b_r)}$ the S -maximal family $W_{S,C} \subset \text{Hilb}_{d,g}^{sc} V_n$ of curves containing C (cf. Definition 4.4). By definition, $W_{(a;b_1, \dots, b_r)}$ contains every smooth connected curve C' on V_n such that C' is contained in a smooth member $S' \in |H|$ and such that $C' \sim al' - \sum_{i=1}^r b_i e'_i$ on S' for a standard basis $\{l', e'_1, \dots, e'_r\}$ of $\text{Pic } S'$ for C' .

PROPOSITION 5.5. — *Suppose that $g \geq 2$ and $g \geq d - n$. If $b_r = 0$, then $W_{(a;b_1, \dots, b_r)}$ is an irreducible component of $(\text{Hilb}_{d,g}^{sc} V_n)_{\text{red}}$ of dimension $d + g + n$ and $\text{Hilb}_{d,g}^{sc} V_n$ is generically non-reduced along $W_{(a;b_1, \dots, b_r)}$.*

Proof. — Let C denote a general member of $W_{(a;b_1, \dots, b_r)}$. Then C is contained in a smooth member $S \in |H|$. Since C is general, so is S in $|H|$. By Lemma 2.8 every line on S is good except for the bad line ℓ_0 on V_7 . If $n = 7$ then ℓ_0 is linearly equivalent to $l - e_1 - e_2$. Since $b_2 = 0$, C intersects ℓ_0 by $C \cdot \ell_0 = (al - b_1 e_1) \cdot \ell_0 = a - b_1 > 0$. We recall that $g \geq d - n$ is equivalent to $\chi(V, \mathcal{I}_C(S)) \geq 1$. Therefore $W_{(a;b_1, \dots, b_r)}$ is an irreducible component of $(\text{Hilb}_{d,g}^{sc} V_n)_{\text{red}}$ by Theorem 4.14 (1), and of dimension $d + g + n$ by Lemma 4.6.

Since $b_r = 0$, the line e_r on S does not intersect C . Since $g \geq 2$, C is not S -normal by Proposition 4.9 (2), and hence we have $H^1(V, \mathcal{I}_C(S)) \neq 0$. Thus $\text{Hilb}_{d,g}^{sc} V_n$ is generically non-reduced along $W_{(a;b_1, \dots, b_r)}$ by Theorem 4.14 (2). □

The next example shows that for every integer $d \geq 12$ the Hilbert scheme of smooth connected curves of degree d on a smooth cubic 3-fold V_3 has a generically non-reduced component.

Example 5.6. — Let $\lambda \in \mathbb{Z}_{\geq 0}$ and let W be one of the S -maximal families

$$\begin{aligned} W_{(\lambda+6; \lambda+1, 1, 1, 1, 1, 0)} &\subset \text{Hilb}_{d, 2d-16}^{sc} V_3 \quad (d = 2\lambda + 13) \quad \text{and} \\ W_{(\lambda+6; \lambda+2, 1, 1, 1, 1, 0)} &\subset \text{Hilb}_{d, \frac{3}{2}d-9}^{sc} V_3 \quad (d = 2\lambda + 12). \end{aligned}$$

Then W is an irreducible component of $(\text{Hilb}^{sc} V_3)_{\text{red}}$ and $\text{Hilb}^{sc} V_3$ is generically non-reduced along W .

It was shown in [13, Theorem 1.4] that for many uniruled 3-folds V the Hilbert scheme $\text{Hilb}^{sc} V$ has infinitely many generically non-reduced components.

BIBLIOGRAPHY

- [1] D. CURTIN, “Obstructions to deforming a space curve”, *Trans. Amer. Math. Soc.* **267** (1981), p. 83-94.
- [2] P. ELLIA, “D’autres composantes non réduites de Hilb \mathbb{P}^3 ”, *Math. Ann.* **277** (1987), p. 433-446.
- [3] G. FLØYSTAD, “Determining obstructions for space curves, with applications to non-reduced components of the Hilbert scheme”, *J. Reine Angew. Math.* **439** (1993), p. 11-44.
- [4] T. FUJITA, “On the structure of polarized manifolds with total deficiency one. I”, *J. Math. Soc. Japan* **32** (1980), p. 709-725.
- [5] ———, “On the structure of polarized manifolds with total deficiency one. II”, *J. Math. Soc. Japan* **33** (1981), p. 415-434.
- [6] V. A. ISKOVSKIĖ, “Fano threefolds. I”, *Math. USSR-Izvstija* **11** (1977), no. 3, p. 485-527.
- [7] ———, “Anticanonical models of three-dimensional algebraic varieties, Current problems in mathematics”, *J. Soviet Math.* **13** (1980), p. 745-814.
- [8] J. O. KLEPPE, “Non-reduced components of the Hilbert scheme of smooth space curves.”, *Space curves, Proc. Conf., Rocca di Papa/Italy 1985, Lect. Notes in Math.* 1266, 181-207, 1987.
- [9] ———, “Liaison of families of subschemes in \mathbf{P}^n ”, in *Algebraic curves and projective geometry (Trento, 1988)*, Lecture Notes in Math., vol. 1389, Springer, Berlin, 1989, p. 128-173.
- [10] ———, “The Hilbert scheme of space curves of small diameter”, *Annales de l’institut Fourier* **56** (2006), no. 5, p. 1297-1335.
- [11] J. KOLLÁR, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 32, Springer-Verlag, Berlin, 1996, viii+320 pages.
- [12] Y. I. MANIN, *Cubic forms*, second ed., North-Holland Mathematical Library, vol. 4, North-Holland Publishing Co., Amsterdam, 1986, Algebra, geometry, arithmetic, Translated from the Russian by M. Hazewinkel, x+326 pages.
- [13] S. MUKAI & H. NASU, “Obstructions to deforming curves on a 3-fold. I. A generalization of Mumford’s example and an application to Hom schemes”, *J. Algebraic Geom.* **18** (2009), no. 4, p. 691-709.
- [14] D. MUMFORD, “Further pathologies in algebraic geometry”, *Amer. J. Math.* **84** (1962), p. 642-648.
- [15] H. NASU, “Obstructions to deforming space curves and non-reduced components of the Hilbert scheme”, *Publ. Res. Inst. Math. Sci.* **42** (2006), no. 1, p. 117-141.

Manuscrit reçu le 30 mars 2009,
 accepté le 3 juillet 2009.

Hirokazu NASU
 Kyoto University
 Research Institute for Mathematical Sciences
 Kyoto 606-8502 (Japan)
 nasu@sie.dendai.ac.jp