

# ANNALES DE L'I. H. P., SECTION B

DAVID J. GRABINER

## **Brownian motion in a Weyl chamber, non-colliding particles, and random matrices**

*Annales de l'I. H. P., section B*, tome 35, n° 2 (1999), p. 177-204

<[http://www.numdam.org/item?id=AIHPB\\_1999\\_\\_35\\_2\\_177\\_0](http://www.numdam.org/item?id=AIHPB_1999__35_2_177_0)>

© Gauthier-Villars, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section B » (<http://www.elsevier.com/locate/anihpb>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>*

# Brownian motion in a Weyl chamber, non-colliding particles, and random matrices

by

**David J. GRABINER**

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-1109  
e-mail: grabiner@math.lsa.umich.edu

---

**ABSTRACT.** – Let  $n$  particles move in standard Brownian motion in one dimension, with the process terminating if two particles collide. This is a specific case of Brownian motion constrained to stay inside a Weyl chamber; the Weyl group for this chamber is  $A_{n-1}$ , the symmetric group. For any starting positions, we compute a determinant formula for the density function for the particles to be at specified positions at time  $t$  without having collided by time  $t$ . We show that the probability that there will be no collision up to time  $t$  is asymptotic to a constant multiple of  $t^{-n(n-1)/4}$  as  $t$  goes to infinity, and compute the constant as a polynomial of the starting positions. We have analogous results for the other classical Weyl groups; for example, the hyperoctahedral group  $B_n$  gives a model of  $n$  independent particles with a wall at  $x = 0$ .

We can define Brownian motion on a semisimple Lie algebra, viewing it as a vector space with the Killing form. Since the Killing form is invariant under the adjoint, the motion induces a process in the Weyl chamber of the Lie algebra, giving a Brownian motion conditioned never to exit the chamber. If there are  $m$  roots in  $n$  dimensions, this shows that the radial part of the conditioned process is the same as the  $n + 2m$ -dimensional Bessel process. The conditioned process also gives physical models, generalizing Dyson's model for  $A_{n-1}$  corresponding to  $u_n$  of  $n$  particles moving in a

---

Supported by an NSF Graduate Fellowship and NSF Postdoctoral Fellowship, and a postdoctoral fellowship from the Center for Discrete Mathematics and Theoretical Computer Science at the Hebrew University of Jerusalem. Some of this work was done while visiting the Mathematical Sciences Research Institute; MSRI is supported by NSF Grant DMS-9022140.

diffusion with a repelling force between two particles proportional to the inverse of the distance between them. © Elsevier, Paris

**RÉSUMÉ.** — Soient  $n$  particules se déplaçant selon des mouvements browniens en dimension 1, le processus étant tué si deux particules se rencontrent. C'est un cas particulier du mouvement brownien contraint à rester dans une chambre de Weyl; le groupe de Weyl pour cette chambre est  $A_{n-1}$ , le groupe symétrique. Pour toutes les positions initiales, nous calculons une formule de déterminant qui donne la fonction de densité pour les positions des particules à l'instant  $t$  lorsqu'elles ne se sont pas rencontrées avant  $t$ . Nous démontrons que la probabilité qu'il n'y ait pas de collision avant le temps  $t$  se comporte comme  $t^{n(n-1)/4}$  quand  $t \rightarrow \infty$ , et nous calculons la constante intervenant en fonction des positions initiales. Nous donnons des résultats analogues pour les autres groupes de Weyl classiques; par exemple, le groupe  $B_n$  donne un modèle de  $n$  particules indépendantes avec un mur en  $x = 0$ .

On peut définir le mouvement brownien sur une algèbre de Lie semisimple en considérant l'algèbre comme un espace vectoriel avec la forme de killing. Puisque la forme de killing est invariante par l'adjoint, le mouvement induit un processus dans la chambre de Weyl. Ceci donne un mouvement brownien conditionné à ne jamais sortir de la chambre. S'il y a  $m$  racines en  $n$  dimensions, la partie radiale du processus conditionné est identique au processus de Bessel en  $(n + 2m)$  dimensions. Le processus donne aussi des modèles physiques qui généralisent le modèle de Dyson pour  $A_{n-1}$  correspondant à  $u_n$ , avec  $n$  particules en diffusion et une force répulsive pour chaque paire de particules proportionnelle à l'inverse de la distance entre ces particules. © Elsevier, Paris

## 1. INTRODUCTION

Let  $n$  particles move in standard Brownian motion in one dimension, with the process terminating if two particles collide. Given the starting positions, we can use a reflection argument to calculate the density function for the particles to be at specified positions at time  $t$  without having collided by time  $t$ . Using this density function and the theory of Lie algebras, we can prove the following results. Theorems 3 and 4 were proved by Dyson [10] for the Brownian motion model, and by Biane [4] for the conditioned process.

**THEOREM 1.** – *For any starting positions, the probability that there will be no collision up to time  $t$  is asymptotic to a constant multiple of  $t^{-n(n-1)/4}$  as  $t$  goes to infinity; the constant is a known polynomial in the starting positions.*

**THEOREM 2.** – *Given that there is no collision up to time  $t$ , the distribution of the radius of the vector whose coordinates are the positions of the particles, divided by the square root of  $t$ , converges in measure to the distribution of the Bessel process with parameter  $n(n+1)/2$  at time 1, which is the radial part of an  $n(n+1)/2$ -dimensional standard normal.*

**THEOREM 3.** – *We can construct an  $n$ -dimensional Brownian motion which is conditioned for no two particles ever to collide. If we take Brownian motion on the space of Hermitian matrices, the induced process on the eigenvalues is the same process. If the starting point is appropriately chosen at the starting radius, the process given by the radial part of the conditioned Brownian motion is the same process as the  $n^2$ -dimensional Bessel process.*

**THEOREM 4.** – *This conditioned process is identical to the process obtained by  $n$  particles moving in a one-dimensional diffusion with constant infinitesimal variance, with a repelling force between two particles proportional to the inverse of the distance between them; that is, its infinitesimal generator has  $\sigma_{ij}^2(\vec{x}, t) = \delta_{ij}$ ,  $\mu_i(\vec{x}, t) = \sum_{i \neq j} 1/(x_i - x_j)$ .*

This is a specific case of Brownian motion in a Weyl chamber; the vector whose coordinates are the locations of the  $n$  particles is constrained to stay inside the chamber. The Weyl group is  $A_{n-1}$ , the symmetric group.

We have similar results for other Weyl groups; Theorems 1 and 2 have analogues for the classical Weyl groups, and Theorems 3 and 4 have analogues for all Weyl groups. For example, the Weyl group  $B_n$  models  $n$  particles in independent Brownian motion with an absorbing wall at  $x = 0$ . In this case, the asymptotic probability is  $ct^{-n^2/2}$  (and the constant is again known), with the radial part corresponding to the  $n(n+1)$ -dimensional Bessel process; the conditioned process has a radial part which is the  $n(2n+1)$ -dimensional Bessel process. This generalizes the results of Pitman and Williams [24, 28] that one-dimensional Brownian motion conditioned to stay positive is the same process as the three-dimensional Bessel process.

Our reflection argument is a generalization of the *reflection principle*, a standard argument in the analysis of both discrete random walks and Brownian motion. In the discrete case, it is used in the classical formula for the Catalan numbers, which enumerate the arrangements of  $n+1$ 's and

$n - 1$ 's so that none of the partial sums are negative. Similarly, it can be used to study Brownian motion in one dimension with an absorbing barrier at  $x = 0$  and a known starting point [14]. It has also been extended to multiple reflections in one dimension to study Brownian motion with two absorbing barriers [11].

The reflection principle has been generalized to multiple dimensions. For example, the *ballot problem*, a classical problem in random walks, asks how many ways there are to walk from the origin to a point  $(\lambda_1, \dots, \lambda_n)$ , taking  $k$  unit-length steps in the positive coordinate directions while staying in the region  $x_1 \geq x_2 \geq \dots \geq x_n$ . The solution is known in terms of the hook-length formula for Young tableaux; a combinatorial proof, using a reflection argument, is given in [27, 29].

The same reflection argument has also been applied to the case of  $n$  independent diffusions, or discrete processes which cannot pass each other without first colliding. Using this method, Karlin and McGregor [17] give a determinant formula for the probability or measure for the  $n$  particles, starting at known positions, not to have collided up to time  $t$  and to be in given positions. Hobson and Werner [16] generalize this argument to  $n$  particles in an interval or circle, and use this to prove a result analogous to Theorem 4 for  $n$  particles on the circle.

Gessel and Zeilberger [12], and independently Biane [3], give a further generalization. For certain “reflectable” random walk-types, we can count the number of  $k$ -step walks between two points of a lattice, staying within a chamber of a Weyl group, in terms of numbers of unconstrained walks. The steps must have certain allowable lengths and directions. In [13], all cases in which this method applies are enumerated, and determinant formulas are given for many important cases, including walks in the classical Weyl chambers.

The argument of [12] can be generalized to Brownian motion in any Weyl chamber or chamber of a Coxeter group, with either absorbing or reflecting boundary conditions. We prove this generalization, and then use the result to compute determinant formulas for Brownian motion in the Weyl chambers of  $A_{n-1}$ ,  $B_n = C_n$ , and  $D_n$ . The  $A_{n-1}$  and  $B_n$  cases are applicable to the independent motion of  $n$  particles in one dimension. The  $A_{n-1}$  formula appears in [17], using the model of  $n$  independent particles rather than motion in a Weyl chamber. The cases of the affine Weyl groups  $\tilde{A}_{n-1}$  and  $\tilde{B}_n$  are studied in [16], also as models of  $n$  independent particles, in a circle or interval.

These determinants factor into forms which can be easily analyzed; this allows us to find the asymptotic probability that there will be no collision

up to time  $t$  for these three cases, for any starting point. In the case of  $A_{n-1}$ , we get a simple formula for the actual measure as well.

Weyl chambers arise in Lie theory as the set of orbits of the adjoint action on a Lie algebra, or conjugation under the associated Lie group. We can define a Brownian motion on the Lie algebra which is invariant under the adjoint by using the Killing form to obtain a norm; for  $A_{n-1}$ , this norm is the square root of the sums of the squares of the absolute values of all the matrix entries. There is thus a natural correspondence between standard Brownian motion on the Lie algebra and some diffusion on the Weyl chamber. Dyson [10] shows that this diffusion has the same generator as the physical model of Theorem 4 for  $A_{n-1}$ . Biane [4] describes the correspondence with the conditioned motion, proving Theorem 3 for  $A_{n-1}$ . These results generalize naturally to other Lie algebras. We can thus use the known properties of random matrices to study the distribution.

DeBlassie [8] uses a different approach to give a more general formula for asymptotics and density functions for a general class of cones, which include the cases discussed here. A cone is defined as the union of all rays from the origin which intersect the unit sphere in a connected open set  $C$ . The Laplace-Beltrami operator  $L_{S^{n-1}}$  on the unit sphere is the non-radial portion of the ordinary Laplacian. If  $\lambda_1$  is the eigenvalue of  $L_{S^{n-1}}$  with smallest absolute value on the space of all  $L^2$  functions on  $C$  which vanish continuously on the boundary of  $C$ , then the asymptotic probability is a constant multiple of

$$t^{-\left[-(n-2)+\sqrt{(n-2)^2+4\lambda_1^2}\right]/4}.$$

This result gives asymptotics for a large class of cones. The coefficients depend on the eigenvalues of the Laplace-Beltrami operator rather than on explicit coefficients; these eigenvalues are known for Weyl chambers [2]. The asymptotics can thus be computed from these formulas; in theory, the explicit density functions can also be computed, but as infinite series. This formula also shows that the asymptotics for all such cones exist and are powers of  $t$ , with no other terms such as logarithms.

The asymptotic probability is known to be a constant multiple of  $t^{-m/2}$  for a wedge of angle  $\pi/m$  in two dimensions [7]. The result holds in general, although the region is only a chamber of a Coxeter group (the dihedral group) if  $m$  is an integer.

O'Connell and Unwin [23] compute an explicit formula for the collision probability of three independent particles, the case  $n = 3$  of Theorem 1. They also study the opposite asymptotic problem to ours, computing an

asymptotic for the probability that  $n$  particles in independent Brownian motion will have a collision up to time  $t$  when  $t$  is small compared to the initial separation of the particles.

Our results are organized as follows. Section 2 contains the basic definitions. In section 3, we prove the basic reflection result, and in section 4, we apply this result to get the determinant formulas. In section 5, we prove Theorems 1 and 2, and their analogues for  $B_n$  and  $D_n$ . In section 6, we construct the conditioned motion, both by Lie theory and by  $h$ -transformation, proving Theorems 3 and 4 and their analogues, and use the  $h$ -transformation to find a physical model for all finite Coxeter groups.

## 2. DEFINITIONS

We will study a process with continuous sample paths in  $\mathbb{R}^n$ , either unconstrained or constrained by a chamber. In the constrained case, we may have either an absorbing boundary condition, causing the process to terminate when it hits a wall, or a reflecting boundary condition. All references to the analogous discrete problem are discussed in [13].

We require that our chamber  $C$  be a chamber of a finite or affine *Coxeter group*. In the finite case,  $C$  is defined by a system of simple roots  $\Delta \subset \mathbb{R}^n$  as

$$(1) \quad C = \{\vec{x} \in \mathbb{R}^n \mid (\alpha, \vec{x}) \geq 0 \text{ for all } \alpha \in \Delta\},$$

and the orthogonal reflections  $r_\alpha: \vec{x} \mapsto \vec{x} - \frac{2(\alpha, \vec{x})}{(\alpha, \alpha)}\alpha$ , generate a finite group  $W$  of linear transformations, the *Coxeter group*. In the affine case, the hyperplanes of reflection which define  $C$  do not all pass through the origin, and the group  $W$  is infinite, but if  $T$  is the subgroup of all translations,  $W/T$  must be finite. In the analogous discrete problem, the steps of the random walk must generate a lattice  $L$  which is stable under the action of  $W$ ; in this case,  $C$  is a *Weyl chamber* and  $W$  a *Weyl group*.

Let  $X(t)$  be a Markov process with continuous sample paths with values in  $\mathbb{R}^n$ ; that is, the distribution of  $X(t_2)$  given  $X(t_1)$  is independent of  $X(t)$  at any  $t$  outside the interval  $[t_1, t_2]$ . We say that the constrained motion is *reflectable* if the increments of the unconstrained motion are symmetric under the Coxeter group; that is, the distribution of  $X(t_2)$  given  $X(t_1)$  is the same as the distribution of  $w(X(t_2))$  given the  $w$ -image  $w(X(t_1))$  as the starting point for any  $w \in W$ .

Standard Brownian motion is reflectable for any Coxeter group. For a finite Coxeter group, in which all planes of reflection pass through the

origin, any diffusion with variance dependent only on time and the radius, and drift dependent on time and symmetric with respect to rotations and reflections about the origin, is reflectable; for example, there could be an absorbing or reflecting barrier at  $|\vec{x}| = R$ .

As another example, consider the case in which each coordinate  $x_i(t)$  is an independent identical diffusion; this could model  $n$  independent particles instead of one particle in  $n$  dimensions. If our Coxeter group is the symmetric group  $A_{n-1} = S_n$  (giving the chamber  $x_1 > x_2 > \dots > x_n$ ), it permutes the particles, so the process is reflectable under this action; this case is discussed in [17]. If the individual diffusions are symmetric about  $x_i = 0$ , then the product process will also be reflectable under the hyperoctahedral group  $B_n$ , which includes all permutations with any number of sign changes.

In the discrete case, reflectability requires the additional condition that the walk cannot go from inside the chamber to outside it without stopping on a wall [12]; this is our condition of continuous sample paths, which is satisfied by any diffusion.

For fixed  $t$ , this process defines a probability measure  $P_t(A) = P\{X(t) \in A\}$  which represents the chance that this process, if started at 0, will be in a set  $A$  at time  $t$ . We assume that this probability measure has a density function  $c_t(\vec{x})$  with respect to Lebesgue measure on  $\mathbb{R}^n$ ; that is,

$$\int_{\vec{x} \in A} c_t(\vec{x}) = P_t(A).$$

Now, we study the case in which the motion is constrained by a chamber, with either absorbing or reflecting boundary conditions. We must now fix the starting point  $\eta$ , since the process now depends on it in a non-trivial way. This generates new stochastic processes  $Y(\eta, t)$  for absorbing boundary conditions and  $Y'(\eta, t)$  for reflecting boundary conditions. These give probability measures  $Q_t(\eta, A)$  and  $Q'_t(\eta, A)$  which give the probability that the process, if started at  $\eta$ , will be in the set  $A$  at time  $t$ , and these have density functions  $b_t(\eta, \lambda)$  and  $b'_t(\eta, \lambda)$ . Note that the total measure  $Q_t(\eta, \mathbb{R}^n)$  will be less than 1 because the process terminates when it reaches a wall of the chamber. The total measure  $Q'_t(\eta, \mathbb{R}^n)$  for reflecting boundary conditions will still be 1.

If our process is a Brownian motion, it has drift  $\mu_i$  and variance  $\sigma_i^2$  in each coordinate direction. If the Coxeter group contains any reflection in the  $x_i, x_j$ -plane other than a sign change of  $x_i$  or  $x_j$ , the reflectability condition requires that  $\sigma_i = \sigma_j$ . Thus, for an irreducible Coxeter group, all of the  $\sigma_i$  must be equal; for a reducible Coxeter group acting on  $\mathbb{R}^{n_1} \oplus \mathbb{R}^{n_2} \oplus \dots \oplus \mathbb{R}^{n_k}$ ,

we can multiply the coordinates in each  $\mathbb{R}^{n_i}$  by a constant factor so that the  $\sigma_i$  are all equal. We can then re-scale time so that all  $\sigma_i = 1$ .

Reflectability also requires the Coxeter group to fix the vector  $\mu$  whose coordinates are the drifts  $\mu_i$ . In all non-trivial cases except for  $A_{n-1}$  on  $\mathbb{R}^n$ , this requires that all  $\mu_i = 0$ , giving standard Brownian motion. For  $A_{n-1}$ , it requires that the  $\mu_i$  all be equal; we can then change coordinates to  $x'_i = x_i - \mu_i t$  to get an equivalent process in which all of the  $\mu'_i$  are zero. Thus, if the unconstrained process has stationary increments, and is thus a Brownian motion [14], we may assume that it is standard Brownian motion; however, we can state the theorems just as easily in terms of the more general reflectable process.

### 3. THE REFLECTION ARGUMENT

**THEOREM 5.** – *If  $c_t$  is the density function for a reflectable continuous stochastic process, then for absorbing boundary conditions, we have*

$$(2) \quad b_t(\eta, \lambda) = \sum_{w \in W} \operatorname{sgn}(w) c_t(w(\lambda) - \eta),$$

and for reflecting boundary conditions,

$$(3) \quad b'_t(\eta, \lambda) = \frac{\sum_{w \in W} c_t(w(\lambda) - \eta)}{\#\{w \in W : w(\lambda) = \lambda\}}.$$

If  $W$  is an affine group rather than a finite group, these may be infinite sums. The integrals over the images of any region must converge absolutely, since the measure of unconstrained motion over the whole space, the set of all  $W$ -images of all points in the chamber, is 1.

*Proof.* – The discrete result analogous to (2) is proved in [12] and [3]; the proof which follows is essentially identical to that in [12] except that the discrete terms “walk” and “step” are replaced by their continuous analogues “path” and “time.”

Every path from  $\eta$  to any  $w(\lambda)$  which does touch at least one wall of the chamber has some first time  $t_0$  at which it touches a wall; let the wall be the hyperplane perpendicular to  $\alpha_i$ , choosing the largest  $i$  if there are several choices [25]. Reflect the path after time  $t_0$  across that hyperplane; the resulting path is a path from  $\eta$  to  $r_{\alpha_i} w(\lambda)$  which also first touches wall  $i$  at time  $t_0$ . This clearly gives a measure-preserving bijection of paths, and since  $r_{\alpha_i}$  has sign  $-1$ , all such paths cancel out in (2). The only paths

which do not cancel in these pairs are the paths which stay within the Weyl chamber, and since  $w(\lambda)$  is inside the Weyl chamber only if  $w$  is the identity, this is the desired measure.

For (3), we note that the map on all paths starting at  $\eta$  which takes every point to its unique image in the chamber  $C$  is measure-preserving, since we have reflecting boundary conditions and increments which are stable under the group  $W$ . This map takes all paths which end at any  $w(\lambda)$  to paths which end at  $\lambda$  itself. If  $\lambda$  is on a wall of the chamber, paths to  $\lambda$  may be counted multiple times, so we must divide by the size of its stabilizer in  $W$ .

In practice, we can ignore this constant factor; it is 1 except for  $\lambda$  on a wall, and this is a set of measure zero unless  $t = 0$ . Eliminating the denominator thus changes the density function only on a set of measure zero, and thus does not change the measure of any measurable set.

#### 4. DETERMINANT FORMULAS FOR THE DENSITY FUNCTIONS

We can now apply this theorem to standard Brownian motion, in the Weyl chambers of  $A_{n-1}$ ,  $B_n = C_n$ , and  $D_n$ , with either absorbing or reflecting boundary conditions. The measure for unconstrained standard Brownian motion is  $c_t(\vec{x}) = \prod_{i=1}^n N_t(x_i)$ , where  $N_t$  is the normal distribution function with mean 0 and variance  $t$ . Since this factors into separate terms for the individual coordinates, we can use the same techniques to compute determinant formulas as in the discrete case [13].

The most interesting case is  $A_{n-1} = S_n$ , the symmetric group. The Weyl chamber is  $x_1 > x_2 > \dots > x_n$ . This Brownian motion thus models  $n$  independent particles in one dimension. With absorbing boundary conditions, collisions are forbidden (the process terminates if one occurs); with reflecting boundary conditions, particles collide elastically with one another.

For absorbing boundary conditions, we write the sum (2) as

$$(4) \quad b_t(\eta, \lambda) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) c_t(\sigma(\lambda) - \eta)$$

and use the value of  $c_t$  to write this as

$$(5) \quad b_t(\eta, \lambda) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{i=1}^n (N_t(\lambda_{\sigma(i)} - \eta_i)).$$

This sum can be written as a determinant, which gives

$$(6) \quad b_t(\eta, \lambda) = \det_{n \times n} |N_t(\lambda_i - \eta_j)|.$$

This determinant gives the measure for  $n$  particles which start at positions  $\eta_i$  and are in independent Brownian motion to be at positions  $\lambda_i$  at time  $t$  without having collided.

For a more general product of  $n$  independent diffusions, with individual density  $p_t(x \rightarrow y)$  for the diffusion which started at point  $x$  to be at point  $y$  at time  $t$ , we can apply the same argument. This gives the following generalization of (6), which first appears in [17].

$$(7) \quad b_t(\eta, \lambda) = \det_{n \times n} |p_t(\eta_j \rightarrow \lambda_i)|.$$

For reflecting boundary conditions, the calculations are the same except that there is no term for  $\text{sgn}(\sigma)$ , and thus we get permanents rather than determinants. This result can also be seen by observing that an elastic collision between two identical particles is equivalent to the two particles passing through each other with no collision, and thus the reflected particles will be at the positions indicated by  $\lambda$  if the unreflected particles are at any permutation of the coordinates of  $\lambda$ .

For  $B_n$ , the hyperoctahedral group, which includes permutations with any number of sign changes, the Weyl chamber is  $x_1 > x_2 > \dots > x_n > 0$ . This also models  $n$  independent particles in one dimension, with an additional wall at  $x = 0$ .

We write  $w \in W$  as a product of an  $\epsilon$  which negates some coordinates and a  $\sigma$  in the symmetric group. We get

$$(8) \quad b_t(\eta, \lambda)(x) = \sum_{\sigma \in S_n} \sum_{\epsilon_i = \pm 1} \text{sgn}(\sigma) \prod_{i=1}^n \epsilon_i \prod_{i=1}^n (N_t(\epsilon_i \lambda_{\sigma(i)} - \eta_i)).$$

Using the multilinearity of the products in the determinant, we can again write this sum as a determinant, with separate terms for  $\epsilon_i = 1$  and  $\epsilon_i = -1$  in each entry. We use  $N_t(x) = N_t(-x)$  to keep the signs of  $\lambda$  positive and get a more elegant formula. This gives

$$(9) \quad b_t(\eta, \lambda) = \det_{n \times n} |N_t(\lambda_i - \eta_j) - N_t(\lambda_i + \eta_j)|.$$

This determinant gives the measure for  $n$  particles which start at  $\eta_i$  to be at  $\lambda_i$  at time  $t$ , neither having collided nor having touched  $x = 0$ .

Again, the same argument applies if  $n$  particles are in general independent diffusions, provided that the diffusions are symmetric about  $x = 0$ . The more general formula is

$$(10) \quad b_t(\eta, \lambda) = \det_{n \times n} |p_t(\eta_j \rightarrow \lambda_i) - p_t(\eta_j \rightarrow -\lambda_i)|.$$

For reflecting boundary conditions, we lose the sign of the  $\sigma$ , which makes the determinant into a permanent, and the sign of the  $\epsilon_i$ , which turns the minus sign between the two  $N_t$  in (9) or  $p_t$  in (10) into a plus sign. Again, the resulting formula is the same as would be obtained by treating elastic collisions as though the particles passed through each other, and allowing particles to pass through the wall at  $x = 0$  instead of bouncing. (In the transformed model, particles at positions  $x$  and  $-x$  no longer collide, but since they collided elastically in the original model and could be considered to pass through each other instead, the effect is the same.)

For  $D_n$ , the even hyperoctahedral group, which includes permutations with an even number of sign changes, the Weyl chamber is  $x_1 > x_2 > \dots > x_n, x_{n-1} > -x_n$ . This does not give a natural model for  $n$  particles in one dimension.

Again, we can write  $w = \epsilon\sigma$ . We take our sum over all possible  $\epsilon$ , and then add an additional factor of  $(1 + \prod_{i=1}^n \epsilon_i)/2$  to annihilate those  $\epsilon$  which are not allowed in  $D_n$ .

$$(11) \quad b_t(\eta, \lambda)(x) = \sum_{\sigma \in S_n} \sum_{\epsilon_i = \pm 1} \operatorname{sgn}(\sigma) \frac{1 + \prod_{i=1}^n \epsilon_i}{2} \prod_{i=1}^n (\epsilon_i N_t(\epsilon_i \lambda_{\sigma(i)} - \eta_i)).$$

We now take the  $\frac{1}{2}$  and the  $(\prod_{i=1}^n \epsilon_i)/2$  terms separately. The  $(\prod_{i=1}^n \epsilon_i)/2$  term is half the sum we had in (9); the  $\frac{1}{2}$  term gives half of (9), but with a plus sign between the terms. Thus we get

$$(12) \quad b_t(\eta, \lambda)(x) = \frac{1}{2} \left[ \det_{n \times n} |N_t(\lambda_i - \eta_j) - N_t(\lambda_i + \eta_j)| + \det_{n \times n} |N_t(\lambda_i - \eta_j) + N_t(\lambda_i + \eta_j)| \right].$$

If we let  $\lambda'$  be obtained from  $\lambda$  by changing the sign of  $\lambda_n$ , this will change the sign of the first term but preserve the second term. Thus the first term alone, with no factor of  $1/2$ , is  $b_t(\eta, \lambda)(x) - b_t(\eta, \lambda')(x)$ , and the second term alone is  $b_t(\eta, \lambda)(x) + b_t(\eta, \lambda')(x)$ . If  $\lambda_n = 0$  or  $\eta_n = 0$ , the first term is zero, so  $b_t(\eta, \lambda)(x)$  is the second term alone, with the factor of  $1/2$ .

For reflecting boundary conditions, we ignore the sign of  $\sigma$ . Thus the determinants become permanents, but we keep the minus signs because they came from the factor  $(1 + \prod_{i=1}^n \epsilon_i)/2$ , which was not from  $\operatorname{sgn}(w)$ .

## 5. ASYMPTOTICS

We can use these formulas to find asymptotics for the probability that the motion will not hit a wall of the chamber by time  $t$ , and for its distribution at time  $t$  given that it has not hit a wall.

### 5.1. Calculating the individual values for $A_{n-1}$

We can eliminate the determinant to get a more explicit formula for (6) at a single point  $\lambda$  if all the coordinates of our starting point  $\eta$  are rational. Re-scaling by  $x_i \rightarrow cx_i$ ,  $t \rightarrow c^2t$  will make all the coordinates integers, and we can then translate all coordinates by  $-\eta_n$  so that we have  $\eta_n = 0$ . (Both of these transformations leave the Weyl chamber  $x_1 > x_2 > \dots > x_n$  unchanged.)

We now write out the normal distributions in (6) explicitly as exponentials, and expand  $(\lambda_i - \eta_j)^2$  as  $\lambda_i^2 - 2\lambda_i\eta_j + \eta_j^2$ :

$$(13) \quad b_t(\eta, \lambda) = \frac{1}{(2\pi t)^{n/2}} \det_{n \times n} \left| \exp \left( \frac{-\lambda_i^2 + 2\lambda_i\eta_j - \eta_j^2}{2t} \right) \right|.$$

Row  $i$  of this matrix contains a constant factor  $\exp(-\lambda_i^2)$ , and column  $j$  a constant factor  $\exp(-\eta_j^2)$ , so we can take these out, and put them in a constant term, which simplifies further because  $|\lambda|^2 = \sum \lambda_i^2$ . This gives us

$$(14) \quad b_t(\eta, \lambda) = \frac{1}{(2\pi t)^{n/2}} \exp \left( \frac{-|\lambda|^2 - |\eta|^2}{2t} \right) \det_{n \times n} \left| \exp \left( \frac{\lambda_i \eta_j}{t} \right) \right|.$$

Since the  $\eta_j$  are all integers, we can write the determinant as the generalized Vandermonde determinant

$$(15) \quad \det_{n \times n} |[\exp(\lambda_i/t)]^{\eta_j}|.$$

If  $\eta_j = n - j$ , this is the standard Vandermonde determinant, equal to

$$(16) \quad \prod_{i>j} [\exp(\lambda_i/t) - \exp(\lambda_j/t)].$$

And for any non-negative integers  $\eta_j$ , it is the product of this Vandermonde determinant and the Schur function [19]

$$s_{\eta_1-n+1, \eta_2-n+2, \dots, \eta_n}(\exp(\lambda_1/t), \dots, \exp(\lambda_n/t))$$

The Schur function can also be defined combinatorially [19], with the coefficient of  $\prod x_i^{n_i}$  in  $s_\mu$  the number of ways to fill in the partition

diagram of  $\mu$ , using the number  $i$  exactly  $n_i$  times, such that the entries are non-decreasing in each row and strictly increasing in each column.

In particular, we can let  $C_\eta = S_\mu(1, 1, \dots, 1)$  be the total number of such tableaux; this is important because it is the approximate value of the Schur function when the  $\lambda_i$  are much less than  $t$ . (This holds because the Schur function is a homogeneous polynomial of degree  $\eta_1 + \dots + \eta_n - (n(n-1))/2$ , with positive coefficients.) This will show the dependence of the asymptotics on the starting point.

This constant is known [19]; it is

$$(17) \quad C_\eta = \prod_{i < j} (\eta_i - \eta_j) / \prod_{i < j} (j - i).$$

This allows us to compute the constant term in the asymptotics.

For any  $\eta$  with the same sum of the coordinates, all multiples of  $1/c$ , our rescaling gives a Schur function whose index is the transformed vector  $c\eta$ . Rescaling to restore the old time values gives a Schur function which is a homogeneous polynomial in the  $\exp(\lambda_i/t)^{1/c}$  whose degree in the  $\exp(\lambda_i/t)$  is  $\eta_1 + \dots + \eta_n - (n(n-1))/2c$ . Thus, for any point  $\eta$ , even one with fractional coordinates, the degree will be bounded by the sum of its coordinates. Thus the determinants, and therefore the  $b_t(\eta, \lambda)$ , for different starting points  $\eta$  and  $\eta'$  both of radius less than a known  $\delta$ , will be in the approximate ratio of  $C_\eta$  to  $C_{\eta'}$ , with an error of  $O(\delta/t)$ . If  $\delta$  is fixed, then when we calculate the asymptotics as  $t \rightarrow \infty$ , the Schur function will converge to  $C_\eta$  at a rate of  $O(1/t)$ .

## 5.2. Asymptotic probability of no collisions

The integral of the value in (14) over the whole Weyl chamber is the probability that  $n$  particles starting at the positions  $\eta_j$  will have no collisions up to time  $t$ . We can use this formula to show that the asymptotic probability as  $t \rightarrow \infty$  is a constant multiple of  $t^{-n(n-1)/4}$ , with the constant depending on  $\eta$ . We will also show that, given that the  $n$ -dimensional Brownian motion has not hit a wall, its radial distribution, rescaled by multiplying the radius by  $1/\sqrt{t}$ , converges to the distribution of the Bessel process with parameter  $n(n+1)/2$  at time 1.

Fix  $t$  very large compared with  $|\eta|^2$ . If  $|\lambda|^2$  is much larger than  $t$ , then the exponential of  $-|\lambda|^2/2t$  in (14) will decay exponentially fast. In particular,  $\lambda_i/t$  can be assumed to be  $O(t^{\epsilon-1/2})$ , so we only need to look at the leading nonzero terms in (15); the error will be a factor of this order when compared with the value of the determinant. In (16), the first nonzero term

of the factor  $[\exp(\lambda_i/t) - \exp(\lambda_j/t)]$  is  $(\lambda_i - \lambda_j)/t$ ; we can then take out the factor of  $t^{-n(n-1)/2}$  and leave only a term involving the  $\lambda$ . Likewise, in the Schur function from (15), we need only keep the constant term  $C_\eta$ .

Thus our probability is asymptotic to the integral over the Weyl chamber of

$$(18) \quad \frac{C_\eta}{(2\pi t)^{n/2}} \exp\left(\frac{-|\lambda|^2 - |\eta|^2}{2t}\right) t^{-n(n-1)/2} \prod_{i>j} (\lambda_i - \lambda_j).$$

And this integral can be computed by using Selberg's integral [20, 22]; we have

$$(19) \quad \int_{\mathbb{R}^n} \exp(-|x|^2/2) \left| \prod_{i>j} (x_i - x_j) \right| dx_i = 2^{3n/2} \prod_{k=1}^n \Gamma((k/2) + 1).$$

This corresponds to our desired integral when we set  $\vec{x} = \lambda/\sqrt{t}$ , and divide by  $n!$  because we are taking our integral over only one of the  $n!$  different Weyl chambers. Dropping the  $\exp(-|\eta|^2/2t)$  (which goes to 1 as  $t \rightarrow \infty$  and thus doesn't affect the leading term), and writing out  $C_\eta$  explicitly again gives our full asymptotic:

$$(20) \quad \frac{\prod_{i<j} (\eta_i - \eta_j)}{\prod_{i<j} (j - i)} \frac{2^{3n/2}}{(2\pi)^{n/2} n!} \prod_{k=1}^n [\Gamma((k/2) + 1)] t^{-n(n-1)/4}.$$

We can also note that the exponential in (18) is spherically symmetric, while  $\prod_{i<j} (\lambda_i - \lambda_j)$  is homogeneous of degree  $n(n-1)/2$ . Thus the density, integrated over the sphere of radius  $r$  at a fixed  $t$ , is a constant multiple of

$$\exp(-r^2/2t) r^{(n(n-1)/2)-1},$$

and thus a constant multiple of the radial distribution of  $(n(n+1)/2)$ -dimensional Brownian motion at time  $t$ . Thus the radial distribution is exactly the same for the distribution of (18) renormalized so that the integral over the Weyl chamber is 1 (i.e., given that no two particles have collided) as for the distribution obtained from standard Brownian motion in  $n(n+1)/2$  dimensions. This proves Theorem 2; if we restrict to  $|\lambda| < ct^{1/2+\epsilon}$ , the  $\lambda_i/t$  terms are all  $O(t^{\epsilon-1/2})$ . Thus, in this region, the ratio of the radial distribution for the constrained motion to the radial distribution for unconstrained motion in  $n(n+1)/2$  dimensions converges uniformly to 1 at a rate of  $O(t^{\epsilon-1/2})$ .

Equivalently, we could fix time and  $\lambda$ , and for a scalar  $\delta$ , take  $\hat{\eta} = \delta\eta$  as our starting point. By the scaling properties of Brownian motion, this is equivalent to keeping  $\eta$  fixed, taking  $\hat{\lambda} = \lambda/\delta$ , and  $\hat{t} = t/\delta^2$ . Thus, as  $\delta \rightarrow 0$ , the ratio of the radial distributions converges uniformly to 1 within the region  $|\lambda| < ct^{1/2+\epsilon}$  at a rate of  $O(\delta^{1-\epsilon})$ , and the probability that either distribution is outside that region goes to 0 exponentially fast.

### 5.3. Asymptotics for $B_n$ : no collisions and a wall

We can use the same technique to get asymptotics for  $B_n$  as for  $A_{n-1}$ . The analogue of Theorem 1 now gives a constant multiple of  $t^{-n^2/2}$  as the asymptotic probability of no collision, and the analogue of Theorem 2 says that the distribution of the radius converges to the  $n^2 + n$ -dimensional Bessel process.

Here, we require that the coordinates all be odd integers. Again, we write out the determinant (9) explicitly:

$$(21) \quad b_t(\eta, \lambda) = \frac{1}{(2\pi t)^{n/2}} \det_{n \times n} \left| \exp \left( \frac{-\lambda_i^2 + 2\lambda_i \eta_j - \eta_j^2}{2t} \right) - \exp \left( \frac{-\lambda_i^2 - 2\lambda_i \eta_j - \eta_j^2}{2t} \right) \right|.$$

As before, we remove the constant factors to get

$$(22) \quad b_t(\eta, \lambda) = \frac{1}{(2\pi t)^{n/2}} \exp \left( \frac{-|\lambda|^2 - |\eta|^2}{2t} \right) \det_{n \times n} \left| \exp \left( \frac{\lambda_i \eta_j}{t} \right) - \exp \left( \frac{-\lambda_i \eta_j}{t} \right) \right|.$$

The determinant here is not an actual Vandermonde determinant. However, in the specific case  $\eta_j = 2n + 1 - 2j$ , we can make it a Vandermonde determinant by elementary operations. Adding  $(-1)^k \binom{2n+1-2j}{k}$  times column  $j+k$  to column  $j$  does not change the determinant, but it changes the entries in column  $j$  to

$$\begin{aligned} \sum_{k=0}^{2n+1-2j} (-1)^k \binom{2n+1-2j}{k} \exp((2n+1-2j-2k)\lambda_i/t) \\ = [\exp(\lambda_i/t) - \exp(-\lambda_i/t)]^{2n+1-2j}. \end{aligned}$$

This is a generalized Vandermonde determinant; we can make it an actual Vandermonde determinant by dividing row  $i$  by  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$ . The resulting determinant is

$$(23) \quad \det_{n \times n} \left| [\exp(\lambda_i/t) - \exp(-\lambda_i/t)]^{2(n-j)} \right|,$$

and its value is

$$(24) \quad \prod_{i>j} [(\exp(\lambda_i/t) - \exp(-\lambda_i/t))^2 - (\exp(\lambda_j/t) - \exp(-\lambda_j/t))^2].$$

Putting this together with the constants we have taken out, we get

$$\begin{aligned} b_t(\eta, \lambda) &= \frac{1}{(2\pi t)^{n/2}} \exp \left( \frac{-|\lambda|^2 - |\eta|^2}{2t} \right) \\ &\quad \times \left( \prod_{i=1}^n [\exp(\lambda_i/t) - \exp(-\lambda_i/t)] \right) \\ &\quad \times \prod_{i>j} [(\exp(\lambda_i/t) - \exp(-\lambda_i/t))^2 - (\exp(\lambda_j/t) - \exp(-\lambda_j/t))^2]. \end{aligned}$$

For a more general starting point, with all coordinates odd integers, we note that  $(\exp(\lambda_i \eta_j/t) - \exp(-\lambda_i \eta_j/t))$  is a polynomial in  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$  with no constant term. We can break these polynomials into their individual terms, giving a large number of determinants, each one a generalized Vandermonde determinant in the  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$ . Each individual determinant is thus the product of (24) and a Schur function; it also contains the product of the  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$  as a factor, since these are constant factors in row  $i$ . Thus each determinant is a product of these factors with some symmetric function in the  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$ . As with  $A_{n-1}$ , the error we get in approximating the Schur functions by their constant term  $C_\eta$  is a factor of  $O(1/t)$ .

And as with  $A_{n-1}$ , we can get asymptotics by integrating this over the Weyl chamber. For large  $t$ , the radial exponential will be exponentially small if  $|\lambda| > t^{1/2+\epsilon}$ , so we can assume that all of the  $\lambda_i/t$  are very small. Thus  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$  can be approximated by its leading nonzero term,  $2\lambda_i/t$ . Thus, as with (18), we get

$$(25) \quad \frac{1}{(2\pi t)^{n/2}} \exp \left( \frac{-|\lambda|^2 - |\eta|^2}{2t} \right) (2/t)^{n^2} \prod_{i=1}^n (\lambda_i) \prod_{i>j} (\lambda_i^2 - \lambda_j^2).$$

This integral can also be computed by using Selberg's integral [20, 22]; we have

$$\begin{aligned} & \int_{\mathbb{R}^n} \exp(-|x|^2/2) \prod_{i=1}^n |x_i| \left| \prod_{i>j} (x_i^2 - x_j^2) \right| dx_i \\ &= \frac{2^{(n^2+3n)/2}}{\pi^{n/2}} \prod_{i=1}^n [\Gamma(1+i/2)\Gamma((1+i)/2)]. \end{aligned}$$

This corresponds to our desired integral when we set  $\vec{x} = \lambda/\sqrt{t}$ , and divide by  $2^n n!$  because that is the number of Weyl chambers. Dropping the  $\exp(-|\eta|^2/2t)$  because it goes to 1 as  $1/t$ , we get our asymptotic probability that Brownian motion started at the specific point  $\eta$  will not hit a wall up to time  $t$ :

$$(26) \quad \frac{2^{(3n^2/2)}}{\pi^n n!} \prod_{k=1}^n [\Gamma(1+k/2)\Gamma((1+k)/2)] t^{-n^2/2}.$$

For a more general starting point, each individual determinant in the sum gives an integral of a homogeneous polynomial in the  $\exp(\lambda_i/t) - \exp(-\lambda_i/t)$  which is of degree at least  $n^2$ , since it contains the previous determinant as a factor. The same technique as above gives an asymptotic which is thus at most  $t^{-n^2/2}$ . Thus the asymptotic is at most a constant multiple of  $t^{-n^2/2}$ , and is less if and only if the coefficient of  $t^{-n^2/2}$  is zero. But the coefficient cannot be zero. Let  $m = \max((\eta_i - \eta_{i+1}/2), \eta_n)$ , and  $\eta'_i = (2n + 1 - 2i)m$ . Then we know that Brownian motion starting at  $\eta'$  decays asymptotically as  $t^{-n^2/2}$ , but Brownian motion starting at  $\eta$  will always hit a wall if it is translated to start at  $\eta'$ , because  $\eta'$  is at least as far as  $\eta$  from every wall. Thus the asymptotic probability will be some constant  $C_\eta$  times the formula (26); we will compute  $C_\eta$  in Section 5.5.

A result analogous to Theorem 2 also holds for  $B_n$ , using the same argument as for  $A_{n-1}$ . The exponential in (25) is spherically symmetric, while the product  $\prod(\lambda_i) \prod_{i>j} (\lambda_i^2 - \lambda_j^2)$  is homogeneous of degree  $n^2$ . The radial distribution for (25), renormalized so that the integral over the Weyl chamber is 1, is thus the same as the radial distribution for unconstrained Brownian motion in  $n^2+n$  dimensions. As with  $A_{n-1}$ , the ratio between the radial distribution for the  $B_n$  constrained motion and the radial distribution for the unconstrained motion in  $n^2$  dimensions converges uniformly to 1 at a rate of  $O(t^{\epsilon-1/2})$  inside the region  $|\lambda| < ct^{1/2+\epsilon}$ , and the probability that either distribution is outside that region goes to 0 exponentially fast. Again, we can rescale by multiplying the radius by  $1/\sqrt{t}$  to get convergence to a fixed distribution.

### 5.4. Asymptotics for $D_n$

The process for  $D_n$  is almost the same as for  $B_n$ , so we won't work it out in full detail; we get a similar result with the same error terms and convergence properties. The analogue of Theorem 1 now gives a constant multiple of  $t^{(-n^2+n)/2}$  as the asymptotic probability of no collision, and the analogue of Theorem 2 says that the distribution of the radius converges to the  $n^2$ -dimensional Bessel process.

Here, it is most natural to let  $\eta_i = n - i$ . For this value of  $\eta$ , we have only the second determinant in (12), with a plus sign between the terms; the last row of the other determinant is zero. For general  $\eta$ , the first determinant is the  $B_n$  determinant, which we know is  $O(t^{-n^2/2})$ , and we will show that the second determinant is asymptotically larger.

The determinant that we get is

$$(27) \quad \det_{n \times n} \left| \exp\left(\frac{\lambda_i \eta_j}{t}\right) + \exp\left(\frac{-\lambda_i \eta_j}{t}\right) \right|.$$

Again, this isn't a Vandermonde determinant, but  $[\exp(\lambda_i/t) + \exp(-\lambda_i/t)]^{\eta_j}$  is equal to  $\exp(\lambda_i \eta_j/t) + \exp(-\lambda_i \eta_j/t)$  plus a sum of lower order terms, so elementary operations which do not change the determinant give us the Vandermonde determinant

$$(28) \quad \det_{n \times n} |[\exp(\lambda_i/t) + \exp(-\lambda_i/t)]^{n-j}|,$$

and its value is

$$(29) \quad \prod_{i>j} [\exp(\lambda_i/t) + \exp(-\lambda_i/t) - \exp(\lambda_j/t) - \exp(-\lambda_j/t)].$$

The leading nonzero term is

$$\prod_{i>j} \frac{(\lambda_i/t)^2 - (\lambda_j/t)^2}{2},$$

and the full integral is

$$(30) \quad \frac{1}{(2\pi t)^{n/2}} \exp\left(\frac{-|\lambda|^2 - |\eta|^2}{2t}\right) (2/t)^{n^2-n} \prod_{i>j} (\lambda_i^2 - \lambda_j^2).$$

Again, we get a result which can be obtained from Selberg's integral [20, 22]; we have

$$(31) \quad \int_{\mathbf{R}^n} \exp(-|x|^2/2) \left| \prod_{i>j} (x_i^2 - x_j^2) \right| dx_i = \frac{2^{(n^2+2n)/2}}{\pi^{n/2}} \prod_{i=1}^n [\Gamma(1+i/2)\Gamma(i/2)].$$

Here, there are  $2^{n-1}n!$  Weyl chambers, which gives the asymptotic

$$(32) \quad \frac{2^{(3n^2-3n+2)/2}}{\pi^n n!} \prod_{k=1}^n [\Gamma(1+k/2)\Gamma((1+k)/2)] t^{(-n^2+n)/2}.$$

For a general starting point whose coordinates are all integers, we use the same technique as for  $B_n$ . The terms in the determinant are all polynomials in  $\exp(\lambda_i/t) + \exp(-\lambda_i/t)$ , so we can again split the sum into individual determinants, each of which is the product of (29) and a Schur function of the  $\exp(\lambda_i/t) + \exp(-\lambda_i/t)$ . Since  $\lambda_i/t$  is small, these are all close to 2, so the full asymptotic is a sum of terms of order  $t^{(-n^2+n)/2}$ . As with  $B_n$ , we can translate to  $\eta'$  which is further from any wall than  $\eta$  to show that the coefficient of  $t^{(-n^2+n)/2}$  cannot be zero.

The result analogous to Theorem 2 follows by the same argument as for the other Weyl groups. The exponential in (30) is spherically symmetric in  $\mathbb{R}^n$ , and the product  $\prod_{i>j}(\lambda_i^2 - \lambda_j^2)$  is homogeneous of degree  $n^2 - n$ . Our argument thus shows that the radial distribution converges to the  $n^2$ -dimensional Bessel process.

### 5.5. The constant factor for a general starting point

Since we know the constant factor in the asymptotic probability of no collision up to time  $t$  for one specific starting point  $\eta$ , and the asymptotic distribution for an arbitrary starting point, we can use the time-reversibility of Brownian motion to compute the asymptotic probability of no collisions for an arbitrary starting point. The argument is the same for all of the Weyl groups.

In each case, the density  $b_t(\eta, \lambda)$  for Brownian motion starting at  $\eta$  to be at  $\lambda$  at time  $t$ , not having collided with a wall up to time  $t$ , is asymptotic for large  $t$  to a product of the form

$$(33) \quad C_\eta C'_\lambda f(t) \exp\left(\frac{-|\lambda|^2 - |\eta|^2}{2t}\right),$$

in which  $C'_\lambda$  and  $f(t)$  are known. Since Brownian motion is symmetric in time, the density  $b_t(\lambda, \eta)$  must be equal to  $b_t(\eta, \lambda)$ . As long as  $t$  is large enough compared to  $|\eta|$  and  $|\lambda|$  for the formula (33) to be valid (which it will be for large  $t$  because of the exponentials), we can reverse the roles of  $\eta$  and  $\lambda$ . Thus we have  $C_\eta C'_\lambda = C_\lambda C'_\eta$ , and since the formula for  $C'$  is known, we see that  $C_\eta$  must be a constant multiple of  $C'_\eta$ ; it is thus a constant multiple of  $\prod(\eta_i - \eta_j)$  for  $A_n$ , of  $\prod(\eta_i^2 - \eta_j^2) \prod \eta_i$  for  $B_n$ , and  $\prod(\eta_i^2 - \eta_j^2)$  for  $D_n$ .

We know the value of the constant from the formulas (20), (26), and (32). For  $A_n$ , we already have the value for a general starting point because we used the Schur functions to compute  $C_\eta$  in (20); we could have instead used this technique. For  $B_n$ , we get

$$\prod_{i < j} \frac{\eta_i^2 - \eta_j^2}{(2j-1)^2 - (2i-1)^2} \prod_{i=1}^n \frac{\eta_i}{2n+1-2i} \\ \times \frac{2^{(3n^2/2)}}{\pi^n n!} \prod_{k=1}^n [\Gamma(1+k/2)\Gamma((1+k)/2)] t^{-n^2/2}.$$

as the asymptotic probability of no collision; for  $D_n$ , we get

$$\prod_{i < j} \frac{\eta_i^2 - \eta_j^2}{(2j-1)^2 - (2i-1)^2} \\ \times \frac{2^{(3n^2-3n+2)/2}}{\pi^n n!} \prod_{k=1}^n [\Gamma(1+k/2)\Gamma((1+k)/2)] t^{(-n^2+n)/2}.$$

## 6. RANDOM MATRICES AND CONDITIONED BROWNIAN MOTION

### 6.1. Brownian motion on a Lie algebra

Instead of viewing our Weyl chamber as the chamber of a Weyl group, we can view it as the space of orbits under the adjoint action of the Lie algebra corresponding to that Weyl group, and then use the theory of Lie algebras to study it. For the Lie algebras  $\mathfrak{u}_n(\mathbb{C})$  and  $\mathfrak{so}_n(\mathbb{R})$  corresponding to Weyl groups  $A_{n-1}$ ,  $B_n$ , and  $D_n$ , the Weyl chamber corresponds to the eigenvalues of the matrices in the algebra. The following construction was first developed by Dyson [10] for  $\mathfrak{u}_n(\mathbb{C})$ ; he computed the properties of the Brownian motion from the specific data rather than using the general Lie theory. The results we use from Lie theory are given in [1, 15].

Given a finite-dimensional semisimple Lie algebra  $\mathfrak{g}$ , we can define a normal distribution or Brownian motion on  $\mathfrak{g}$  by viewing it as a vector space and using its Killing form, which is non-degenerate and invariant under the action of the adjoint, as the inner product. If the Lie algebra corresponds to a compact semisimple Lie group  $G$ , the inner product is invariant under conjugation by  $G$ . For any semisimple Lie algebra, the Brownian motion at time  $t$  will have a Gaussian distribution on the Lie

algebra, with the Killing form. If the Lie algebra is an algebra of matrices, we have a Gaussian distribution on the matrices on that Lie algebra. This allows us to use all of the known results about random matrices [22] to study the motion, and its eigenvalues in particular.

In particular, on  $u_n(\mathbb{C})$ , the Lie algebra of skew-Hermitian matrices, we can take standard Brownian motion on the imaginary part of each diagonal entry  $M_{ii}$ , and  $1/\sqrt{2}$  times standard Brownian motion on each matrix entry  $M_{ij}$  with  $i < j$ , with  $M_{ji} = -\overline{M_{ij}}$ ; this is Dyson's model [10]. For a skew-Hermitian matrix, the Hilbert-Schmidt norm is the sum of the squares of the absolute values of the matrix entries, which is equal to the norm obtained from the Killing form. This makes the Hilbert-Schmidt norm of  $M$  equivalent to the radius of a Brownian motion in  $n^2$  dimensions.

Likewise, on  $\mathfrak{so}_n(\mathbb{R})$ , the Lie algebra of skew-symmetric matrices, we can take standard Brownian motion on each matrix entry  $M_{ij}$  with  $i < j$ , and take  $M_{ji} = -M_{ij}$ . This makes  $1/\sqrt{2}$  times the Hilbert-Schmidt norm of  $M$  equivalent to the radius of a Brownian motion in  $n^2 - n$  dimensions. (This is actually  $1/\sqrt{2}$  times the Killing form; we use this normalization because the eigenvalues come in pairs  $\pm i\lambda_j$ , and we want to count only one of each pair, rather than both as in the standard Hilbert-Schmidt norm.)

Since this Brownian motion is invariant under the adjoint in  $\mathfrak{g}$  or  $G$ , it induces a diffusion on the Weyl chamber. For the Weyl groups  $A_{n-1}$ ,  $B_n = C_n$ , and  $D_n$  with Lie algebras  $u_n(\mathbb{C})$ ,  $\mathfrak{so}_{2n+1}(\mathbb{R})$ ,  $\mathfrak{sp}_{2n}(\mathbb{R})$ , and  $so_{2n}(\mathbb{R})$ , the Weyl chamber is in a natural correspondence with the space of eigenvalues, obtained by dividing the independent eigenvalues by  $i$  and arranging them in decreasing order; for other Lie algebras, we can study the Weyl chamber, but cannot place coordinates on it which directly correspond to the eigenvalues of random matrices. We can use the Lie algebra analogue of the Weyl Integration formula for Lie groups to study the process in the Weyl chamber in terms of the corresponding process on  $\mathfrak{g}$ .

**LEMMA 1** [6, IX 6.3.2 (11)]. – *Let  $\mathfrak{g}$  be a finite-dimensional Lie algebra with  $m$  roots. Let  $\delta$  be  $(2\pi)^m$  times the product of all the positive roots of  $\mathfrak{g}$ . The Jacobian of the adjoint map on  $\mathfrak{g}$  at a point  $x$  is  $\delta(x)\bar{\delta}(x)$ .*

**COROLLARY 6** [6, IX 6.3.4 (13)]. – *If  $f$  is a function on  $\mathfrak{g}$  which is invariant under the adjoint map, the integral of  $f$  over the Lie algebra  $\mathfrak{g}$  is equal to the integral of  $\delta\bar{\delta}f$  over the Weyl chamber.*

This shows that the measure induced from  $\mathfrak{g}$  at a point in the Weyl chamber is proportional to  $\delta\bar{\delta}$ . The distribution in the Weyl chamber of the induced diffusion started at the origin is thus proportional to  $\delta\bar{\delta}$  times the distribution in  $\mathfrak{g}$  of standard Brownian motion started at the origin.

In particular, we note that  $\delta$  is positive real on the interior of the Weyl chamber, and zero on the walls. This suggests that we have a process which can never touch a wall of the Weyl chamber; we will prove this later by showing that  $1/\delta$  is a martingale for the process on the Weyl chamber.

## 6.2. Construction by $h$ -transformation

We can also use Doob's  $h$ -transformation [9] to construct a process which always stays in the Weyl chamber. We will first construct the conditioned Brownian motion in this way, and then show that the two processes are actually identical; this allows us to prove Theorem 3 and develop the physical model of Theorem 4. This method was developed by Biane [4] on  $A_{n-1}$ , and generalizes naturally to other Lie groups.

**6.2.1. General properties.** – Given standard Brownian motion in any number of dimensions, we can use the process of  $h$ -transformation to construct a Brownian motion satisfying certain conditions. For any non-negative harmonic function  $h$ , the measure for the transformed Brownian motion to go to  $\lambda$  at time  $t$  after starting at  $\eta$  is  $h(\lambda)$  times the measure for untransformed Brownian motion with the same starting point, renormalized by an appropriate constant so that the total measure on all paths is 1.

In particular, suppose that we have an open connected region  $D$  and a harmonic function  $h$  which is zero on the boundary of the region and positive on the interior. Then the function  $1/h$  (taken to have value  $+\infty$  on the boundary of  $D$ ) is a martingale for  $h$ -transformed Brownian motion [9, 2.X.1]. If our starting point is inside  $D$  and we take as a stopping time either a fixed  $t$  or the time that the process reaches the boundary, the expectation must be finite, and thus the probability that Brownian motion reaches the boundary before time  $t$  must be zero. Thus the  $h$ -transformed Brownian motion will be conditioned to stay in the interior of the region.

The density function for this transformed motion will be  $h$  times the density function for the untransformed motion on the same region, normalized appropriately. That is, its value at a point  $\lambda$  and time  $t$  will be  $h(\lambda)$  times the value  $b_t(\eta, \lambda)$  which gives the measure for unconstrained motion to go from  $\eta$  to  $\lambda$  in time  $t$  while staying within the region, since the transformation puts a new measure on the same set of paths.

The same technique can also be applied to a discrete random walk with a set  $S$  of steps, provided that  $h$  is harmonic in the discrete lattice; that is, we need

$$(34) \quad \sum_{s \in S} h(x + s) / |S| = h(x).$$

The transformed discrete walk now has probability  $h(x + s)/(h(x)|S|)$  instead of  $1/|S|$  of going from  $x$  to  $x + s$  in a given step. Thus, given a starting point  $\eta$ , the probability of going to  $\lambda$  in a given number of steps of the transformed random walk is  $h(\lambda)/h(\eta)$  times the probability for the untransformed walk to go to  $\lambda$  while staying within the region in which  $h$  is positive.

In order to make it impossible for the walk to leave the region  $D$ , we need  $h$  to be zero on all points which can be reached from the interior of  $D$  in a single step. Thus, only if the discrete walk is reflectable (as defined above) is it sufficient for  $h$  to be zero on the continuous boundary of  $D$ .

**6.2.2. Finding the function.** – The properties of conditioning, as well as our asymptotics, suggest that  $\delta$  itself should be our  $h$ . It can be checked algebraically that  $h$  is harmonic for each group; however, it can also be proved naturally.

**THEOREM 7.** – *For any finite Coxeter group  $W$ , the product of all the positive roots is a harmonic function, both for the continuous Laplacian and for the discrete Laplacian*

$$L_S h(\vec{x}) = \frac{1}{|S|} \sum_{\vec{s} \in S} [h(\vec{x} + \vec{s}) - h(\vec{x})],$$

for any set  $S$  which is symmetric under the group  $W$ .

*Proof.* – This result in the continuous case is due to [2]; the discrete argument is a simple generalization which is mentioned in [4].

By the properties of root systems, a reflection in any simple root changes the sign of only that root, while permuting the other positive roots. Thus the product  $h$  of all  $m$  roots is antisymmetric in all the simple roots, and by applying repeated reflections, we see that it is antisymmetric in every root. It is also of degree  $m$ .

The continuous Laplacian is spherically symmetric, and thus symmetric under  $W$ . The discrete Laplacian is symmetric under  $W$  because the set  $S$  is. Applying the continuous Laplacian to a polynomial decreases the degree by 2, while applying the discrete Laplacian decreases the degree by at least 1 since  $h(\vec{x} + \vec{s}) - h(\vec{x})$  is of lower degree than  $h$ . Thus the application of the Laplacian to  $h$  gives a polynomial which is of degree less than  $m$  which is still antisymmetric in  $W$ .

Now, any polynomial which is antisymmetric in  $W$  must be zero on every one of the hyperplanes of reflection. If it is not identically zero, it must have all  $m$  roots as factors, so it must be of degree at least  $m$ . Thus

the Laplacian must annihilate our polynomial  $h$ , so  $h$  is harmonic for either the discrete or continuous walk.

It follows that the transformation by this  $h$  gives the same process as the process generated by Lie theory. We have already computed the asymptotic density that Brownian motion started at a fixed  $\eta$  will remain in the Weyl chamber for time  $t$  and be at  $\lambda$  at that time. Transforming by  $h$  has the effect of multiplying the measure of all paths from a fixed  $\eta$  to an arbitrary  $\lambda$  which stay within the chamber by a factor of  $h(\lambda)$  (and a normalizing constant). Thus the density function for the transformed Brownian motion which starts at  $\eta$  to be at  $\lambda$  at time  $t$  is proportional to  $h(\lambda)b_t(\eta, \lambda)$ , and this converges to a constant multiple of  $h(\lambda)^2$  as  $t$  becomes large. This is the same factor  $\delta\bar{\delta}$  which we obtained from the Weyl Integration Formula. In particular, this now shows that the Lie theory process also stays within the Weyl chamber almost surely.

### 6.3. The radial process

Since the  $h$ -transformed process is identical to the process generated by Lie theory, we can prove Theorem 3 and the analogous theorem for any finite Weyl group, generalizing the argument of Biane [4] for  $A_{n-1}$ . We again use the Killing form to obtain a norm on  $\mathfrak{g}$  as a vector space. If there are  $m$  positive roots, then  $\mathfrak{g}$  has dimension  $n + 2m$  [15]. The norm is thus given by the Bessel process in  $n + 2m$  dimensions. The Killing form is invariant under the adjoint, so the diffusion in the Weyl chamber has its radius given by the same process.

To properly state this result for Brownian motion conditioned never to leave the Weyl chamber, we cannot allow the process to start at the origin, which is not a point in the chamber. However, the square of the product of the roots is a homogeneous function of degree  $2m$ ; it thus gives an identical distribution on any fixed radius. Thus, if we start the motion on the Lie algebra at 0, then at any later time, given the fixed radius, the distribution on the sphere will be given by the square of the product of the roots, and the process can be continued from that time on as a Bessel process. This allows us to state the general theorem, formalizing Theorem 3.

**THEOREM 8.** – *For any Weyl group  $W$  acting on  $\mathbb{R}^n$  with  $m$  roots, consider the process which starts at a fixed radius  $r_0$ , with the starting point chosen on the sphere of radius  $r_0$  by a distribution with density proportional to the square of the product of the roots. Then the radius of the position of transformed motion at time  $t$  gives a Bessel process with parameter  $n + 2m$ .*

The simplest case of this theorem is for  $B_1$ , which is one-dimensional motion with an absorbing boundary at  $x = 0$ . In this case, the only root is  $x$ , so  $m = 1$ , and we have the result of [24] and [28] that Brownian motion conditioned never to hit 0 is the same as the three-dimensional Bessel process. In our more general cases, the number of dimensions for the Bessel process is  $n + 2m = n^2$  for  $A_{n-1}$  (acting in  $n$  dimensions because the Lie algebra is  $\mathfrak{u}_n(\mathbb{C})$ ; it acts in  $n - 1$  dimensions for the Lie algebra  $\mathfrak{su}_n(\mathbb{C})$ , giving a Bessel process in  $n^2 - 1$  dimensions),  $n(2n + 1)$  for  $B_n$ , and  $n(2n - 1)$  for  $D_n$ .

#### 6.4. The infinitesimal generator and physical models

We have constructed the conditioned process as a transformation of Brownian motion. We can also construct it as a diffusion with its known infinitesimal generator. In this form, both the multidimensional models and the models of  $n$  independent particles lead to natural physical models.

We will use the notation of [18] for infinitesimal generators of diffusions. The drift vector  $\mu$  is defined by

$$\mu_i(\vec{x}, t) = \lim_{\Delta t \downarrow 0} \frac{1}{\Delta t} \{E(X_i(t + \Delta t)) - X_i(t) | \vec{X}(t) = \vec{x}\},$$

and the infinitesimal variance matrix is defined by

$$\begin{aligned} \sigma_{i,j}(\vec{x}, t) &= \lim_{\Delta t \downarrow 0} \frac{1}{\Delta t} \\ &\{[E(X_i(t + \Delta t)) - X_i(t)][E(X_j(t + \Delta t)) - X_j(t)] | \vec{X}(t) = \vec{x}\}, \end{aligned}$$

We will omit the variable  $t$  in the infinitesimal generators of our diffusions, because they are independent of time.

It can easily be checked that Brownian motion transformed by a harmonic (and thus necessarily  $C^2$ ) function  $h$  has infinitesimal drift  $\mu = \nabla h/h = \nabla(\log h)$  and infinitesimal variance  $\sigma_{ij} = \delta_{ij}$ . [4]

This makes the computation easy, because our  $h$  is the product  $\prod_{\alpha \in \Delta} \alpha$  of all the roots, viewed as linear functions of the  $x_i$ . If we write  $\alpha(\vec{x})$  as the dot-product  $(\alpha, \vec{x})$ , then we have

$$(35) \quad \frac{\nabla h}{h} = \sum_{\alpha \in \Delta} \nabla(\alpha, \vec{x}) = \sum_{\alpha \in \Delta} \alpha \frac{1}{(\alpha, \vec{x})}.$$

Since  $(\alpha, \vec{x})/|\alpha|$  is the distance from  $\vec{x}$  to the hyperplane orthogonal to  $\alpha$ , while  $\alpha/|\alpha|$  is the unit vector in the direction of  $\alpha$ , this term in the drift

is the inverse of the distance between  $\vec{x}$  and the hyperplane, directed away from the hyperplane. We thus have the following physical model.

**THEOREM 9.** – *For any finite Coxeter group, Brownian motion conditioned to stay within a chamber is equivalent to the motion of a particle in a diffusion with constant infinitesimal velocity, and a repulsive force from every hyperplane of reflection (not merely the walls of the chamber) inversely proportional to the distance from that hyperplane.*

We can also look at the  $n$  coordinates as individual motions in one dimension. If the root  $\alpha$  contains  $cx_i$ , we get a term  $c/(\alpha, \vec{x})$  in the sum.

In particular, for  $A_n$ , the drift  $\mu_i$  is  $\sum_{i \neq j} 1/(x_i - x_j)$ . Thus each particle is subject to a repelling force from every other particle (not merely its neighbors), inversely proportional to the distance between them. This proves Theorem 4; this result is originally due to Dyson [10].

The model for  $B_n$  is not as natural as a model of particles. The drift  $\mu_i$  is

$$\frac{1}{x_i} + \sum_{i \neq j} \left( \frac{1}{x_i - x_j} + \frac{1}{x_i + x_j} \right)$$

That is, each particle is repelled by every other particle, and by the wall at 0 (the  $1/x_i$  term), but also by the mirror image of every other particle (the  $1/(x_i + x_j)$  term), as if the wall at  $x = 0$  was also a mirror reflecting all forces. For  $D_n$ , we have only the terms of  $1/(x_i - x_j)$  and  $1/(x_i + x_j)$ ; this means that the mirror reflects forces but is itself permeable to particles.

These models are more natural as models of the eigenvalues. For  $D_n$ , a matrix in the Lie algebra  $\mathfrak{so}_{2n}(\mathbb{R})$  has eigenvalues  $\pm i\lambda_j$ ; this model thus says that the eigenvalues in different pairs  $\pm i\lambda_j$  and  $\pm i\lambda_k$  for  $j \neq k$  repel one another other, although the pair  $\pm i\lambda_j$  do not repel each other. For  $B_n$ , a matrix in the Lie algebra  $\mathfrak{so}_{2n+1}(\mathbb{R})$  has eigenvalues  $\pm i\lambda_j$  and 0; this model thus says that the unconnected eigenvalues  $\pm i\lambda_j$  and  $\pm i\lambda_k$  for  $j \neq k$  repel each other, and each eigenvalue is also repelled by the fixed eigenvalue at 0.

## 7. OPEN PROBLEMS

The discussion of Brownian motion on a Lie algebra is valid for the exceptional Lie groups as well, but the techniques for computing the specific asymptotics do not appear to work. For a chamber of a general Coxeter group, the Lie algebra technique is not meaningful. In either case,

is it possible to get the same type of asymptotics, with constant terms in particular?

We have shown that the same harmonic functions which we used to transform Brownian motion can be used to transform discrete random walks. Is it possible to use these results to compute asymptotics for the discrete walks, including the constants on leading terms?

Brownian motion can be defined on a general manifold [21]. This allows us to apply the argument of Theorem 5 whenever we have a suitable chamber. As before, the Brownian motion must be symmetric under the reflections in any wall of the chamber, and the reflections in the walls must generate a discrete group which partitions the manifold into chambers. For example, since 2-dimensional Brownian motion is conformally invariant, we can define a Brownian motion on the modular surface [21]. Our chamber can be the standard fundamental domain; if we use the standard map of the modular surface to the upper half-plane, our chamber is bounded by  $x > -1/2$ ,  $x < 1/2$ , and  $x^2 + y^2 > 1$ . Can the resulting formulas be used to compute properties of this Brownian motion, such as asymptotic survival probabilities, hitting times, and physical models?

#### ACKNOWLEDGEMENTS

Sections 2-4 are part of my thesis, "Walks and Representation Theory," which was written at Harvard University. I would like to thank Peter Magyar for his collaboration in work on the discrete problem; Itai Benjamini and Gil Kalai, for suggestions related to the asymptotic results; Arun Ram, for a suggestion which simplified the computation of the determinant formulas; Dan Stroock, for suggesting that I study reflecting boundary conditions; the referee of an earlier version of this paper, for suggesting the Lie theory argument and connection to random matrices; Wendelin Werner, for suggesting some important references; my thesis advisors, Persi Diaconis and Richard Stanley, for their guidance in my thesis research; and Yuval Peres, for his guidance in the research beyond my thesis.

#### REFERENCES

- [1] J. F. ADAMS, Lectures on Lie Groups, University of Chicago Press, 1969.
- [2] P. BÉRARD and G. BESSON, Spectres et groupes cristallographiques II: domaines sphériques, *Ann. Inst. Fourier*, Vol. 30, 3, 1980, pp. 237–248.
- [3] P. BIANCE, Minuscule weights and random walks on lattices, *Quant. Prob. Rel. Topics*, Vol. 7, 1992, pp. 51–65.

- [4] P. BIANE, Permutation model for semi-circular systems and quantum random walks, *Pacific J. Math.*, Vol. **171**, 1995, pp. 373–387.
- [5] N. BOURBAKI, Groupes et Algèbres de Lie, Chapters 4, 5, 6. Hermann, Paris, 1968.
- [6] N. BOURBAKI, Groupes et Algèbres de Lie, Chapter 9. Masson, Paris, 1982.
- [7] D. L. BURKHOLDER, Exit times of Brownian motion, harmonic majorization, and Hardy spaces, *Adv. Math.*, Vol. **26**, 1977, pp. 182–205.
- [8] R. D. DEBLASSIE, Exit times from cones in  $\mathbb{R}^n$  of Brownian motion, *Prob. Th. Rel. Fields*, Vol. **74**, 1987, pp. 1–29.
- [9] J. L. DOOB, Classical Potential Theory and its Probabilistic Counterpart, Springer-Verlag, New York, 1984.
- [10] F. J. DYSON, A Brownian-motion model for the eigenvalues of a random matrix, *J. Math. Phys.*, Vol. **3**, 1962, pp. 1191–1198.
- [11] D. FREEDMAN, Brownian Motion and Diffusion, Holden-Day, San Francisco; second edition published by Springer-Verlag, New York, 1971.
- [12] I. M. GESSEL and D. ZEILBERGER, Random walk in a Weyl chamber, *Proc. Amer. Math. Soc.*, Vol. **115**, 1992, pp. 27–31.
- [13] D. J. GRABINER and P. MAGYAR, Random walks in Weyl chambers and the decomposition of tensor powers, *J. Alg. Combin.*, Vol. **2**, 1993, pp. 239–260.
- [14] J. M. HARRISON, Brownian Motion and Stochastic Flow Systems, John Wiley and Sons, New York, 1985.
- [15] J. E. HUMPHREYS, Introduction to Lie Algebras and Representation Theory, Springer-Verlag, New York, 1972.
- [16] D. HOBSON and W. WERNER, Non-colliding Brownian motion on the circle, *Bull. London Math. Soc.*, Vol. **28**, 1996, pp. 643–650.
- [17] S. P. KARLIN and G. MACGREGOR, Coincidence probabilities, *Pacific. J. Math.*, Vol. **9**, 1959, pp. 1141–1164.
- [18] S. P. KARLIN and H. M. TAYLOR, A Second Course in Stochastic Processes, Academic Press, New York, 1981.
- [19] I. G. MACDONALD, Symmetric Functions and Hall Polynomials, Oxford University Press, 1979.
- [20] I. G. MACDONALD, Some conjectures for root systems, *SIAM J. Math. Anal.*, Vol. **13**, 1982, pp. 988–1007.
- [21] H. P. MCKEAN, Stochastic Integrals, Academic Press, New York, 1969.
- [22] M. L. MEHTA, Random Matrices: Second Edition, Academic Press, New York, 1991.
- [23] N. O'CONNELL and A. UNWIN, Collision times and exit times from cones: A duality, *Stoch. Proc. Appl.*, Vol. **43**, 1992, pp. 291–301.
- [24] J. PITMAN, One-dimensional Brownian motion and the three-dimensional Bessel process, *Ann. Appl. Prob.*, Vol. **7**, 1975, pp. 511–526.
- [25] R. A. PROCTOR, Reflection and algorithm proofs of some more Lie group dual pair identities, *J. Combin. Th. A*, Vol. **62**, 1993, pp. 107–127.
- [26] A. SELBERG, Bemerkinger om et multipelt integral, *Norsk Matematisk Tidsskrift*, Vol. **26**, 1944, pp. 71–78.
- [27] T. WATANABE and S. G. MOHANTY, On an inclusion-exclusion formula based on the reflection principle, *Discrete Math.*, Vol. **64**, 1987, pp. 281–288.
- [28] D. WILLIAMS, Path decomposition and continuity of local time for one-dimensional diffusions, *Proc. London Math. Soc.*, Vol. **28**, 1974, pp. 738–768.
- [29] D. ZEILBERGER, Andre's reflection proof generalized to the many-candidate ballot problem, *Discrete Math.*, Vol. **44**, 1983, pp. 325–326.

(Manuscript received November 6, 1997;  
Revised August 5, 1998.)