

COMPOSITIO MATHEMATICA

R. S. DAHIYA

Certain theorems on n-dimensional operational calculus

Compositio Mathematica, tome 18, n° 1-2 (1967), p. 17-24

http://www.numdam.org/item?id=CM_1967__18_1-2_17_0

© Foundation Compositio Mathematica, 1967, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Certain theorems on n -dimensional operational calculus

by

R. S. Dahiya

1

In the present paper we obtain a few theorems in n -dimensional operational calculus starting from given operational relations in one variable. These theorems are next applied to the evaluation of a few operational results. In the theorems, we shall take $D(\sigma_{10}, \dots, \sigma_{n0})$ to represent the set of points for which $R(P_i) > R(P_{i0}) = \sigma_{i0} > 0$ and unless otherwise stated, D will represent the set of points for which $R(P_i) > 0$, $i = 1, 2, \dots, n$.

2

THEOREM 1. Let

- (i) $\phi(p) \subset f(x)$
- (ii) $p\sigma(p) \subset f(x^2)$
- (iii) $H(p) \subset x^{-n}\phi(1/x^2)$

where $f(x)$, $f(x^2)$ and $x^{-n}\phi(1/x^2)$ are each integrable L in $(0, \infty)$ or the definition integral in each of (i), (ii), (iii) is absolutely convergent for $R(P) > 0$.

If x_1, \dots, x_n , be a set of real numbers > 0 , then

$$(2.1) \quad \frac{\sigma(\sqrt{1/x_1 + \dots + 1/x_n})}{(x_1 \dots x_n)^{\frac{n}{2}}} \supset_n \frac{\pi^{(n-1)/2} 2^n (p_1 \dots p_n)}{\sqrt{p_1 + \dots + p_n}} \cdot H(\sqrt{p_1 + \dots + p_n})$$

provided that

(a) the integral in (2.1) exists as an absolutely convergent integral for $(p_1, \dots, p_n) \in D(\sigma_{10}, \dots, \sigma_{n0})$; or

(b) the original in (2.1) is integrable L in $(0, \infty)$ with respect to the variables x_1, \dots, x_n ; (the operational variables p_1, \dots, p_n corresponding to x_1, \dots, x_n).

From (i), we have

$$(2.2) \quad f(x^2) \supset \frac{p}{\sqrt{\pi}} \int_0^\infty e^{-\frac{1}{2}p^2 u^2} \phi\left(\frac{1}{u^2}\right) du$$

where the integrals on the right is absolutely convergent. So that, by (ii), we can write (2.2) as

$$(2.3) \quad \sigma(p) = \frac{1}{\sqrt{\pi}} \int_0^\infty e^{-\frac{1}{2}p^2 u^2} \phi\left(\frac{1}{u^2}\right) du$$

writing (2.3) in the form

$$(2.4) \quad \frac{\pi^{(1-n)/2} \sigma(\sqrt{1/x_1 + \dots + 1/x_n})}{(x_1 \dots x_n)^{\frac{n}{2}}} \\ = \pi^{-n/2} \int_0^\infty (x_1 \dots x_n)^{-\frac{n}{2}} \exp\left(-\frac{1}{4} \sum \frac{u^2}{x_i}\right) \phi\left(\frac{1}{u^2}\right) du.$$

We multiply both sides by $\exp(-\sum p_i x_i)$, integrate with respect to x_i between the limits $(0, \infty)$ and then change the order of integration in the resulting integral on the right, permissible by Fubini's theorem, on account of the absolute convergence.

This gives

$$(2.5) \quad \pi^{(1-n)/2} \int_0^\infty \dots \int_0^\infty \exp(-\sum p_i x_i) \frac{\sigma(\sqrt{1/x_1 + \dots + 1/x_n})}{(x_1 \dots x_n)^{\frac{n}{2}}} dx_1 \dots dx_n \\ = \pi^{-n/2} \int_0^\infty \phi\left(\frac{1}{u^2}\right) du \int_0^\infty x_1^{-\frac{n}{2}} \exp\left(-\frac{1}{4} \frac{u^2}{x_1} - p_1 x_1\right) dx_1 \dots \\ \int_0^\infty x_n^{-\frac{n}{2}} \exp\left(-p_n x_n - \frac{1}{4} \frac{u^2}{x_n}\right) dx_n.$$

Evaluating the inner integrals on the right by

$$(A) \quad \int_0^\infty x^{-\frac{n}{2}} e^{-px - \frac{1}{4}(a^2)/x} dx = \frac{2\sqrt{\pi}}{a} e^{-a\sqrt{p}}.$$

We obtain, on account of the operational relation (iii), the required theorem.

3

THEOREM 2. *Let*

- (i) $\phi(p) \subset f(x)$
 (ii) $\psi(p) \subset x^{-\frac{1}{2}}\phi\left(\frac{1}{x}\right)$
 (iii) $\lambda(p) \subset x^{1-n}f(x^4)$

where $f(x)$, $x^{-\frac{1}{2}}\phi(1/x)$ & $x^{1-n}f(x^4)$ are each integrable L in $(0, \infty)$ or the definition integral in each of (i), (ii) and (iii) is absolutely convergent for $R(P) > 0$. If x_1, \dots, x_n be a set of positive real numbers, then

$$(3.1) \quad \frac{\psi\left[\frac{1}{64}(1/x_1 + \dots + 1/x_n)^2\right]}{(x_1 \dots x_n)^{\frac{1}{2}}(1/x_1 + \dots + 1/x_n)^2} \supset_n \frac{2^{n-4}\pi^{(n+1)/2}(p_1 \dots p_n)}{\sqrt{p_1 + \dots + p_n}} \cdot \lambda(\sqrt{p_1 + \dots + p_n})$$

provided that

(a) the definition integral in (3.1) is absolutely convergent for $(p_1, \dots, p_n) \in D(\sigma_{10}, \dots, \sigma_{n0})$; or

(b) the original in (3.1) is integrable L in $(0, \infty)$ with respect to the variables x_1, \dots, x_n ; (the operational variables p_1, \dots, p_n corresponding to x_1, \dots, x_n).

4

THEOREM 3. *Let*

- (i) $\phi(p) \subset f(x)$
 (ii) $\phi(\sqrt{p}) \subset F\left(\frac{1}{x}\right)$
 (iii) $H(p) \subset \frac{1}{x^n}f(x)$.

Then

$$(4.1) \quad \frac{F(1/x_1 + \dots + 1/x_n)}{(x_1 \dots x_n)^{\frac{1}{2}}(1/x_1 + \dots + 1/x_n)^{\frac{1}{2}}} \supset_n \frac{2^n \pi^{(n-1)/2}(p_1 \dots p_n)}{\sqrt{p_1 + \dots + p_n}} \cdot H(\sqrt{p_1 + \dots + p_n}).$$

The proofs of theorems 2 and 3 are on the same lines.

5

As applications of theorems 1, 2 and 3, we shall find some two-dimensional correspondences.

(a) Let

$$f(x) = J_{2n,n}(\mathfrak{B}\sqrt[3]{x}); \text{ so that } \phi(p) = J_{2n}\left(\frac{2}{\sqrt{p}}\right)$$

$$f(x^2) = J_{2n,n}(\mathfrak{B}\sqrt[3]{x^2}); \text{ so that } \sigma(p) = \frac{1}{p} e^{-(2/p^2)} I_n\left(\frac{2}{p^2}\right)$$

and $n = 2$, then

$$x^{-n}\phi\left(\frac{1}{x^2}\right) = x^{-2}J_{2n}(2x);$$

so that

$$H(p) = \frac{\Gamma(2n-1)}{\Gamma(2n+1)p^{2n-2}} {}_2F_1\left[\begin{matrix} n-\frac{1}{2}, n; \\ 2n+1; \end{matrix} -\frac{4}{p^2}\right].$$

Hence we obtain, from (2.1)

$$\begin{aligned} \frac{e^{-(2xy)/(x+y)}}{xy(x+y)^{\frac{1}{2}}} I_n\left(\frac{2xy}{x+y}\right) \supset_2 \frac{2\sqrt{\pi}}{n(2n-1)} \frac{pq}{(\sqrt{p}+\sqrt{q})^{2n-1}} \\ \cdot {}_2F_1\left[\begin{matrix} n-\frac{1}{2}, n; \\ 2n+1; \end{matrix} -\frac{4}{(\sqrt{p}+\sqrt{q})^2}\right]. \end{aligned}$$

(b) Let

$$f(x) = x^{\nu/2} J_{\nu}(2a\sqrt{x});$$

so that

$$\phi(p) = a^{\nu} p^{-\nu} e^{-a^2/p},$$

$$f(x^2) = x^{\nu} J_{\nu}(2ax);$$

so that

$$p\sigma(p) = \frac{a^{\nu} 2^{2\nu} \Gamma(\nu+\frac{1}{2})p}{\sqrt{\pi}(p^2+4a^2)^{\nu+\frac{1}{2}}}, \quad R(\nu) > -\frac{1}{2},$$

$$x^{-n}\phi\left(\frac{1}{x^2}\right) = a^{\nu} x^{2\nu-2} e^{-a^2 x^2};$$

$$\text{so that } H(p) = \frac{\Gamma(2\nu-2)p}{(8a^2)^{\nu-\frac{1}{2}}} e^{(p^2/8a^2)} D_{2\nu-1}\left(\frac{p}{\sqrt{2a}}\right), \quad R(a) < 0.$$

Hence from (2.1), we get

$$\begin{aligned} \frac{(xy)^{\nu-1}}{(x+y+4a^2xy)^{\nu+\frac{1}{2}}} \supset_2 \frac{\sqrt{\pi}\Gamma(2\nu-1)}{a^{3\nu-1}2^{5\nu-\frac{3}{2}}} \\ pq e^{((\sqrt{p}+\sqrt{q})^2/8a^2)} D_{2\nu-1}\left(\frac{\sqrt{p}+\sqrt{q}}{\sqrt{2a}}\right), \quad \begin{matrix} R(\nu) > -\frac{1}{2} \\ R(a) < 0 \end{matrix}. \end{aligned}$$

(c) Let

$$f(x) = \sqrt{x} L_1(\sqrt{x});$$

so that
$$\phi(p) = \frac{1}{2p} e^{\frac{1}{2}p} \operatorname{Erf} e \left(\frac{1}{2\sqrt{p}} \right)$$

$$f(x^2) = x L_1(x);$$

so that
$$p\sigma(p) = \frac{p}{(p^2-1)^{\frac{3}{2}}} - \frac{2\sqrt{2}p^{-\frac{1}{2}}}{\sqrt{\pi}(1-p^2)^{-\frac{1}{2}}} P^{-\frac{1}{2}} \left(\frac{1}{p} \right)$$

$$x^{-n} \phi \left(\frac{1}{x^2} \right) = \frac{1}{2} e^{x^2/4} \operatorname{Erf} c \left(\frac{x}{2} \right);$$

so that
$$H(p) = \frac{p}{2\sqrt{\pi}} e^{-p^2} \operatorname{Ei}(p^2).$$

Hence from (2.1), we get

$$\begin{aligned} & 2pq e^{-(\sqrt{p}+\sqrt{q})^2} \operatorname{Ei}(\sqrt{p}+\sqrt{q}) C_2 \frac{1}{(x+y-xy)^{\frac{3}{2}}} \\ & - \frac{2\sqrt{2}(xy-x-y)^{\frac{3}{2}}}{\sqrt{\pi}(xy)^{\frac{3}{2}}(x+y)^{\frac{3}{2}}} P^{-\frac{1}{2}} \left(\sqrt{\frac{xy}{x+y}} \right). \end{aligned}$$

(d) Let

$$f(x) = \frac{x^\nu}{\Gamma(\nu+1)};$$

so that
$$\sigma(p) = \frac{1}{p^\nu}; \quad R(\nu) > -1.$$

$$f(x^2) = \frac{x^{2\nu}}{\Gamma(\nu+1)};$$

so that
$$\sigma(p) = \frac{\Gamma(2\nu+1)}{\Gamma(\nu+1)} p^{-1-2\nu}, \quad R(\nu) > -\frac{1}{2}$$

and $n = 3$, then

$$x^{-n} \phi \left(\frac{1}{x^2} \right) = x^{2\nu-3}; \quad \text{so that } H(p) = \Gamma(2\nu-2) p^{3-2\nu}.$$

Thus we obtain, from (2.1)

$$\begin{aligned} & \frac{(xyz)^{\nu-1}}{(xy+yz+zx)^{\nu+\frac{1}{2}}} \supset_3 \frac{8\pi\Gamma(2\nu-2)\Gamma(\nu+1)}{\Gamma(2\nu+1)} \\ & \times \frac{pqr}{(\sqrt{p}+\sqrt{q}+\sqrt{r})^{2\nu-2}}, \quad R(\nu) > -\frac{1}{2}. \end{aligned}$$

(e) Let

$$f(x) = (2x)^{\nu-1} e^{-2\sqrt{x}};$$

so that
$$\phi(p) = \Gamma(2\nu) p^{1-\nu} e^{1/2\nu} D_{-2\nu} \left(\sqrt{\frac{2}{p}} \right)$$

$$x^{-\frac{1}{2}} \phi(x) = \Gamma(2\nu) x^{\nu-\frac{3}{2}} e^{x/2} D_{-2\nu}(\sqrt{2x});$$

so that
$$\psi(p) = \frac{\sqrt{2\pi} \Gamma(2\nu-1) p}{2^{\nu-\frac{1}{2}} (p-1)^{\nu-\frac{1}{2}}} P_{-1}^{1-2\nu} \left(\frac{1}{\sqrt{p}} \right), R(\nu) > \frac{1}{2}$$

and $n = 2$, then

$$x^{1-n} f(x^4) = 2^{\nu-1} x^{4\nu-5} e^{-2x^2};$$

so that
$$\lambda(p) = \frac{\Gamma(4\nu-4)}{2^{8(\nu-1)}} p e^{(p^2/16)} D_{4-4\nu} \left(\frac{p}{2} \right).$$

Hence from theorem 2, we obtain

$$\begin{aligned} & \frac{(xy)^{2\nu-\frac{1}{2}}}{[(x^2+y^2)-64x^2y^2]^{\nu-\frac{1}{2}}} P_{-1}^{1-2\nu} \left(\frac{xy}{8(x+y)} \right) \\ & \supset_2 \frac{\pi \Gamma(4\nu-4) pq}{2^{8\nu-4} \Gamma(2\nu-1)} e^{(\sqrt{p}+\sqrt{q})^2/16} D_{4-4\nu} \left(\frac{\sqrt{p}+\sqrt{q}}{2} \right), R(\nu) > +\frac{1}{2}. \end{aligned}$$

(f) Let

$$f(x) = x^{\nu-1} e^{-(x^2/8a)};$$

so that
$$\phi(p) = 2^\nu a^{\nu/2} \Gamma(\nu) p e^{a p^2} D_{-\nu}(2\sqrt{ap})$$

$$\phi(\sqrt{p}) = \Gamma(\nu) 2^\nu a^{\nu/2} \sqrt{p} e^{a p} D_{-\nu}(2\sqrt{ap});$$

so that
$$F \left(\frac{1}{x} \right) = \frac{\Gamma(\nu)(2a)^{\nu/2}}{\Gamma((\nu+1)/2)} x^{(\nu-1)/2} (x+2a)^{-\nu/2}$$

and $n = 2$, then

$$x^{-n} f(x) = x^{\nu-3} e^{-(x^2/8a)};$$

so that
$$H(p) = 2^{\nu-2} a^{(\nu/2)-1} \Gamma(\nu-2) p e^{a p^2} D_{2-\nu}(2\sqrt{ap}).$$

Hence from (4.1), we obtain

$$\begin{aligned} & \frac{(xy)^{(\nu-3)/2}}{(x+y)^{\nu/2} (xy/(x+y)+2a)^{(\nu+1)/2}} \supset_2 \frac{\sqrt{\pi} 2^{\nu/2} \Gamma((\nu+1)/2)}{a(\nu-1)(\nu-2)} pq \\ & \exp(a(\sqrt{p}+\sqrt{q})^2) D_{2-\nu}[2\sqrt{a}(\sqrt{p}+\sqrt{q})] R(\nu) > -2. \end{aligned}$$

(g) Let

$$f(x) = x^{\nu+1} J_{\nu}(ax);$$

so that
$$\phi(p) = \frac{2^{\nu+1} \Gamma(\nu + \frac{3}{2}) a^{\nu} p^2}{\sqrt{\pi} (p^2 + a^2)^{\nu + \frac{3}{2}}}, R(\nu) > -\frac{1}{2}.$$

$$\phi(\sqrt{p}) = \frac{2^{\nu+1} \Gamma(\nu + \frac{3}{2})}{\sqrt{\pi}} \frac{a^{\nu} p}{(p + a^2)^{\nu + \frac{3}{2}}};$$

so that
$$F\left(\frac{1}{x}\right) = \frac{2^{\nu+1} a^{\nu}}{\sqrt{\pi}} x^{\nu + \frac{1}{2}} e^{-a^2 x}$$

and $n = 2$, then

$$x^{-n} f(x) = x^{\nu-1} J_{\nu}(ax);$$

so that
$$H(p) = \frac{\Gamma(2\nu)p}{(p^2 + a^2)^{(\nu/2)}} P_{\nu-1}^{-\nu} \left(\frac{p}{\sqrt{p^2 + a^2}} \right).$$

Hence from (4.1), we get

$$\begin{aligned} \frac{(xy)^{\nu-\frac{1}{2}}}{(x+y)^{\nu+1}} e^{-a^2 xy/(x+y)} \supset_2 \frac{\pi \Gamma(2\nu)}{2^{\nu-1} a^{\nu} [(\sqrt{p} + \sqrt{q})^2 + a^2]^{-(\nu/2)}} \\ P_{\nu-1}^{-\nu} \left[\frac{\sqrt{p} + \sqrt{q}}{\sqrt{(\sqrt{p} + \sqrt{q})^2 + a^2}} \right], R(\nu) > -\frac{1}{2}. \end{aligned}$$

(h) Let

$$f(x) = x^3 J_{\frac{3}{4}}^2(x);$$

so that
$$\phi(p) = \frac{5 \cdot 2^5}{\pi p^6} {}_2F_1 \left[2, \frac{7}{2}; \frac{5}{2}; -\frac{4}{p^2} \right]$$

$$\phi(\sqrt{p}) = \frac{5 \cdot 2^5}{\pi p^3} {}_2F_1 \left[2, \frac{7}{2}; \frac{5}{2}; -\frac{4}{p} \right];$$

so that
$$F\left(\frac{1}{x}\right) = \frac{5 \cdot 2^5 x^3}{\pi \Gamma(4)} {}_2F_2 \left[2, \frac{7}{2}; \frac{5}{2}, 4; -4x \right]$$

and $n = 2$, then

$$x^{-n} f(x) = x J_{\frac{3}{4}}^2(x);$$

so that
$$H(p) = \frac{2^4}{\pi \Gamma(4) p^5} {}_2F_1 \left[([2, 3; 4, -\frac{4}{p^2}] \right).$$

Hence from (4.1), we get

$$\frac{(xy)^2}{(x+y)^{\frac{7}{2}}} {}_2F_2 \left[2, \frac{7}{2}; \frac{5}{2}, 4; -\frac{4(x+y)}{xy} \right] \supset_2 \frac{4\sqrt{\pi} pq}{5(\sqrt{p}+\sqrt{q})^5} {}_2F_1 \left[2, 3; 4; -\frac{4}{(\sqrt{p}+\sqrt{q})^2} \right].$$

(i) In the theorem taking $n = 3$ and let

$$f(x) = \frac{x^\nu}{\Gamma(\nu+1)};$$

so that $\phi(p) = \frac{1}{p^\nu}$, $R(\nu) > -1$

$$x^{-n} f(x) = \frac{x^{\nu-3}}{\Gamma(\nu+1)};$$

so that $H(p) = \frac{\Gamma(\nu-2)}{\Gamma(\nu+1)p^{\nu-3}}$

$$\phi(\sqrt{p}) = p^{-(\nu/2)};$$

so that $F\left(\frac{1}{x}\right) = \frac{x^{\nu/2}}{\Gamma((\nu+2)/2)}$.

Hence from (4.1), we get

$$\frac{(xyz)^{(\nu-2)/2}}{(xy+yz+zx)^{(\nu+1)/2}} \supset_3 \frac{8\pi\Gamma((\nu+2)/2)}{\nu(\nu-1)(\nu-2)} \times \frac{pqr}{(\sqrt{p}+\sqrt{q}+\sqrt{r})^{\nu-2}}, R(\nu) > -2.$$

Finally I wish to thank Prof. S. C. Mitra for his suggestions and guidance in the preparation of this paper.

REFERENCES

ERDÉLYI, A. and others

[1] Tables of Integrals Transforms, Vol. 1, Bateman-Project.

VOELKER & DOETSCH

[2] Die Zweidimensionale Laplace-Transformation.

McLACHLAN, N. W., HUMBERT, P. et POLI, L.

[8] (1950) Memorial des sciences.

(Oblatum 17-12-64)

Deptt. of Maths.

Birla college of Engineering

Pilani (Raj.) India.