

# COMPOSITIO MATHEMATICA

ROBERT KAUFMAN

JANG-MEI WU

## **Singularity of parabolic measures**

*Compositio Mathematica*, tome 40, n° 2 (1980), p. 243-250

[http://www.numdam.org/item?id=CM\\_1980\\_\\_40\\_2\\_243\\_0](http://www.numdam.org/item?id=CM_1980__40_2_243_0)

© Foundation Compositio Mathematica, 1980, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## SINGULARITY OF PARABOLIC MEASURES\*

Robert Kaufman and Jang-Mei Wu

### Abstract

We show by example that a recent result of Dahlberg on harmonic measure for the Laplace equation can not be extended to parabolic measure for heat equation. The example is based on the non-self-adjointness of the heat operator; the methods are estimations of Green's function and construction of special boundary curves.

Let  $f(t)$  be a continuous function on  $(-\infty, \infty)$  and  $\Omega \subseteq R^2$  be the region  $\{(x, t): x > f(t)\}$ . Let  $m$  be the measure on  $\partial\Omega$  defined by  $m(E) =$  the Lebesgue measure of  $\{t: (f(t), t) \in E\}$ . If  $\Omega$  is Dirichlet regular for the heat equation (or adjoint heat equation), for a fixed point  $(y, s) \in \Omega$ , the parabolic measure (or adjoint parabolic measure) of a Borel set  $E \subseteq \partial\Omega$  at  $(y, s)$ , denoted by  $w^{(y,s)}(E)$  (or  $w^{*(y,s)}(E)$ ), is defined to be the value at  $(y, s)$  of the solution of the heat equation (or adjoint heat equation) on  $\Omega$  with boundary value 1 on  $E$  and 0 on  $\partial\Omega \setminus E$  in the Brelot-Peron-Wiener sense.

In case  $f(t) \equiv 0$ , and  $\Omega = \{x > 0\}$  it is known that  $m, w^{(x_0, t_0)}, w^{*(x_0, t_0)}$  are mutually absolutely continuous on  $\{(0, t): s_0 \leq t \leq t_0\}$ .

Let  $\lambda^{(y,s)}$  be the harmonic measure on  $\partial\Omega$  at  $(y, s)$  corresponding to the Laplace equation  $\partial^2/\partial x^2 + \partial^2/\partial t^2 = 0$ . It is known by conformal mapping that

**THEOREM A:** *If  $f$  is Lip 1, then  $\lambda^{(y,s)}$  and  $m$  are mutually absolutely continuous on  $\partial\Omega$ .*

In fact, Dahlberg [1] has proved Theorem A for  $x \in R^n, n \geq 2$ . The proof depends explicitly on the self-adjointness of the Laplace equation. Domains with Lip 1 boundaries are the most general regions on which the boundary behavior of harmonic functions has been extensively studied.

\* Research partially supported by an NSF-Grant at Illinois and an XL-Grant at Purdue.

Because the affine transformations  $\{x \rightarrow ax + b, t \rightarrow a^2t + c\}$  are the only diffeomorphisms that preserve solutions of heat equation [2], regions with  $\text{Lip} \frac{1}{2}$  boundaries are very natural for studying solutions of heat equation. In [5], Petrowski proved that if  $f(t)$  is  $\text{Lip} \frac{1}{2}$  then every point on  $\partial\Omega$  is a regular point for heat equation.

Richard A. Hunt proposed the problem whether  $m, w$  and  $w^*$  are mutually absolutely continuous on  $\partial\Omega$  if  $f(t)$  is  $\text{Lip} \frac{1}{2}$ . In [6], the first author proved the following

**THEOREM B:** *Suppose that  $f(t)$  is  $\text{Lip} \frac{1}{2}$  and  $E$  is a set on  $\partial\Omega$  with  $m$  measure zero, then  $E$  is composed of two parts, one with  $w^{(y,s)}$  measure zero, the other with  $w^{*(y,s)}$  measure zero for each  $(y, s) \in \Omega$ .*

In this note, we show by example that  $m, w$  and  $w^*$  can be mutually singular on  $\partial\Omega$ , for certain  $\text{Lip} \frac{1}{2}$  function  $f(t)$ .

### 1. Lemmas on parabolic functions

We call solutions of heat equation parabolic functions.

For fixed  $(y, s) \in R^2$ , we denote by  $W(x, t; y, s)$  the fundamental solution of heat equation defined by

$$W(x, t; y, s) = [4\pi(t - s)]^{-1/2} \exp\left[-\frac{|x - y|^2}{4(t - s)}\right] \quad \text{for } t > s$$

$$0 \quad \text{for } t \leq s.$$

Suppose  $\Omega$  is Dirichlet regular for heat equation, we say  $g$  is the Green's function for  $\Omega$ , if for each fixed  $(y, s) \in \Omega$ ,  $W(x, t; y, s) - g(x, t; y, s)$  is the bounded parabolic function in  $\Omega$  with boundary value  $W(x, t; y, s)$  for  $(x, t) \in \partial\Omega$ .

**LEMMA 1:** *Suppose  $D = \{(x, t): x > 2\sqrt{|t|}\}$ ,  $(X, -T)$  is a point in  $D$  with  $T > 0$  and  $g(x, t)$  is the Green's function on  $D$  with pole at  $(X, -T)$ . Then there are constants  $c$  and  $C$  depending on  $(X, -T)$  so that  $g(x, 0) \leq Cx^2$  for  $0 < x < c$ .*

**PROOF:** Let  $b = X - 2\sqrt{T}$  and  $S = \{(x, t); x = X - b/2 \text{ and } -T \leq t \leq 0\}$ . Let  $C_1 = \sup\{g(x, t): (x, t) \in S\}$  and

$$C_2 = \inf\left\{\frac{1}{\sqrt{8\pi T}} - W(x, t; 0, -2T): (x, t) \in S\right\}.$$

The level curve  $\gamma$  defined by  $W(x, t; 0, -2T) = 1/\sqrt{8\pi T}$  satisfies the equation:

$$\frac{1}{\sqrt{4\pi(t+2T)}} e^{-(x^2/4(t+2T))} = \frac{1}{\sqrt{8\pi T}}$$

or

$$x^2 = -2(t+2T) \log\left(1 + \frac{t}{2T}\right).$$

If  $(x, t) \in \gamma$  and  $-T \leq t < 0$  then  $x^2 < -4t$ . Therefore if  $(x, t) \in D$  and  $t \geq -T$ , then  $(1/\sqrt{8\pi T}) - W(x, t; 0, -2T) \geq 0$ . Let  $V$  be the region  $D \cap \{(x, t): t > -T\} \cap \{x < X - b/2\}$ . We observe by the definitions of  $C_1$  and  $C_2$  that

$$C_2 g(x, t) \leq C_1 \left( \frac{1}{\sqrt{8\pi T}} - W(x, t; 0, -2T) \right)$$

for  $(x, t) \in S$ . When  $(x, t)$  is on the part of  $\partial V$  with  $t = -T$  or on the part of  $\partial V$  with  $x = 2\sqrt{|t|}$ , the above inequality also holds because the left side is zero and the right side is positive. In view of the maximum principle for the solutions of heat equations [3, Chap. 2], we obtain

$$\begin{aligned} g(x, 0) &\leq \frac{C_1}{C_2} \left[ \frac{1}{\sqrt{8\pi T}} - W(x, 0; 0, -2T) \right] \\ &= \frac{C_1}{C_2} \frac{1}{\sqrt{8\pi T}} (1 - e^{-x^2/8T}) \\ &\leq Cx^2 \end{aligned}$$

for  $(x, t) \in S$ . When  $(x, t)$  is on the part of  $\partial V$  with  $t = -T$  or on the part of  $\partial V$  with  $x = 2\sqrt{|t|}$ , the above inequality also holds because the

Suppose  $f$  is  $\text{Lip } \frac{1}{2}$  satisfying  $|f(t) - f(\tau)| \leq M|t - \tau|^{1/2}$ . For  $a > 0$  we denote by  $\Delta(t, a) = \{(f(s), s): |s - t| \leq a\}$  and  $A(t, a) = (f(t) + 10M\sqrt{a}, t + 2a)$ . Under these assumptions, we may reformulate Lemma 1.4 in [4] and Lemma 2.2 in [6] as follows.

**LEMMA 2:** *There exist positive constants  $C, c$  depending on  $M$  only, so that*

$$w^{(y,s)}(\Delta(t, r)) \leq Cw^{(y,s)}(\Delta(t, a))w^{A(t,a)}(\Delta(t, r))$$

whenever  $0 < r < a/2$ ,  $(y, s) \in \Omega$  and  $|y - f(t)|^2 + |s - t| > ca$ .

LEMMA 3: Let  $(x_0, t_0), (y_0, s_0)$  be two fixed points in  $\Omega$  with  $t_0 > s_0 > a > 0$ , and  $g$  be the Green's function on  $\Omega$ . Then there are positive constants  $C, \mu, \rho$  depending on  $a, M, (x_0, t_0)$  and  $(y_0, s_0)$  so that

$$w^{(y_0, s_0)}(\Delta(t, r)) \leq Cr^{1/2}g(x_0, t_0; f(t) + \mu\sqrt{r}, t)$$

whenever  $-a < t < a$  and  $0 < r < \rho$ .

### 2. A test for singular measures

Suppose that  $\mu$  is a positive Borel measure on  $[0, 1]$ , and that  $\mu$  is not totally singular to Lebesgue measure  $m$ ; then  $d\mu/dx \geq c > 0$  on some set  $E$  of measure  $m(E) > 0$ . Thus

$$\begin{aligned} \lim \mu([x, x + h])h^{-1} &= \mu'(x) \text{ and} \\ \lim \mu([x - h, x])^{-1} &= \mu'(x) \end{aligned}$$

as  $h \rightarrow 0^+$ , when  $x \in E$ . Letting  $h = 1, \frac{1}{2}, \frac{1}{3}, \dots$  we can apply Egoroff's theorem to find a set  $E_0 \subseteq E$ , with  $m(E_0) > 0$ , and a sequence  $\epsilon_n > 0$  decreasing to 0, so that

$$\begin{aligned} |\mu([x, x + h]) - h\mu'(x)| &\leq \epsilon_n h, \text{ and} \\ |\mu([x - h, x]) - h\mu'(x)| &\leq \epsilon_n h, \end{aligned}$$

when  $(n + 1)^{-1} \leq h \leq n^{-1}$ , and  $x \in E_0$ . Let now  $r(n)$  be a polygonal function on  $(0, 1]$ , defined by the conditions  $r(n^{-1}) = \epsilon_{n-1}$  for  $n \geq 2$  and  $r$  is constant on  $[\frac{1}{2}, 1]$ . Then

$$|\mu([a, b]) - (b - a)\mu'(x)| \leq (b - a)r(b - a)$$

whenever  $0 \leq a < x < b \leq 1$  and  $x \in E_0$ .

Let  $0 < \delta < \frac{1}{4}$  and observe that by the Lebesgue density theorem,  $E_0$  must meet one of the sets  $[(k + \frac{5}{8})N^{-1}, (k + \frac{3}{4})N^{-1}]$  ( $1 \leq k \leq N - 2$ ) whenever  $N$  is sufficiently large. We apply the inequality on  $\mu$ -measures to intervals  $[a, b_1], [a, b_2], [a, b_3]$  with

$$\begin{aligned} a &= (k + \frac{1}{2})N^{-1}, \quad b_1 = (k + 1 - \delta)N^{-1}, \quad b_2 = (k + 1 + \delta)N^{-1}, \\ &\quad b_3 = (k + 1 + \frac{1}{2})N^{-1}. \end{aligned}$$

We write these inequalities as

$$E_i: |\mu([a, b_i]) - (b_i - a)\mu'(x)| \leq N^{-1}r(N^{-1}), \quad i = 1, 2, 3.$$

Combination of  $E_1$  and  $E_2$  gives

$$|\mu((b_1, b_2]) - (b_2 - b_1)\mu'(x)| \leq 2N^{-1}r(N^{-1}),$$

and comparison with  $E_3$  yields

$$|\mu((b_1, b_2]) - 2\delta\mu([a, b_3])| \leq 3N^{-1}r(N^{-1}).$$

Let us write  $I = [(k + \frac{1}{2})N^{-1}, (k + 1 + \frac{1}{2})N^{-1}]$ ,  $I_\delta = ((k + 1 - \delta)N^{-1}, (k + 1 + \delta)N^{-1}]$ , that is  $I = [a, b_3]$ ,  $I_\delta = (b_1, b_2]$ . We divide the last inequality by  $\mu(I)$ , which exceeds  $c/(2N)$  for  $N \geq N_0$ . We obtain

$$\mu(I_\delta)/\mu(I) \geq 2\delta + o(1), \quad N \rightarrow +\infty.$$

### 3. Construction of curves

Let  $h(t)$  be a function on  $[0, 1]$  subject to the following conditions

- 1)  $0 \leq h \leq 1$ ,
- 2)  $h(t) = 4t^{1/2}$  for  $0 \leq t \leq \frac{1}{3}$ ,
- 3)  $h(t) = h(1 - t)$ ,
- 4)  $h$  is of class  $C^1$  on  $[\frac{1}{4}, \frac{3}{4}]$ ,
- 5)  $|h(t) - h(s)| \leq 4|t - s|^{1/2}$  for  $0 \leq s \leq t \leq 1$ .

Let  $h_n(t)$  be the function on  $[0, 1]$  with period  $1/n$ , such that  $h_n(t) = h(nt)/n^{1/2}$  for  $0 \leq t \leq 1/n$ .

Let  $\ell_n(t)$  be a function of class  $C^1[0, 1]$ , periodic with period  $1/n$ , such that

- 6)  $0 \leq \ell_n \leq h_n$ ,
- 7)  $\ell_n(0) = 0$  and  $\ell_n(t) = h_n(t)$  for  $n^{-3} \leq t \leq n^{-1} - n^{-3}$ ,
- 8)  $|\ell_n(t) - \ell_n(s)| \leq 5|t - s|^{1/2}$ .

We shall choose a sequence  $(n_j)$  and set  $f_k(t) = \sum_1^k \ell_{n_j}(t)$ ,  $f(t) = \sum_1^\infty \ell_{n_j}(t)$ . We require the following properties of  $f$  and  $f_k$ :

- 9)  $|f(t) - f(s)| \leq 8|t - s|^{1/2}$ .
- 10)  $0 \leq f(t) - f_k(t) \leq n_k^{-3/2}$ .

11) The inequalities  $3|t - \tau|^{1/2} \leq f_k(t) - f_k(\tau) \leq 6|t - \tau|^{1/2}$  hold whenever  $\tau = i/n_k$  ( $1 \leq i \leq n_k - 1$ ) and  $n_k^{-3} \leq |t - \tau| \leq (4n_k)^{-1}$ .

To obtain 10) we observe the inequality  $0 \leq f(t) - f_k(t) \leq \sum_{k+1}^\infty n_j^{-1/2}$ , and simply choose  $n_{j+1} > 16n_j^3$ . This choice is compatible with the rest of the construction and we don't mention it again.

To obtain 11) and 9) we let  $B_{k-1}$  be an upper bound for  $|f'_{k-1}|$ , so that

$$\begin{aligned} |f_k(t) - f_k(\tau) - \ell_{n_k}(t) + \ell_{n_k}(\tau)| &\leq B_{k-1}|t - \tau| \\ &\leq B_{k-1}(4n_k)^{-1/2}|t - \tau|^{1/2}, \end{aligned}$$

for the numbers  $\tau, t$  mentioned in 11). By 7) and 2) we find

$$\ell_{n_k}(t) - \ell_{n_k}(\tau) = 4|t - \tau|^{1/2}$$

for these numbers, because  $n_k\tau$  is an integer, and we obtain 11) by taking  $B_{k-1}(4n_k)^{-1/2} < 1$ . To obtain 9) we suppose that  $|f_p(t) - f_p(s)| \leq (6 - p^{-1})|t - s|^{1/2}$ , for  $p = k - 1$ . (This is true when  $p = 1$ ). Then

$$|f_k(t) - f_k(s)| \leq |f_{k-1}(t) - f_{k-1}(s)| + |\ell_{n_k}(t) - \ell_{n_k}(s)|.$$

Since  $\ell_{n_k} \leq n_k^{-1/2}$ , we have the inequality

$$|f_k(t) - f_k(s)| \leq (6 - (k - 1)^{-1})|t - s|^{1/2} + 2n_k^{-1/2}.$$

Thus the required estimate is valid when  $(k^2 - k)^{-1}|t - s|^{1/2} \geq 2n_k^{-1/2}$  or  $|t - s| \geq 4n_k^{-1}(k^2 - k)^2$ . But when the last inequality is violated, we can use the estimation

$$|f_k(t) - f_k(s)| \leq B_{k-1}|t - s| + 5|t - s|^{1/2};$$

this yields the inequality in question provided  $B_{k-1}|t - s| \leq |t - s|^{1/2} \cdot \frac{1}{2}$ , or  $|t - s| \leq (2B_{k-1})^{-2}$ . One estimate or the other is available for large  $n_k$ .

#### 4. The theorem

We retain the notations from §3 and extend the function  $f$ , constructed in §3, to  $(-\infty, \infty)$  by defining  $f(t) = f(0)$  for  $t < 0$  and  $f(t) = f(1)$  for  $t > 1$ . We let  $\Omega$  be  $\{(x, t): x > f(t)\}$ ,  $w$  be the parabolic measure on  $\partial\Omega$  evaluated at  $(X, T)$  and  $w^*$  be the adjoint parabolic measure on  $\partial\Omega$  evaluated at  $(Y, S)$  where  $(X, T)$  and  $(Y, S)$  are two fixed points in  $\Omega$  with  $T > 1$  and  $S < 0$ .

We observe, with the aid of maximum principle, that for  $E \subseteq \{(f(t), t): 0 < t < 1\}$ , if  $w(E) = 0$  then  $w^{(x,t)}(E) = 0$  for every  $(x, t) \in \Omega$ ; and if  $w^*(E) = 0$  then  $w^{*(x,t)}(E) = 0$  for every  $(x, t) \in \Omega$ .

**THEOREM:** *None of the three measures:  $m$ ,  $w$  and  $w^*$  on  $\partial\Omega$  is absolutely continuous with respect to another. In fact  $m$ ,  $w$  and  $w^*$  are totally singular with respect to each other on  $\{(f(t), t): 0 < t < 1\}$ .*

We first prove the following lemma and assume as we may that  $(X, T) = (10, 100)$ .

LEMMA 4: *There are positive absolute constants  $C$  and  $\rho < \frac{1}{32}$ , so that whenever  $\tau = i/n_k$  ( $1 \leq i \leq n_k - 1$ ),  $n_k^{-1} < \delta < \rho$ ,*

$$I_k = \{(f(t), t): |t - \tau| < (16n_k)^{-1}\} \text{ and}$$

$$E_k = \{(f(t), t): |t - \tau| < \delta n_k^{-1}\}$$

then  $w(E_k) \leq C\delta^{3/2}w(I_k)$  for sufficiently large  $k$ .

PROOF: For a fixed  $\tau = i/n_k$ , we let  $A = (f(\tau) + 5/\sqrt{n_k}, \tau + 1/8n_k)$  and  $B$  be  $A + (0, 1/8n_k)$ . From 9) and Lemma 2 it follows that for some absolute constant  $C$ ,

$$w(E_k) \leq Cw(I_k)w^A(E_k).$$

Let  $\Phi$  be the map  $(x, t) \rightarrow (\sqrt{n_k}(x - f(\tau)), n_k(t - \tau))$  and  $G$  be the Green's function on  $\Phi(\Omega)$ . We note that  $\partial\Phi(\Omega) = \Phi(\partial\Omega)$  is the graph of a  $\text{Lip } \frac{1}{2}$  function with 8 as an upper bound for the  $\text{Lip } \frac{1}{2}$  constant and  $\Phi$  preserves parabolic functions (i.e.  $v$  is parabolic on  $\Phi(\Omega)$  if and only if  $v(\Phi)$  is parabolic on  $\Omega$ ). Let  $\bar{w}$  be the parabolic measure on  $\partial\Phi(\Omega)$ , thus  $\bar{w}^{\Phi(A)}(\Phi(E_k)) = w^A(E_k)$ . Because  $\Phi(A) = (5, \frac{1}{8})$  and  $\Phi(B) = (5, \frac{1}{4})$ , it follows from Lemma 3 that there exist absolute constants  $C$ ,  $\mu$  and  $\rho < \frac{1}{32}$ , so that

$$w^A(E_k) = \bar{w}^{\Phi(A)}(\Phi(E_k))$$

$$\leq C\sqrt{\delta}G(\Phi(B); \mu\sqrt{\delta}, 0)$$

if  $0 < \delta < \rho$ .

Let  $\alpha(t) = 2|t - \tau|^{1/2} + f(\tau) - 10n_k^{-3/2}$ . From 9), 10) and 11) it follows that  $\alpha(t) \leq f(t)$  whenever  $|t - \tau| < (4n_k)^{-1}$ . Therefore  $\Phi(\Omega) \cap \{(x, t): |t| < \frac{1}{4}\} \subseteq \{(x, t): x > 2|t|^{1/2} - 10n_k^{-1}\}$ . let  $\tilde{G}$  be the Green's function on  $\{(x, t): x > 2|t|^{1/2} - 10n_k^{-1}\}$ . We recall that  $\Phi(B) = (5, \frac{1}{4})$  and  $n_k^{-1} < \delta < \rho$ , and obtain, by the maximum principle and the adjoint form of Lemma 1, that

$$G(\Phi(B); \mu\sqrt{\delta}, 0)$$

$$\leq \tilde{G}(\Phi(B); \mu\sqrt{\delta}, 0) = \tilde{G}^*(\mu\sqrt{\delta}, 0; \Phi(B))$$

$$\leq C(\mu\sqrt{\delta} + 10n_k^{-1})^2 \leq C\delta,$$

for absolute constants  $C$ . Thus  $w^A(E_k) \leq C\delta^{3/2}$ . This proves Lemma 4.

PROOF OF THE THEOREM: From the above lemma and the test for singular measures in §2 we see that  $w$  is totally singular to  $m$  on

$\{(f(t), t): 0 < t < 1\} \equiv S$ , that is, there is a set  $E \subseteq S$  of  $m$  measure zero but  $w(E) = w(S)$ . Similarly there is a set  $E^* \subseteq S$  of  $m$  measure zero but  $w^*(E^*) = w^*(S)$ . From these properties and Theorem B, we conclude that  $w$  and  $w^*$  are mutually singular. The theorem follows easily.

## REFERENCES

- [1] B.E.J. DAHLBERG: On estimates of harmonic measures. *Arch. Rational Mech. Anal.* 65, No. 3 (1977) 275–288.
- [2] E.G. EFFROS and J.L. KAZDAN: On the Dirchlet problem for the heat equation. *Indiana Univ. Math. J.* 20 (1971) 683–693.
- [3] A. FRIEDMAN: *Partial differential equations of parabolic type*. Prentice-Hall, 1964.
- [4] J.T. KEMPER: Temperatures in several variables: kernel functions, representations and parabolic boundary values. *Trans. Amer. Math. Soc.*, 167 (1972) 243–262.
- [5] I.G. PETROWSKI: Zur Ersten Randwertaufgaben der Wärmeleitungsgleichung. *Compositio Math.* 1 (1935) 383–419.
- [6] J.-M. WU: On parabolic measures and subparabolic functions. *Trans. Amer. Math. Soc.* 251 (1979) 171–186.

(Oblatum 27-IV-1978 & 23-X-1978)

University of Illinois  
Urbana, Illinois