

RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

L. CARLITZ

Saalschützian transforms

Rendiconti del Seminario Matematico della Università di Padova,
tome 50 (1973), p. 95-118

http://www.numdam.org/item?id=RSMUP_1973__50__95_0

© Rendiconti del Seminario Matematico della Università di Padova, 1973, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*
<http://www.numdam.org/>

Saalschützian Transforms.

L. CARLITZ (*)

1. Put

$$(a)_n = a(a+1), \dots, (a+n-1), \quad (a)_0 = 1.$$

Saalschütz's theorem [1, p. 9], [3, p, 48] reads

$$(1.1) \quad \sum_{k=0}^n \frac{(-n)_k (a)_k (b)_k}{k! (c)_k (d)_k} = \frac{(c-a)_n (c-b)}{(c)_n (c-a-b)_n},$$

where

$$(1.2) \quad c + d = a + b - n + 1.$$

If we replace b by $c-b$, (1.1) becomes

$$\sum_{k=0}^n \frac{(-n)_k (a)_k (c-b)_k}{k! (c)_k (a-b-n+1)_k} = \frac{(c-a)_n (b)_n}{(c)_n (b-a)_n}.$$

Since

$$(a-b-n+1)_k = (-1)^k (b-a+n-k)_k,$$

$$\frac{(b-a)_n}{(a-b+n-1)_k} = (-1)^k (b-a)_{n-k},$$

we get

$$(1.3) \quad \frac{1}{(b)_n} \sum_{k=0}^n \binom{n}{k} (a)_k (b-a)_{n-k} \frac{(c-b)_k}{(c)_k} = \frac{(c-a)_n}{(c)_n}.$$

(*) Indirizzo dell'A.: Dept. of Mathematics, Duke University - Durham. N.C. 27706, U.S.A.

Supported in part by NSG grant GP-17031.

We may think of (1.3) as a linear transformation from the set of rational functions

$$\frac{(c-b)_n}{(c)_n} \quad (n = 0, 1, 2, \dots)$$

to the set

$$\frac{(c-a)_n}{(c)_n} \quad (n = 0, 1, 2, \dots).$$

We accordingly define the linear transformation

$$(1.4) \quad T_{a,b}: x_n = \frac{1}{(b)_n} \sum_{k=0}^n \binom{n}{k} (a)_k (b-a)_{n-k} y_k \quad (n = 0, 1, 2, \dots).$$

We shall assume that neither a nor b is equal to zero or a negative integer.

It follows at once from (1.4) that

$$(1.5) \quad T_{a,a} = I,$$

the identity transformation. We shall show that

$$(1.6) \quad T_{a,b} T_{b,a} = I$$

and that

$$(1.7) \quad T_{a,b} T_{c,d} = T_{c,d} T_{a,b}.$$

Moreover a relation of the form

$$(1.8) \quad T_{a_1, b_1} T_{a_2, b_2} \dots T_{a_n, b_n} = I$$

holds only in a « trivial » way.

In the second part of the paper we show that analogous results are implied by the q -analog of Saalschütz's theorem:

$$(1.9) \quad \sum_{k=0}^n \frac{(q^{-n})_k (a)_k (b)_k}{(q)_k (c)_k (d)_k} q^k = \frac{(c/a)_n (c/b)_n}{(c)_n (c/ab)_n},$$

where now

$$(a)_n = (1-a)(1-qa) \dots (1-q^{n-1}a), \quad (a)_0 = 1$$

and

$$(1.10) \quad cd = q^{1-n}ab.$$

2. Returning to the definition (1.4), we ask when is

$$(2.1) \quad T_{a,b} = T_{c,a}?$$

Clearly this requires

$$\frac{1}{(b)_n} (a)_k (b-a)_{n-k} = \frac{1}{(d)_n} (c)_k (d-c)_{n-k} \quad (n = 0, 1, 2, \dots; k = 0, 1, \dots, n).$$

For $k = n$ this becomes

$$\frac{(a)_n}{(b)_n} = \frac{(c)_n}{(d)_n} \quad (n = 0, 1, 2, \dots).$$

It follows that

$$\frac{a}{b} = \frac{c}{d}, \quad \frac{a+1}{b+1} = \frac{c+1}{d+1}, \quad \frac{a+2}{b+2} = \frac{c+2}{d+2}, \dots$$

and therefore

$$(a+x)(d+x) \equiv (b+x)(c+x).$$

For $x = -a$, this implies $b = a$ or $c = a$. If $b = a$ we get $T_{c,a} = I$ so that $c = d$. If $c = a$ it follows that $b = d$. Hence (2.1) holds if and only if

$$(i) \quad a = b, \quad c = d$$

or

$$(ii) \quad a = c, \quad b = d.$$

We show now that

$$(2.2) \quad T_{a,b} T_{b,c} = T_{a,c}.$$

PROOF. Put

$$(2.3) \quad y_k = \frac{1}{(c)_k} \sum_{j=0}^k \binom{k}{j} (b)_j (c-b)_{k-j} z_j \quad (k = 0, 1, 2, \dots).$$

Then by (1.4) and (2.3)

$$(2.4) \quad x_n = \frac{1}{(b)_n} \sum_{j=0}^n \binom{n}{k} (a)_k (b-a)_{n-k} \cdot \frac{1}{(c)_k} \sum_{j=0}^k \binom{k}{j} (b)_j (c-b)_{k-j} z_j = \\ = \frac{1}{(b)_n} \sum_{j=0}^n \binom{n}{j} (b)_j z_j \sum_{k=j}^n \binom{n-j}{k-j} (a)_k (b-a)_{n-k} \frac{(c-b)_{k-j}}{(c)_k}.$$

The inner sum is equal to

$$\sum_{k=0}^{n-j} \binom{n-j}{k} a_{j+k} (b-a)_{n-j-k} \frac{(c-b)_k}{(c)_{j+k}} = \\ = \frac{(a)_j}{(c)_j} \sum_{k=0}^{n-j} \binom{n-j}{k} \frac{(a+j)_k (c-b)_k}{(c+j)_k} (b-a)_{n-j-k} = \\ = \frac{(a)_j (b-a)_{n-j}}{(c)_j} \sum_{k=0}^n \frac{(-n+j)_k (a+j)_k (c-b)_k}{k! (c+j)_k (a-b-n+j+1)_k} = \\ = \frac{(a)_j (b-a)_{n-j}}{(c)_j} \frac{(c-a)_{n-j} (b+j)_{n-j}}{(c+j)_{n-j} (b-a)_{n-j}} = \frac{(a)_j (c-a)_{n-j} (b+j)_{n-j}}{(c)_n}.$$

Hence (2.4) becomes

$$x_n = \frac{1}{b_n} \sum_{j=0}^n \binom{n}{j} \frac{(a)_j (b)_j (c-a)_{n-j} (b+j)_{n-j}}{(c)_n} z_j = \frac{1}{(c)_n} \sum_{j=0}^n \binom{n}{j} (a)_j (c-a)_{n-j} z_j.$$

This evidently proves (2.2).

As an immediate corollary of (2.2) we have

$$(2.5) \quad T_{a,b} T_{b,a} = I.$$

We show next that

$$(2.6) \quad T_{b,c} T_{a,b} = T_{a,c}.$$

PROOF. Put

$$x_n = \frac{1}{(c)_n} \sum_{k=0}^n \binom{n}{k} (b)_k (c-b)_{n-k} y_k, \\ y_k = \frac{1}{(b)_k} \sum_{j=0}^k \binom{k}{j} (a)_j (b-a)_{k-j} z_j.$$

Then

$$\begin{aligned} x_n &= \frac{1}{(c)_n} \sum_{k=0}^n \binom{n}{k} (b)_k (c-b)_{n-k} \cdot \frac{1}{(b)_k} \sum_{j=0}^k \binom{k}{j} (a)_j (b-a)_{k-j} z_j = \\ &= \frac{1}{(c)_n} \sum_{j=0}^n \binom{n}{j} (a)_j z_j \sum_{k=j}^n \binom{n-j}{k-j} (c-b)_{n-k} (b-a)_{k-j}. \end{aligned}$$

The inner sum is equal to

$$\begin{aligned} \sum_{k=0}^{n-j} \binom{n-j}{k} (c-b)_{n-j-k} (b-a)_k &= (c-b)_{n-j} \sum_{k=0}^{n-j} \frac{(-n+j)_k (b-a)_k}{k! (b-c-n+j+1)_k} = \\ &= (c-b)_{n-j} \frac{(a-c-n+j+1)_{n-j}}{(b-c-n+j+1)_{n-j}} = (c-b)_{n-j} \frac{(c-a)_{n-j}}{(c-b)_{n-j}} = (c-a)_{n-j} \end{aligned}$$

by Vandermonde's theorem. Thus

$$x_n = \frac{1}{(c)_n} \sum_{j=0}^n \binom{n}{j} (a)_j (c-a)_{n-j} z_j,$$

which proves (2.6).

It is now easy to prove that

$$(2.7) \quad T_{a,b} T_{c,d} = T_{c,d} T_{a,b}.$$

Indeed, by (2.2) and (2.7),

$$\begin{aligned} T_{a,b} T_{c,d} &= T_{c,b} T_{a,c} T_{c,d} \\ &= T_{c,b} T_{a,d} \\ &= T_{c,b} T_{b,d} T_{a,b} \\ &= T_{c,d} T_{a,b}. \end{aligned}$$

We shall however give a second proof of (2.7) that makes use of an explicit formula for the transformation $T_{a,b} T_{c,d}$. Put

$$\begin{aligned} x_n &= \frac{1}{(b)_n} \sum_{k=0}^n \binom{n}{k} (a)_k (b-a)_{n-k} y_k, \\ y_k &= \frac{1}{(d)_k} \sum_{j=0}^k \binom{k}{j} (c)_j (d-c)_{k-j} z_j. \end{aligned}$$

Then

$$\begin{aligned}
 (2.8) \quad x_n &= \frac{1}{(b)_n} \sum_{k=0}^n \binom{n}{k} \frac{(a)_k (b-a)_{n-k}}{(d)_k} \sum_{j=0}^k \binom{k}{j} (c)_j (d-c)_{k-j} z_j = \\
 &= \frac{1}{(b)_n} \sum_{j=0}^n \binom{n}{j} (c)_j z_j \sum_{k=j}^n \binom{n-j}{k-j} (a)_k (b-a)_{n-k} \frac{(d-c)_{k-j}}{(d)_k} = \\
 &= \frac{1}{(b)_n} \sum_{j=0}^n \binom{n}{j} (c)_j z_j \sum_{k=0}^{n-j} \binom{n-j}{k} (a)_{j+k} (b-a)_{n-j-k} \frac{(d-c)_k}{(d)_{j+k}} = \\
 &= \frac{1}{(b)_n} \sum_{j=0}^n \binom{n}{j} \frac{(a)_j (c)_j}{(d)_j} z_j \sum_{k=0}^{n-j} \binom{n-j}{k} (a+j)_k (b-a)_{n-j-k} \frac{(d-c)_k}{(d+j)_k}.
 \end{aligned}$$

By Vandermode's theorem

$$\frac{(d-c)_k}{(d+j)_k} = \sum_{s=0}^k \frac{(-k)_s (c+j)_s}{s! (d+j)_s} = \sum_{s=0}^k (-1)^s \binom{k}{s} \frac{(c+j)_s}{(d+j)_s}.$$

Thus

$$\begin{aligned}
 &\sum_{k=0}^{n-j} \binom{n-j}{k} (a+j)_k (b-a)_{n-j-k} \frac{(d-c)_k}{(d+j)_k} = \\
 &= \sum_{k=0}^{n-j} \binom{n-j}{k} (a+j)_k (b-a)_{n-j-k} \sum_{s=0}^k (-1)^s \binom{k}{s} \frac{(c+j)_s}{(d+j)_s} = \\
 &= \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \frac{(c+j)_s}{(d+j)_s} \sum_{k=s}^{n-j} \binom{n-j-s}{k-s} (a+j)_k (b-a)_{n-j-k} = \\
 &= \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \frac{(c+j)_s}{(d+j)_s} \sum_{k=0}^{n-j-s} \binom{n-j-s}{k} (a+j)_{s+k} (b-a)_{n-j-s-k} = \\
 &= \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \frac{(a+j)_s (c+j)_s}{(d+j)_s} (b-a)_{n-j-s} \cdot \\
 &\cdot \sum_{k=0}^{n-j-s} \frac{(-n+j+s)_k (a+j+s)_k}{k! (a-b-n+j+s+1)_k} = \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \cdot \\
 &\cdot \frac{(a-j)_s (c+j)_s}{(d+j)_n} (b-a)_{n-j-s} \frac{(-b-n+1)_{n-j-s}}{(a-b-n+j+s+1)_{n-j-s}} = \\
 &= \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \frac{(a+j)_s (c+j)_s}{(d+j)_s} (b-a)_{n-j-s} \frac{(b+j+s)_n}{(b-a)_{n-s}} = \\
 &= (b)_n \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \frac{(a+j)_s (c+j)_s}{(b)_{j+s}}.
 \end{aligned}$$

Therefore (2.8) becomes

$$(2.9) \quad x_n = \sum_{j=0}^n \binom{n}{j} \frac{(a)_j(c)_j}{(b)_j(d)_j} z_j \sum_{s=0}^{n-j} (-1)^s \binom{n-j}{s} \frac{(a+j)_s(c+j)_s}{(b+j)_s(d+j)_s}$$

or, if we prefer,

$$(2.10) \quad x_n = \sum_{k=0}^n \binom{n}{k} \frac{(a)_k(c)_k}{(b)_k(d)_k} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} z_j.$$

Clearly (2.9) represents both $T_{a,b}T_{c,d}$ and $T_{c,d}T_{a,b}$. For that matter it also represents both $T_{a,d}T_{b,c}$ and $T_{b,c}T_{a,d}$. In particular, (2.10) implies (2.7).

3. It is natural to ask whether

$$(3.1) \quad T_{a,b}T_{c,d} = T_{e,f},$$

for properly chosen e, f . By (2.8) this is equivalent to

$$(3.2) \quad \frac{(a)_j(c)_j}{(b)_n(d)_j} \sum_{k=0}^{n-j} \binom{n-j}{k} (a+j)_k(b-a)_{n-j-k} \frac{(d-c)_k}{(d+j)_k} = \frac{1}{(f)_n} (e)_j(f)_{n-j}.$$

For $j = n$, (3.2) reduces to

$$(3.3) \quad \frac{(a)_n(c)_n}{(b)_n(d)_j} = \frac{(e)_n}{(f)_n}.$$

This implies

$$\frac{ac}{bd} = \frac{e}{f}, \quad \frac{(a+1)(c+1)}{(b+1)(d+1)} = \frac{e+1}{f+1}, \quad \frac{(a+2)(c+2)}{(b+2)(d+2)} = \frac{e+2}{f+2}, \dots$$

and therefore

$$(3.4) \quad (a+x)(c+x)(f+x) \equiv (b+x)(d+x)(e+x).$$

For $x = -0$, (3.4) becomes

$$(b-a)(d-a)(e-a) = 0.$$

(i) If $b = a$, (3.1) reduces to $T_{c,d} = T_{e,f}$, so that

$$c = d, \quad e = f \quad \text{or} \quad c = e, \quad d = f;$$

(ii) if $d = a$, (3.1) becomes $T_{c,b} = T_{e,f}$, so that

$$c = b, \quad e = f \quad \text{or} \quad c = e, \quad b = f;$$

(iii) if $e = a$, (3.1) reduces to $T_{c,d} = T_{b,f}$, so that

$$c = d, \quad b = f \quad \text{or} \quad c = b, \quad d = f.$$

Thus in every case, (3.1) holds only in a « trivial » way, that is, as a consequence of (2.2) or (2.6).

Consider next the equation

$$(3.5) \quad T_{a,b} T_{c,d} = T_{e,f} T_{g,h}.$$

By (2.10), (3.5) is equivalent to

$$(3.6) \quad \frac{(a)_k (c)_k}{(b)_k (d)_k} = \frac{(e)_k (g)_k}{(f)_k (h)_k}.$$

Exactly as above, (3.6) implies

$$(3.7) \quad (a+x)(c+x)(f+x)(h+x) = (b+x)(d+x)(e+x)(g+x).$$

Taking $x = -a$, it follows that $a = b, d, e$ or g . If $a = b$, (3.5) reduces to

$$T_{c,d} = T_{e,f} T_{g,h};$$

if $a = d$, it reduces to

$$T_{d,b} = T_{e,f} T_{g,h};$$

if $a = e$, we get

$$T_{c,d} = T_{b,f} T_{g,h};$$

if $a = g$, we get

$$T_{c,d} = T_{e,f} T_{b,h}.$$

Thus in every case, (3.5) reduces to (3.1).

The most general relation can be reduced to the form

$$(3.8) \quad T_{a_1, b_1} T_{a_2, b_2} \dots T_{a_n, b_n} = I.$$

As above, (3.8) implies

$$(3.9) \quad (a_1 + x) \dots (a_n + x) \equiv (b_1 + x) \dots (b_n + x).$$

It follows from (3.10) that $a_n = b_i$ for some i . If $i = n$, (3.8) reduces to

$$T_{a_1, b_1} \dots T_{a_{n-1}, b_{n-1}} = I .$$

If $i \neq n$, we may assume $i = n - 1$, in which case (3.8) becomes

$$T_{a_1, b_1} \dots T_{a_{n-1}, b_{n-1}} T_{a_n, b_n} = I .$$

Again the number of factors on the left has been reduced.

We conclude that (3.8) holds only in a « trivial » way.

4. We may write (1.3) in the following form.

$$(4.1) \quad (x - a)_n = \frac{1}{(b)_n} \sum_{k=0}^n \binom{n}{k} (a)_k (b - a)_{n-k} (x - b)_k (x + k)_{n-k} .$$

This suggests the following interpolation problem. Let $f(x)$ be a polynomial of degree $\leq n$. We consider the representation of $f(x)$ in the form

$$(4.2) \quad f(x) = \sum_{k=0}^n A_{n,k} (x - a)_k (x + k)_{n-k} ,$$

where a is an arbitrary constant and $A_{n,k} = A_{n,k}(a)$ is independent of x .

To determine the coefficients $A_{n,k}$ we first take $x = a$. This gives

$$(4.3) \quad (a)_n A_{n,0} = f(a) .$$

Next, for $x = a - 1$, (4.2) reduces to

$$f(a - 1) = (a - 1)_n A_{n,0} - (a)_{n-1} A_{n,1} ,$$

so that

$$(4.4) \quad (a)_n A_{n,1} = (a - 1)f(a) - (a + n - 1)f(a - 1) .$$

For $x = a - 2$, (4.2) becomes

$$f(a - 2) = (a - 2)_n A_{n,0} - 2(a - 1)_n A_{n,1} + 2(a)_{n-2} A_{n,2} .$$

This yields

$$(4.5) \quad \begin{aligned} 2(a)_{n,2} = & a(a - 1)f(a) - 2(a - 1)(a + n - 1)f(a - 1) + \\ & + (a + n - 1)(a + n - 2)f(a - 2) . \end{aligned}$$

At the next step we get

$$(4.6) \quad 6(a)_n A_{n,3} = a(a-1)(a+1)f(a) - 3a(a-1)a(a+n-1)f(a-1) + \\ + 3(a-1)(a+n-1)(a+n-2)f(a-2) - \\ - (a+n-1)(a+n-2)(a+n-3)f(a-3).$$

This suggests the general result

$$(4.7) \quad k!(a)_n A_{n,k} = \sum_{j=0}^k (-1)^j \binom{k}{j} (a-1)_{k-j} (a+n-j)_j f(a-j).$$

To prove (4.7) we note first that

$$(4.8) \quad 1 = \frac{1}{(a)_k} \sum_{j=0}^k (-1)^j \binom{k}{j} (x-a)_j (x+j)_{k-j}.$$

Indeed

$$\frac{1}{(x)_k} \sum_{j=0}^k (-1)^j \binom{k}{j} (x-a)_j (x+j)_{k-j} = \\ = \sum_{j=0}^k (-1)^j \binom{k}{j} \frac{(x-a)_j}{(x)_j} = \sum_{j=0}^k \frac{(-k)_j (x-a)_j}{j!(a)_j} = \frac{(a)_k}{(x)_k},$$

by Vandermonde's theorem. This evidently proves (4.8).

Now replace x by $x+m$ in (4.8) and then multiply both sides by $(x-a)_m$. We get

$$(x-a)_m = \frac{1}{(a)_k} \sum_{j=0}^k (-1)^j \binom{k}{j} (x-a)_{m+j} (x+m+j)_{k-j} = \\ = \frac{1}{(a)_k} \sum_{j=m}^{k+m} (-1)^{j-m} \binom{k}{j-m} (x-a)_j (x+j)_{k+m-j}.$$

Replacing k by $n-m$, this becomes

$$(4.9) \quad (x-a)_m = \frac{1}{(a)_{n-m}} \sum_{j=m}^n (-1)^{j-m} \binom{n-m}{j-m} (x-a)_j (x+j)_{n-j}.$$

In the next place if $f(x)$ is a polynomial of degree $\leq n$, we may put

$$(4.10) \quad f(x) = \sum_{m=0}^n C_m (x-a)_m$$

where

$$(4.11) \quad C_m = \frac{1}{m!} \delta^m f(a)$$

and

$$\begin{aligned} \delta f(x) &= f(x) - f(x-1), \\ \delta^k f(x) &= \delta \cdot \delta^{k-1} f(x) = \sum_{j=0}^k (-1)^j \binom{k}{j} f(x-j). \end{aligned}$$

By (4.9) and (4.10)

$$\begin{aligned} (4.12) \quad f(x) &= \sum_{m=0}^n \frac{1}{(a)_{n-m}} C_m \sum_{j=m}^n (-1)^{j-m} \binom{n-m}{j-n} (x-a)_j (x+j)_{n-j} = \\ &= \sum_{j=0}^n (x-a)_j (x+j)_{n-j} \sum_{m=0}^j (-1)^{j-m} \binom{n-m}{j-m} \frac{C_m}{(a)_{n-m}}. \end{aligned}$$

By (4.11),

$$\begin{aligned} \sum_{m=0}^j (-1)^{j-m} \binom{n-m}{j-m} \frac{C_m}{(a)_{n-m}} &= \\ &= \sum_{m=0}^j (-1)^{j-m} \binom{n-m}{j-m} \frac{1}{m!(a)_{n-m}} \sum_{k=0}^m (-1)^k \binom{m}{k} f(a-k) = \\ &= (-1)^j \sum_{k=0}^j f(a-k) \sum_{m=k}^j (-1)^{m-k} \binom{n-m}{j-m} \binom{m}{k} \frac{1}{m!(a)_{n-m}}. \end{aligned}$$

Since

$$\begin{aligned} \sum_{m=k}^j (-1)^{m-k} \binom{n-m}{j-m} \binom{m}{k} \frac{1}{m!(a)_{n-m}} &= \\ &= \sum_{m=0}^{j-k} (-1)^m \binom{n-k-m}{j-k-m} \binom{k+m}{k} \frac{1}{(k+m)!(a)_{n-k-m}} = \\ &= \sum_{m=0}^{j-k} (-1)^m \frac{(n-k-m)!}{(n-j)!(j-k-m)!k!m!(a)_{n-k-m}} = \\ &= \frac{(n-k)!}{k!(n-j)!(j-k)!(a)_{n-k}} \sum_{m=0}^{j-k} \frac{(-j+k)_m (-a-n+k+1)_m}{m!(-n+k)_m} = \\ &= \frac{(n-k)!}{k!(n-j)!(j-k)!(a)_{n-k}} \frac{(a-1)_{j-k}}{(-n+k)_{j-k}} = (-1)^{j-k} \frac{(a-1)_{j-k}}{k!(j-k)!(a)_{n-k}}, \end{aligned}$$

we have

$$\sum_{m=0}^j (-1)^{j-m} \binom{n-m}{j-m} \frac{C_m}{(a)_{n-m}} = \sum_{k=0}^j (-1)^k \frac{(a-1)_{j-k}}{k!(j-k)!(a)_{n-k}} f(a-k).$$

Hence (4.12) becomes

$$f(x) = \sum_{j=0}^n (x-a)_j (x+j)_{n-j} \cdot \frac{1}{j!(a)_n} \sum_{k=0}^j (-1)^k \binom{j}{k} (a-1)_{j-k} (a+n-k)_k f(a-k),$$

in agreement with (4.7).

It remains to show that the representation (4.2) is unique. We shall assume that the parameter a is not equal to zero or a negative integer. If the representation (4.2) is not unique, then for some n , there exist numbers

$$B_k = B_k(a) \quad (0 \leq k \leq n),$$

not all zero, such that

$$(4.13) \quad \sum_{k=0}^n B_k (x-a)_k (x+k)_{n-k} = 0.$$

For $x = a$, (4.13) reduces to

$$(a)_n B_0 = 0,$$

so that $B_0 = 0$. Let

$$B_0 = \dots = B_{m-1} = 0, \quad B_m \neq 0$$

and take $x = a - m$. Then (4.13) reduces to

$$\sum_{k=m}^n B_k (-m)_k (a-m+k)_{n-k} = 0,$$

so that

$$(-1)^m m! (a)_{n-m} B_m = 0, \quad B_m = 0.$$

We have therefore proved the following

THEOREM A. *Let $f(x)$ be a polynomial of degree $\leq n$. Assume that a is not equal to zero or a negative integer. Then $f(x)$ is uniquely repre-*

sentable in the form

$$(4.14) \quad f(x) = \sum_{k=0}^n A_{n,k}(x-a)_k(x+k)_{n-k},$$

where the coefficients $A_{n,k}$ are determined by (4.7).

If a is equal to zero or a negative integer, the representation (4.14) is in general not possible. For example, for $a = 0$, (4.14) reduces to

$$f(x) = \sum_{k=0}^n A_{n,k}(x)_n,$$

so that $f(x)$ is a constant multiple of $(x)_n$. If $x = -m$, $m \geq 0$, (4.14) becomes

$$f(x) = \sum_{k=0}^n A_{n,k}(x+m)_k(x+k)_{n-k},$$

so that

$$(x)_m f(x) = \sum_{k=0}^n A_{n,k}(x)_{m+k}(x+k)_{n-k} = (x)_n \sum_{k=0}^n A_{n,k}(x+k)_m.$$

Hence, for $m \geq 0$, $f(x)$ is divisible by $(x+m)_{n-m}$.

5. We turn now to the q -analog of Saalschützer's theorem [1, p. 68], [3, p. 97]

$$(5.1) \quad \sum_{k=0}^n \frac{(q^{-n})_k(a)_k(b)_k}{(q)_k(c)_k(d)_k} q^k = \frac{(c/a)_n(c/b)_n}{(c)(c/ab)_n}$$

where now

$$(5.2) \quad (a)_n = (1-a)(1-qa) \dots (1-q^{n-1}a), \quad (a)_0 = 1$$

and

$$(5.3) \quad cd = q^{1-n}ab.$$

The following corollaries of (5.1) will be used. First, taking $b = d = 0$, we get

$$(5.4) \quad \sum_{k=0}^n \frac{(q^{-n})_k(a)_k}{(q)_k(c)_k} q^k = \frac{(c/a)_n}{(c)_n} a^n.$$

This may be called the q -analog of Vandermonde's theorem.

Next, if we let $n \rightarrow \infty$, (5.1) reduces to

$$(5.5) \quad \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(q)_k (c)_k} \left(\frac{c}{ab}\right)^k = \prod_{k=0}^{\infty} \frac{(1 - q^k c/a)(1 - q^k c/b)}{(1 - q^k c)(1 - q^k c/ab)}.$$

This is the q -analog of Gauss's theorem. Note that for $b = q^{-n}$, (5.5) becomes

$$(5.6) \quad \sum_{k=0}^n \frac{(q^{-n})_k (a)_k}{(q)_k (c)_k} \left(\frac{c}{ab}\right)^k = \frac{(c/a)_n}{(c)_n}.$$

Replacing k by $n - k$, (5.6) becomes

$$\sum_{k=0}^n \frac{(q^{-n})_k (q^{1-n}/c)_k}{(q)_k (q^{1-n}/a)_k} = (-1)^n q^{-\frac{1}{2}n(n-1)} \frac{(c-a)_n}{(c)_n}.$$

If we now replace q^{1-n}/c and q^{1-n}/a by a and c , respectively, we get (5.4).

In (5.1) replace b by c/b . Then (5.1) becomes

$$\sum_{k=0}^n \frac{(q^{-n})_k (a)_k (c/b)_k}{(q)_k (c)_k (q^{1-n} a/b)_k} q^k = \frac{(c/a)_n (b)_n}{(c)_n (b/a)_n}.$$

Since

$$\begin{aligned} (q^{-n})_k &= (-1)^k q^{-n k + \frac{1}{2}k(k-1)} (q)_n / (q)_{n-k}, \\ (q^{1-n} a/b)_k &= (-1)_k q^{-n k + \frac{1}{2}k(k+1)} (q^{n-k} b/a)_k (a/b)^k, \\ \frac{(b/a)_n}{(q^{1-n} a/b)_k} &= (-1)^k q^{n k - \frac{1}{2}k(k+1)} (b/a)_{n-k} (b/a)^k, \end{aligned}$$

we get

$$(5.7) \quad \frac{1}{(b)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (a)_k (b/a)_{n-k} \frac{(c/b)_k}{(c)_k} \left(\frac{b}{a}\right)^k = \frac{(c/a)_n}{(c)_n},$$

where

$$(5.8) \quad \begin{bmatrix} n \\ k \end{bmatrix} = \frac{(q)_n}{(q)_k (q)_{n-k}}.$$

We may think of (5.7) as a linear transformation from the set of rational functions

$$\frac{(x/b)_n}{(x)_n} \quad (n = 0, 1, 2, \dots)$$

to the set

$$\frac{(x/a)_n}{(x)_n} \quad (n = 0, 1, 2, \dots).$$

We define the linear transformation

$$(5.9) \quad T_{a,b}: x_n = \frac{1}{(b)_n} \sum_{k=0}^n \binom{n}{k} (a)_k (b/a)_{n-k} (b/a)^k y^k \quad (n = 0, 1, 2, \dots).$$

We assume that the parameters a, b, c, \dots are not equal to

$$(5.10) \quad q^{-t} \quad (t = 0, 1, 2, \dots).$$

It follows from (5.9) that

$$(5.11) \quad T_{a,a} = I \text{ (identity)}.$$

Also, as in the ordinary case, it is easy to show that $T_{a,b} = T_{c,d}$ if and only if

$$(i) \quad a = b, \quad c = d$$

or

$$(ii) \quad a = c, \quad b = d.$$

Indeed, by (5.9), $T_{a,b} = T_{c,d}$ if and only if

$$\frac{1}{(b)_n} (a)_k (b/a)_{n-k} (b/a)^k = \frac{1}{(d)_n} (c)_k (d/c)_{n-k} (d/c)^k \quad (0 \leq k \leq n).$$

For $k = n$, this reduces to

$$\frac{(a)_n}{(b)_n} \left(\frac{b}{a}\right)^n = \frac{(c)_n}{(d)_n} \left(\frac{d}{c}\right)^n \quad (n = 1, 2, 3, \dots).$$

Hence

$$\frac{1 - q^n a}{1 - q^n b} \frac{a}{b} = \frac{1 - q^n c}{1 - q^n d} \frac{c}{d} \quad (n = 1, 2, 3, \dots),$$

so that

$$\frac{1 - ax}{1 - bx} \frac{b}{a} \equiv \frac{1 - cx}{1 - dx} \frac{d}{c}.$$

We shall now show that

$$(5.12) \quad T_{a,b} T_{b,c} = T_{a,c}.$$

PROOF. Put

$$y_k = \frac{1}{(c)_k} \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix} (b)_j (c/b)_{k-j} (c/b)^j z_j \quad (k = 0, 1, 2, \dots).$$

Then, by (5.9) and (5.12),

$$\begin{aligned} x_n &= \frac{1}{(b)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (a)_k (b/a)_{n-k} (b/a)^k \cdot \frac{1}{(c)_k} \sum_{j=0}^k (b)_j (c/b)_{k-j} (c/b)^j z_j = \\ &= \frac{1}{(b)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (b)_j (c/b)^j z_j \sum_{k=j}^n \begin{bmatrix} n-j \\ k-j \end{bmatrix} (a)_k (b/a)_{n-k} (b/a)^k \frac{(c/b)_{k-j}}{(c)_k} = \\ &= \frac{1}{(b)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (b)_j (c/b)^j z_j \sum_{k=0}^{n-j} \begin{bmatrix} n-j \\ k \end{bmatrix} (a)_{j+k} (b/a)_{n-j-k} (b/a)^{j+k} \frac{(c/b)_k}{(c)_{j+k}} = \\ &= \frac{1}{(b)_n} \sum_{j=0}^n \frac{(a)_j (b)_j}{(c)_j} (c/a)^j z_j \sum_{k=0}^{n-j} \begin{bmatrix} n-j \\ k \end{bmatrix} (q^j a)_k (b/a)_{n-j-k} (b/a)^k \frac{(c/b)_k}{(q^j c)_k}. \end{aligned}$$

By (5.7),

$$\sum_{k=0}^{n-j} \begin{bmatrix} n-j \\ k \end{bmatrix} (q^j a)_k (b/a)_{n-j-k} (b/a)^k \frac{(c/b)_k}{(q^j c)_k} = \frac{(c/a)_{n-j} (q^j b)_{n-j}}{(q^j c)_{n-j}}$$

so that

$$\begin{aligned} x_n &= \frac{1}{(b)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} \frac{(a)_j (b)_j}{(c)_j} (c/a)^j \frac{(c/a)_{n-j} (q^j b)_{n-j}}{(q^j c)_{n-j}} z_j = \\ &= \frac{1}{(c)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (a)_j (c/a)_{n-j} (c/a)^j z_j. \end{aligned}$$

This proves (5.12).

We show next that

$$(5.13) \quad T_{b,c} T_{a,b} = T_{a,c}.$$

PROOF. Put

$$\begin{aligned} x_n &= \frac{1}{(c)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (b)_k (c/b)_{n-k} (c/b)^k y_k, \\ y_k &= \frac{1}{(b)_k} \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix} (a)_j (b/a)_{k-j} (b/a)^j z_j. \end{aligned}$$

Then

$$\begin{aligned} x_n &= \frac{1}{(c)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (c/b)_{n-k} (c/b)^k \cdot \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix} (a)_j (b/a)_{k-j} (b/a)^j z_j = \\ &= \frac{1}{(c)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (a)_j (b/a) z_j \sum_{k=0}^{j=0} \begin{bmatrix} n-j \\ k \end{bmatrix} (c/b)_{n-j-k} (b/a)_k (c/b)^k. \end{aligned}$$

It will therefore suffice to show that

$$(5.14) \quad \sum_{k=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (c/b)_{n-k} (b/a)^k (c/b)^k = (c/a)_n.$$

To prove (5.14), we make use of the identity

$$(5.15) \quad \sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} x^n = \prod_{n=0}^{\infty} \frac{1 - q^n a x}{1 - q^n x}.$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{x^n}{(q)_n} \sum_{k=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (c/b)_{n-k} (b/a)_k (c/b)^k &= \sum_{k=0}^{\infty} \frac{(b/a)_k}{(q)_k} (c/b)^k x^k \sum_{n=0}^{\infty} \frac{(c_k b)_n}{(q)_n} x^n = \\ &= \prod_{k=0}^{\infty} \frac{1 - q^n c x/a}{1 - q^n c x/b} \cdot \prod_{n=0}^{\infty} \frac{1 - q^n c x/b}{1 - q^n x} = \prod_{n=0}^{\infty} \frac{1 - q^n c x/a}{1 - q^n x} = \sum_{n=0}^{\infty} \frac{(c/a)_n}{(q)_n} x^n \end{aligned}$$

and (5.14) follows at once.

6. It is evident from (5.12) and (5.13) that

$$(6.1) \quad T_{a,b} T_{b,c} = T_{b,c} T_{a,b}.$$

We shall now prove that

$$(6.2) \quad T_{a,b} T_{c,d} = T_{c,d} T_{a,b}.$$

PROOF. By (5.12) and (5.13),

$$\begin{aligned} T_{a,b} T_{c,d} &= T_{c,b} T_{a,c} T_{c,d} \\ &= T_{c,b} T_{a,d} \\ &= T_{c,b} T_{b,d} T_{a,b} \\ &= T_{c,d} T_{a,b}. \end{aligned}$$

We shall now give another proof of (6.2) that makes use of an explicit formula for the transformation $T_{a,b}T_{c,d}$. Put

$$x_n = \frac{1}{(b)_n} \sum_{k=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (a)_k (b/a)_{n-k} (b/a)^k y_k,$$

$$y_k = \frac{1}{(d)_k} \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix} (c)_j (d/c)_{k-j} (d/c)^j z_j.$$

Then

$$(6.3) \quad x_n = \frac{1}{(b)_n} \sum_{k=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (a)_k (b/a)_{n-k} (b/a)^k \frac{1}{(d)_k} \sum_{j=0}^k \begin{bmatrix} k \\ j \end{bmatrix} (c)_j (d/c)_{k-j} (d/c)^j z_j =$$

$$= \frac{1}{(b)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (c)_j (d/c)^j z_j \sum_{k=j}^n \begin{bmatrix} n-j \\ k-j \end{bmatrix} (a)_k (b/a)_{n-k} \cdot (d/c)_{k-j} (b/a)^k / (d)_k =$$

$$= \frac{1}{(b)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} \frac{(a)_j (c)_j}{(d)_j} \left(\frac{bd}{ac}\right)^j z_j \cdot \sum_{k=0}^{n-j} \begin{bmatrix} n-j \\ k-j \end{bmatrix} (q^j a)_k (b/a)_{n-j-k} \frac{(d)_k}{(q^j d)_k} (b/a)^k.$$

Let

$$(6.4) \quad S_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (a)_k (b/a)_{n-k} \frac{(d/c)_k}{(d)_k} (b/a)^k,$$

$$(6.5) \quad R_n = (b)_n \sum_{s=0}^n (-1)^s \begin{bmatrix} n \\ s \end{bmatrix} \frac{(a)_s (c)_s}{(b)_s (d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{bd}{ac}\right)^s.$$

We shall show that

$$(6.6) \quad R_n = S_n.$$

PROOF. Clearly (6.6) is equivalent to

$$(6.7) \quad \sum_{n=0}^{\infty} R_n \frac{x^n}{(q)_n} = \sum_{n=0}^{\infty} S_n \frac{x^n}{(q)_n}.$$

We rewrite (5.15) in the form

$$(6.8) \quad \sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} x^n = \frac{e(a)}{e(ax)},$$

where

$$(6.9) \quad e(x) = \sum_{n=0}^{\infty} \frac{x^n}{(q)_n} = \prod_{n=0}^{\infty} (1 - q^n x)^{-1}.$$

Then

$$\begin{aligned} \sum_{n=0}^{\infty} S_n \frac{x^n}{(q)_n} &= \sum_{k=0}^{\infty} \frac{(a)_k (d/c)_k}{(q)_k (d)_k} (bx/a)^k \sum_{n=0}^{\infty} \frac{(b/a)_n}{(q)_n} x^n = \\ &= \sum_{k=0}^{\infty} \frac{(a)_k (d/c)_k}{(q)_k (d)_k} (bx/a)^k \frac{e(x)}{e(bx/a)}, \end{aligned}$$

$$\begin{aligned} \sum_{n=0}^{\infty} R_n \frac{x^n}{(q)_n} &= \sum_{s=0}^{\infty} (-1)^s \frac{(a)_s (c)_s}{(d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{bdx}{ac}\right)^s \sum_{n=0}^{\infty} \frac{(q^s b)_n}{(q)_n} x^n = \\ &= \sum_{s=0}^{\infty} (-1)^s \frac{(a)_s (c)_s}{(q)_s (d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{bdx}{ac}\right)^s \frac{e(x)}{e(q^s bx)}. \end{aligned}$$

Thus (6.7) is equivalent to

$$(6.10) \quad \sum_{k=0}^{\infty} \frac{(a)_k (d/c)_k}{(q)_k (d)_k} (bx/a)^k = \sum_{s=0}^{\infty} (-1)^s \frac{(a)_s (c)_s}{(q)_s (d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{bdx}{ac}\right)^s \frac{e(bx/a)}{e(q^s bx)}.$$

The right hand side of (6.10) is equal to

$$\begin{aligned} \sum_{s=0}^{\infty} (-1)^s \frac{(a)_s (c)_s}{(q)_s (d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{bdx}{ac}\right)^s \sum_{j=0}^{\infty} \frac{(q^s a)_j}{(q)_j} (bx/a)^j &= \\ &= \sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} \left(\frac{bx}{a}\right)^n \sum_{s=0}^n (-1)^s \begin{bmatrix} n \\ s \end{bmatrix} \frac{(c)_s}{(d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{d}{c}\right)^s = \\ &= \sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} \left(\frac{bx}{a}\right)^n \sum_{s=0}^n \frac{(q^{-n})_s (c)_s}{(q)_s (d)_s} \left(\frac{q^n d}{c}\right)^s = \sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} \left(\frac{bx}{a}\right)^n \frac{(d/c)_n}{(d)_n}, \end{aligned}$$

by (5.6). This evidently proves (6.10) and therefore proves (6.6). For an identity containing (6.10) see [2].

Returning to (6.3), we have

$$\begin{aligned} x_n &= \frac{1}{(b)_n} \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} \frac{(a)_j (c)_j}{(d)_j} \left(\frac{bd}{ac}\right)^j z_j \cdot \\ &\quad \cdot (q^j b)_{n-j} \sum_{s=0}^{n-j} (-1)^s \begin{bmatrix} n-j \\ s \end{bmatrix} \frac{(q^j a)_s (q^j c)_s}{(q^j b)_s (q^j d)_s} q^{\frac{1}{2}s(s-1)} \left(\frac{bd}{ac}\right)^s. \end{aligned}$$

Rearranging and simplifying, we get

$$(6.11) \quad x_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} \frac{(a)_k (c)_k}{(b)_k (d)_k} \left(\frac{bd}{ac}\right)^k \sum_{s=0}^k (-1)^s \begin{bmatrix} k \\ s \end{bmatrix} q^{\frac{1}{2}s(s-1)} z_{k-s}.$$

Thus we have an explicit formula for the transformation $T_{a,b}T_{c,d}$. Clearly (6.11) implies (6.1).

7. We may rewrite (5.7) in the following form.

$$(7.1) \quad (x/a)_n = \frac{1}{(b)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (a)_k (b/a)_{n-k} (b/a)^k (x/b)_k (q^k x)_{n-k}.$$

This suggests the following problem. Let $f(x)$ be a polynomial of degree $\leq n$. Assume that $f(x)$ can be written in the form

$$(7.2) \quad f(x) = \sum_{k=0}^n A_{n,k} (x/a)_k (q^k x)_{n-k},$$

where a is an arbitrary constant and

$$A_{n,k} = A_{n,k}(a, q)$$

is independent of x . We shall show how to evaluate the coefficients $A_{n,k}$.

Taking $x = a$ in (7.2), we get

$$(7.3) \quad (a)_n A_{n,0} = f(a).$$

Next, for $x = q^{-1}a$, (7.2) reduces to

$$f(q^{-1}a) = (q^{-1}a)_n A_{n,0} - q^{-1}(1-q)(a)_{n-1} A_{n,1},$$

so that

$$(7.4) \quad (1-q)(a)_n A_{n,1} = q(1-q^{-1}a)f(a) - q(1-q^{n-1}a)f(q^{-1}a).$$

For $x = q^{-2}a$, (7.2) becomes

$$f(q^{-2}a) = (q^{-2}a)_n A_{n,0} - q^{-2}(1-q^2)(q^{-1}a)_{n-1} A_{n,1} + q^{-1}(1-q^2)(a)_{n-2} A_{n,2}.$$

This yields

$$(7.5) \quad (q)_2(a)_n A_{n,2} = q^2(1-q^{-1}a)(1-a)f(a) - q^2(1+q)(1-q^{-1}a)(1-q^{n-1}a)f(q^{-1}a) + q(1-q^{n-1}a)(1-q^{n-2}a)f(q^{-2}a).$$

For $x = q^{-3}a$, we get

$$(7.6) \quad (q)_3(a)_n A_{n,3} = q^3(q^{-1}a)_3 f(a) - q^3(1+q+q^2)(q^{-1}a)_2 (q^{n-1}a) f(q^{-1}a) + q^4(1+q+q^2)(q^{-1}a)_1 (q^{n-2}a)_2 f(q^{-2}a) - q^6(q^{n-3}a)_3 f(q^{-3}a).$$

This suggests the general formula

$$(7.7) \quad (q)_k(a)_k A_{n,k} = q^k \sum_{j=0}^k (-1)^j \begin{bmatrix} k \\ j \end{bmatrix} (q^{n-j}a)_j q^{\frac{1}{2}j(j-1)} f(q^{-j}a).$$

PROOF OF (7.7). To begin with, we have

$$(7.8) \quad (a)_n = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\frac{1}{2}k(k-1)} (x/a)_k (q^k x)_{n-k} a^k.$$

Indeed, since

$$(1 - q^n) \dots (1 - q^{n-k+1}) = (-1)^k q^{n k - \frac{1}{2}k(k-1)} (q^{-n})_k,$$

it follows from (5.6) that

$$\begin{aligned} \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\frac{1}{2}k(k-1)} (x/a)_k (q^k x)_{n-k} a^k &= \\ &= (x)_n \sum_{k=0}^n \frac{(q^{-n})_k (x/a)_k}{(q)_k (x)_k} (q^n a)^k = (x)_n \frac{(a)_n}{(x)_n} = (a)_n. \end{aligned}$$

This proves (7.8). Now replace x by $q^m x$ in (7.8) and multiply by $(x/a)_m$. Then

$$(a)_n (x/a)_m = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\frac{1}{2}k(k-1)} (x/a)_{k+m} (q^{k+m} x)_{n-k} a^k a^k.$$

Changing the notation slightly, this becomes

$$(7.9) \quad (x/a)_m = \frac{(a)_{n-m}}{1} \sum_{k=m}^n (-1)^{k-m} \begin{bmatrix} n-m \\ k-m \end{bmatrix} q^{\frac{1}{2}(k-m)(k-m-1)} \cdot (x/a)_k (q^k x)_{n-k} a^{k-m}.$$

Next, if $f(x)$ is an arbitrary polynomial of degree $\leq n$, we may put

$$(7.10) \quad f(x) = \sum_{m=0}^n C_m (x/a)_m,$$

where $C_m = C_m(a, q)$ is independent of x . It is easily verified that

$$(7.11) \quad (q)_k C_k = (qa)^k \delta^k f(a),$$

where

$$(7.12) \quad \begin{cases} \delta f(x) = \frac{f(x) - f(q^{-1}x)}{x}, \\ \delta^k f(x) = \delta \cdot \delta^{k-1} f(x) = x^{-k} \sum_{j=0}^k (-1)^j \begin{bmatrix} k \\ j \end{bmatrix} q^{\frac{1}{2}j(j-1)} f(q^{-j}x). \end{cases}$$

It follows from (7.9) and (7.10) that

$$\begin{aligned} f(x) &= \sum_{m=0}^n \frac{C_m}{(a)_{n-m}} \sum_{k=m}^n (-1)^{k-m} \begin{bmatrix} n-m \\ k-m \end{bmatrix} q^{\frac{1}{2}(k-m)(k-m-1)} (x/a)_k (q^k x)_{n-k} a^{k-m} = \\ &= \sum_{k=0}^n (x/a)_k (q^k x)_{n-k} \sum_{m=0}^k (-1)^{k-m} \begin{bmatrix} n-m \\ k-m \end{bmatrix} q^{\frac{1}{2}(k-m)(k-m-1)} a^{k-m} \frac{C_m}{(a)_{n-m}}. \end{aligned}$$

By (7.11) and (7.12),

$$(q)_m C_m = q^m \sum_{j=0}^m (-1)^j \begin{bmatrix} m \\ j \end{bmatrix} q^{\frac{1}{2}j(j-1)} f(q^{-j}a),$$

so that

$$\begin{aligned} \sum_{m=0}^k (-1)^{k-m} \begin{bmatrix} n-m \\ k-m \end{bmatrix} q^{\frac{1}{2}(k-m)(k-m-1)} a^{k-m} \frac{C_m}{(a)_{n-m}} &= \sum_{m=0}^k (-1)^{k-m} \begin{bmatrix} n-m \\ k-m \end{bmatrix} \cdot \\ &\cdot q^{\frac{1}{2}(k-m)(k-m-1)} a^{k-m} \cdot \frac{q^m}{(q)_m (a)_{n-m}} \sum_{j=0}^m (-1)^j \begin{bmatrix} m \\ j \end{bmatrix} q^{\frac{1}{2}j(j-1)} f(q^{-j}a) = \\ &= \sum_{j=0}^k (-1)^j q^{\frac{1}{2}j(j-1)} f(q^{-j}a) \sum_{m=j}^k (-1)^{k-m} q^{\frac{1}{2}(k-m)(k-m-1)+m} a^{k-m} \cdot \\ &\cdot \frac{(q)_m (a)_{n-m}}{1} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} n-m \\ k-m \end{bmatrix} = \sum_{k-j}^k (-1)^j q^{\frac{1}{2}j(j-1)} f(q^{-j}a) \cdot \\ &\cdot \sum_{m=0}^{j-1} (-1)^m q^{\frac{1}{2}m(m-1)+k-m} a^m \cdot \frac{1}{(q)_{k-m} (a)_{n-k+m}} \begin{bmatrix} k-m \\ j \end{bmatrix} \begin{bmatrix} n-k+m \\ m \end{bmatrix}. \end{aligned}$$

The inner sum is equal to

$$\begin{aligned} \sum_{m=0}^{k-j} (-1)^m q^{\frac{1}{2}m(m-1)+k-m} a^m \frac{(q)_{n-k+m}}{(q)_j (q)_{k-j-m} (q)_m (a)_{n-k+m}} &= \\ = \frac{q^k}{(q)_j (q)_{k-j} (a)_{n-k}} \sum_{m=0}^{k-j} \frac{(a^{-k+j})_m (q^{n-k+1})_m}{(q)_m (q^{n-k} a)_m} q^{(k-j-1)m} a^m &= \\ = \frac{q^k}{(q)_j (q)_{k-j} (a)_{n-k}} \frac{(q^{-1}a)_{k-j}}{(q^{n-k} a)_{k-j}} = \frac{q^k (q^{-1}a)_{k-j}}{(q)_j (q)_{k-j} (a)_{n-j}}. \end{aligned}$$

We have therefore

$$\begin{aligned}
 (7.13) \quad f(x) &= \sum_{k=0}^n (x/a)_k (q^k x)_{n-k} \sum_{j=0}^k (-1)^s q^{\frac{1}{2}s(s-1)} f(q^{-j} a) \cdot \\
 &\quad \cdot \frac{q^k (q^{-1} a)_{k-j}}{(q_j)(q)_{k-j} (a)_{n-j}} = \sum_{k=0}^n (x/a)_k (q^k x)_{n-k} \frac{q^k}{(q)_k (a)_n} \cdot \\
 &\quad \cdot \sum_{j=0}^k (-1)^j \begin{bmatrix} k \\ j \end{bmatrix} (q^{-1} a)_{k-j} (q^{n-j} a)_j q^{\frac{1}{2}j(j-1)} f(q^{-j} a) .
 \end{aligned}$$

It remains to show that the coefficients $A_{n,k}$ in (7.2) are uniquely determined. Otherwise we should have

$$(7.14) \quad \sum_{k=0}^n B_k (x/a)_k (q^k x)_{n-k} = 0 ,$$

where not all the B_k equal zero. For $x = a$, (7.14) implies $B_0 = 0$. Assume that

$$B_0 = \dots = B_{m-1} = 0 , \quad B_m \neq 0 .$$

For $x = q^{-m} a$, (7.14) reduces to

$$(q^{-m})_m (a)_{n-m} B_m = 0 .$$

Since a is not equal to q^{-k} , $k = 0, 1, 2, \dots$, we get $B_m = 0$.

We have therefore proved the following.

THEOREM B. *Assume $a \neq q^{-k}$, where k is a nonnegative integer. Let $f(x)$ be a polynomial of degree $\leq n$. Then the coefficients in*

$$f(x) = \sum_{k=0}^n A_{n,k} (x/a)_k (q^k a)_{n-k}$$

are uniquely determined by

$$(q)_k (a)_k A_{n,k} = q^k \sum_{j=0}^k (-1)^j \begin{bmatrix} k \\ j \end{bmatrix} (q^{-1} a)_{k-j} (q^{n-j} a)_j q^{\frac{1}{2}j(j-1)} f(q^{-j} a) .$$

REFERENCES

- [1] W. N. BAILEY, *Generalized Hypergeometric Series*, Cambridge, 1935.
- [2] N. A. HALL, *An algebraic identity*, Journal of the London Mathematical Society, **11** (1936), 276.
- [3] L. J. SLATER, *Generalized Hypergeometric Functions*, Cambridge, 1966.

Manoscritto pervenuto in redazione il 1 settembre 1972.