

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

D. FORTUNATO

G. PALMIERI

**Remarks on the Yamabe problem and the
Palais-Smale condition**

Rendiconti del Seminario Matematico della Università di Padova,
tome 75 (1986), p. 47-65

http://www.numdam.org/item?id=RSMUP_1986__75__47_0

© Rendiconti del Seminario Matematico della Università di Padova, 1986, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Remarks on the Yamabe Problem and the Palais-Smale Condition.

D. FORTUNATO - G. PALMIERI (*) (**)

0. Introduction.

Let M be a C^∞ compact Riemannian manifold of dimension $n \geq 3$. Let $g(x)$ be its metric and $R(x)$ its scalar curvature. An important problem concerning the scalar curvature is the Yamabe problem: does there exist a metric g' conformal to g , such that the scalar curvature \bar{R} of the metric is constant?

Since the pionering paper of Yamabe [11] appeared, several authors have studied this problem (cp. [1, 9, 10] and the references contained in [3]).

If we consider the conformal deformation in the form $g' = u^{4/(n-2)} g$ (with $u \in C^\infty$, $u > 0$), the Yamabe problem is reduced (cp. [1, 9, 11]) to the following eigenvalue problem:

$$(0.1) \quad \begin{cases} \text{find } u \in C^\infty, u > 0 \text{ and } \bar{R} \in \mathbb{R} \text{ s.t. ,} \\ \gamma \Delta u + R(x)u = \bar{R}u^{2^*-1}, \end{cases}$$

where $\gamma = 4(n-1)/(n-2)$, Δ denotes the Laplace-Beltrami operator corresponding to g and $2^* = 2n/(n-2)$.

We denote by H^1 the Sobolev space on M , i.e. H^1 is the completion

(*) Indirizzo degli AA.: Dipartimento di Matematica, Università di Bari, Via G. Fortunato, 70125 Bari, Italy.

(**) Work supported by Ministero Pubblica Istruzione Italy (fondi 40% e 60%) and by G.N.A.F.A. of C.N.R.

of the space $C^\infty(M)$ with respect to the norm

$$\|\varphi\| = \left(\int_M (|\varphi|^2 + |\nabla\varphi|^2) dM \right)^{\frac{1}{2}}$$

where $|\nabla\varphi|^2 = g^{ij}\varphi_{x_i}\varphi_{x_j}$.

$L^\alpha(\alpha > 1)$ denotes the space of the functions on M which are α -integrable. Moreover we set

$$|\varphi|_\alpha = \left(\int_M |\varphi|^\alpha dM \right)^{1/\alpha}.$$

Clearly if u minimizes the functional

$$(0.2) \quad \psi(u) = \int_M (\gamma|\nabla u|^2 + R(x)u^2) dM$$

on the manifold in H^1

$$(0.3) \quad V = \{u \in H^1 \mid |u|_{2^*} = 1\}$$

u is a weak solution of (0.1) and \bar{R} is the minimum of ψ on V . Then by a regularity result of Trudinger (cp. [9, Th. 3]) u is C^∞ . Moreover, since $|\nabla|u|| = |\nabla u|$ (cp. [3, prop. 3.49]), we can assume that $u \geq 0$. Then (cp. [3, prop. 3.75]) we deduce that $u > 0$.

Therefore the Yamabe problem is solved if the functional ψ can be minimized on V . Since H^1 is not compactly embedded into L^{2^*} it is not easy to find a minimum directly. In [1, 9, 11] the following approximation-procedure has been used. Since the embedding $H^1 \hookrightarrow L^q$ ($q < 2^*$) is compact the problem

$$-\gamma\Delta u + R(x)u = \omega_q u^{q-1} \quad |u|_q = 1$$

can be easily solved; then, by taking the limit for $q \rightarrow 2^*$, it is possible, in some cases, to solve (0.1).

If Ω is a bounded domain in \mathbb{R}^n , set

$$S = \inf \left\{ \int_\Omega |\nabla\varphi|^2 dx : |\varphi|_{2^*} = 1, \quad \varphi \in C_0^\infty(\mathbb{R}^n) \right\} \quad (n \geq 3).$$

It is known (cp. [2, 8]) that S is independent of Ω and

$$S = \frac{\omega_n^{2/n}}{4} n(n-2)$$

where ω_n denotes the area of the unit n -dimensional sphere S_n .

Using the « approximation-method » Aubin [1, Th. 7] proved the following Theorem.

THEOREM 0.1. *Suppose that*

$$\inf \{ \psi(u) : u \in V \} < S \cdot \gamma .$$

Then there exists $u \in V$ which minimized $\psi|_V$.

In [1] Aubin has also pointed out that

$$\mu = \inf \{ \psi(u) : u \in V \}$$

is a conformal invariant and that the inequality

$$\mu \leq \gamma S$$

holds for any compact n -dimensional Riemmanian manifold.

In this paper we prove that $\psi|_V$ satisfies the Palais-Smale (P-S) condition in the range $]-\infty, \gamma S[$ (cp. lemma 1.1). Then Theorem 0.1 can be proved directly using standard variational arguments.

Moreover we show that $\psi|_V$ does not satisfy the (P-S) condition at $c = \gamma S$ (cp. Th. 2.1, 2.2).

It can be proved also that slight perturbations of (0.1) have always a solution. More precisely we consider the following perturbed eigenvalue problem:

$$(0.4) \quad \begin{cases} \text{find } u \in C^\infty, u > 0 \text{ and } \bar{R} \in \mathbb{R} \text{ s.t. ,} \\ -\gamma \Delta u + (R(x) - \varepsilon)u = \bar{R}u^{2^*-1}, & \varepsilon > 0, \end{cases}$$

and we prove the following

THEOREM 0.2. *Let $n \geq 4$, than for any $\varepsilon > 0$, (0.4) possesses a solution.*

Observe that Theorem 0.2 is similar to a result obtained by Brezis and Nirenberg (cp. [4, Th. 11]) in the context of elliptic boundary value problems on a bounded domain $\Omega \subset \mathbb{R}^n$.

Let us finally recall that another direct proof of Theorem 0.1 is contained in [10].

1. Proof of Theorem 0.1, 0.2.

The following lemma plays a fundamental role in proving Theorems 0.1, 0.2.

LEMMA 1.1. *The functional $\psi|_V$ defined by (0.2) and (0.3) satisfies the Palais-Smale condition in $]-\infty, \gamma S[$ in the following sense:*

(P-S) If $\{u_j\} \subset V$ s.t. as $j \rightarrow \infty$

$$(1.1) \quad d\psi|_V(u_j) \rightarrow 0 \text{ strongly in } H^{-1}$$

and

$$(1.2) \quad \psi(u_j) \rightarrow c, \quad c < \gamma S$$

Then $\{u_j\}$ contains a subsequence converging strongly in H^1 .

PROOF. Let $\{u_j\} \subset V$ be a sequence which satisfies (1.1) and (1.2). Then, by (1.2), we easily deduce that we can select a subsequence, which we continue to denote by $\{u_j\}$, such that

$$(1.3) \quad \begin{cases} u_j \rightarrow u \text{ weakly in } H^1, \\ u_j \rightarrow u \text{ strongly in } L^p, \end{cases} \quad 1 \leq p < 2^*.$$

Set $\varphi(u) = |u|_{2^*}^{2^*}$ $u \in H^1$.

From (1.1) we deduce that there exists a sequence $\{\lambda_j\} \subset \mathbb{R}$ s.t.

$$(1.4) \quad d\psi(u_j) - \lambda_j d\varphi(u_j) \rightarrow 0 \quad \text{in } H^{-1} = (H^1)'$$

Then, since $|u_j|_{2^*} = 1$ and by using (1.2), we obtain

$$(1.5) \quad \lambda_j = \frac{\langle d\psi(u_j), u_j \rangle}{2^*} + o(1) = \frac{2}{2^*} \psi(u_j) + o(1) = \frac{2}{2^*} c + o(1).$$

We show now that u solves the equation

$$(1.6) \quad -\gamma \Delta u + R(x)u - cu|u|^{2^*-2} = 0.$$

Let $\zeta \in C^\infty(M)$, then by (1.4), (1.3), (1.5) we deduce that

$$\begin{aligned} o(1) &= \langle d\psi(u_j) - \lambda_j d\varphi(u_j), \zeta \rangle = \langle d\psi(u_j), \zeta \rangle - \frac{2}{2^*} c \langle d\varphi(u_j), \zeta \rangle + o(1) = \\ &= 2 \langle -\gamma \Delta u + R(x)u, \zeta \rangle - 2c \int_M u|u|^{2^*-2} \cdot \zeta \, dM + o(1). \end{aligned}$$

Then u is a weak solution of (1.6). Then (cp. [9, Th. 3]) u is a C^∞ solution of (1.6).

To show that $u_j \rightarrow u$ strongly in $H^1(M)$, let

$$v_j = u_j - u.$$

Testing (1.4) with v_j , we obtain

$$\begin{aligned} (1.7) \quad o(1) &= \langle d\psi(u_j), v_j \rangle - \lambda_j \langle d\varphi(u_j), v_j \rangle = \\ &= 2 \int_M [\gamma (\nabla u | \nabla v_j) + |\nabla v_j|^2] \, dM - \\ &\quad - \lambda_j \langle d\varphi(u_j), v_j \rangle. \end{aligned}$$

By (1.3) we have

$$(1.8) \quad \int_M [\gamma (\nabla u | \nabla v_j) + R(x) (uv_j + v_j^2)] \, dM = o(1).$$

Whence from (1.7), (1.8) we deduce that

$$\begin{aligned} (1.9) \quad 2\gamma |\nabla v_j|_2^2 &= \lambda_j \langle d\varphi(u_j), v_j \rangle + o(1) = \\ &= 2^* \lambda_j \int_M |u + v_j|^{2^*-2} (u + v_j) v_j \, dM + o(1) \end{aligned}$$

Now we claim that

$$(1.10) \quad \gamma \|v_j\|^2 = \gamma |\nabla v_j|_2^2 + o(1) = c |v_j|_2^{2^*} + o(1).$$

In fact, following an analogous argument as in [5, lemma 2.1], we obtain :

$$\begin{aligned}
 \left| \int_M [(u + v_j)|u + v_j|^{2^*-2}v_j - |v_j|^{2^*}] dM \right| &= \\
 &= \left| \int_M \int_0^{u(x)} \frac{\partial}{\partial \xi} [(v_j + \xi)|v_j + \xi|^{2^*-2}] v_j d\xi dM \right| = \\
 &= (2^* - 1) \left| \int_M \int_0^1 |v_j + tu|^{2^*-2} v_j u dt dM \right| \leq \\
 &\leq \text{const} \left(\int_M (|u| |v_j|^{2^*-1} + |v_j| |u|^{2^*-1}) dM \right).
 \end{aligned}$$

Then, by (1.3) and since $u \in L^{2^*}(M)$, we have

$$(1.11) \quad \int_M (u + v_j) |u + v_j|^{2^*-2} v_j dM = |v_j|_{2^*}^{2^*} + o(1).$$

So (1.10) easily follows from (1.9), (1.11) and (1.5). If $c \leq 0$ from (1.10) we deduce that

$$\|v_j\| = o(1).$$

Whence

$$u_j \rightarrow u \text{ strongly in } H^1.$$

Now suppose that

$$(1.12) \quad 0 < c < \gamma S.$$

Using Theorem 2.21 in [3] (cp. also [5, lemma 2.5]) we have

$$(1.13) \quad \|v_j\|^2 \geq S |v_j|_{2^*}^{2^*} + o(1).$$

By (1.10), (1.13)

$$\begin{aligned}
 \|v_j\|^{2^*} &\geq S^{2^*/2} \frac{\gamma}{c} \|v_j\|^2 + o(1), \\
 (1.14) \quad \|v_j\|^2 &\left(\frac{\gamma}{c} S^{2^*/2} - \|v_j\|^{2^*-2} \right) \leq o(1).
 \end{aligned}$$

Let us now prove that

$$(1.15) \quad \|v_j\|^2 \leq \frac{c}{\gamma},$$

Easy calculations show that

$$(1.16) \quad \begin{aligned} \psi(u_j) = \psi(u + v_j) &= \gamma \int_M (|\nabla u|^2 + |\nabla v_j|^2) dM + \\ &+ \int_M R(x) u^2 dM + o(1). \end{aligned}$$

By (1.16) and (1.2) we deduce that

$$(1.17) \quad \gamma \|v_j\|^2 = c - \int_M (\gamma |\nabla u|^2 + R(x) u^2) dM + o(1).$$

Moreover, since u solves (1.6), we have

$$(1.18) \quad \int_M (\gamma |\nabla u|^2 + R(x) u^2) dM = c |u|_2^{2^*}.$$

From (1.17) and (1.18) we easily deduce (1.15).

By (1.14) and (1.15) we have

$$(1.19) \quad \|v_j\|^2 \left(\frac{\gamma}{c} S^{2^*/2} - \left(\frac{c}{\gamma} \right)^{(2^*-2)/2} \right) \leq o(1).$$

Since $c < \gamma S$, we have

$$(1.20) \quad \frac{\gamma}{c} S^{2^*/2} > \left(\frac{c}{\gamma} \right)^{(2^*-2)/2}.$$

Finally from (1.19) and (1.20) we deduce that

$$\|v_j\| \rightarrow 0 \quad \text{Q.E.D. .}$$

PROOF OF THEOREM 0.1. The proof of Theorem 0.1 can be deduced from lemma 1.1 using standard variational arguments. Suppose

$$\mu = \inf \psi|_V < \gamma S$$

and arguing by contradiction assume that μ is not a critical value for $\psi|_V$. Since the (P-S) condition holds in $]-\infty, \gamma S[$ by well known results (cp. [6, 7]) there exists $\varepsilon > 0$ and an homeomorphism

$$\eta: V \rightarrow V \text{ s.t.}$$

$$(u \in V, \psi(u) \leq \mu + \varepsilon) \Rightarrow (\psi(\eta(u)) \leq \mu - \varepsilon).$$

Obviously this contradicts $\mu = \inf \psi|_V$. **Q.E.D.**

REMARK 1.2. From the proofs given before it is easy to see that lemma 1.1 and Theorem 0.1 still hold if $R(x)$ is any smooth function which does not represent necessarily the scalar curvature of M .

Moreover lemma 1.1 and Theorem 0.1 hold also, with obvious changes, if we replace M with a bounded open set $\Omega \subset \mathbb{R}^n$, $n \geq 3$, and H^1 with the Sobolev space $H_0^1(\Omega)$. In this setting Theorem 0.1 becomes a variant of Lemma 1.2 in [4].

REMARK 1.3. In order to apply Theorem 0.1 it is important to find conditions which guarantee

$$(1.21) \quad \mu = \inf \psi|_V < \gamma S.$$

It is easy to see that if we assume $\int_M R(x) dM < (\text{meas } M)^{(n-2)/n} \gamma S$, then (1.21) is verified (cp. also [1, 3, 9]).

Other classes of manifolds which satisfy (1.21) can be found in [1, 3].

2. Remarks about the (P-S) condition and Proof of Theorem 0.2.

For any Riemannian manifold [1, lemma 4] we have

$$\mu = \inf \psi|_V \leq \gamma S.$$

Moreover there exist manifolds, which we call critical, for which

$$\mu = \gamma S.$$

An example of critical manifold is the n -dimensional sphere $S_n (n \geq 3)$ [1, cor. 4].

For this class of manifolds the (P-S) condition does not hold in $\mu = \gamma S$.

In fact the following Theorem holds

THEOREM 2.1. *Let M be a critical Riemannian manifold i.e.*

$$\mu = \inf \psi|_V = \gamma S .$$

Then $\psi|_V$ does not satisfy the (P-S) condition in $\mu = \gamma S$, i.e. there exists a sequence $\{u_j\} \subset V$ s.t. $\psi(u_j) \rightarrow \gamma S$ and $d\psi|_V(u_j) \rightarrow 0$ and which is not precompact in H^1 .

PROOF. By a result of Aubin [3, Th. 2.21] there exists a sequence $\{\omega_j\} \subset V$ s.t.

$$|\omega_j|_2 \rightarrow 0 \quad \text{and} \quad |\nabla \omega_j|_2^2 \rightarrow S .$$

Then

$$(2.1) \quad \psi(\omega_j) \rightarrow \gamma S .$$

Since the manifold M is critical, by (2.1) we deduce that $\{\omega_j\}$ is a minimizing sequence for $\psi|_V$.

Now we argue by contradiction and suppose that $\psi|_V$ satisfies the (P-S) condition at $\mu = \gamma S$. Then by using standard variational arguments (cp. prop. A.1 in Appendix)

$$d\psi|_V(\omega_j) \rightarrow 0 .$$

Then $\{\omega_j\}$ is precompact in H^1 , therefore it contains a subsequence, which we continue to denote by $\{\omega_j\}$, such that

$$\omega_j \rightarrow \omega \text{ strongly in } H^1 \text{ (and in } L^{2^*}) .$$

Then, since $|\omega_j|_{2^*} = 1$, we have $|\omega|_{2^*} = 1$. On the other hand $|\omega|_2 = 0$. Therefore we get a contradiction. **Q.E.D.**

Also manifolds which are not critical do not satisfy the (P-S) condition at $c = \gamma S$. More precisely the following Theorem holds

THEOREM 2.2. *Let M be a C^∞ Riemannian compact manifold of dimension $n \geq 3$. Then the functional $\psi|_V$ does not satisfy the (P-S) condition in $c = \gamma S$.*

In order to prove this result we need to introduce some notations.

Let δ be a positive number less than the injectivity radius of M . If $P \in M$ we set $B(P, \delta)$ the ball of M centered at P and radius δ . Using geodesic local coordinates, if $Q \in B(P, \delta)$ we have

$$(2.2) \quad Q = (r, \theta) \in [0, \delta] \times S_{n-1}, \quad r = r(P, Q) = d(P, Q).$$

The metric tensor g can be expressed by

$$(2.3) \quad ds^2 = dr^2 + r^2 g_{\theta_i \theta_j}(r, \theta) d\theta^i d\theta^j.$$

Set

$$(2.4) \quad |g| = \det(g_{\theta_i \theta_j}).$$

Fix $\alpha > 0$. Suppose $R(P) > 0$ we choose

$$\alpha^2 = R(P)/n(n-1).$$

It is not restrictive to assume also

$$(2.5) \quad \alpha\delta < \pi.$$

In order to prove Theorem 2.2 we need to introduce a suitable sequence in V .

Following Aubin [1] we consider the sequence of functions

$$k \in \mathbb{N}, \quad v_k(Q) = \begin{cases} \left(\frac{1}{k} + \frac{1 - \cos \alpha r}{\alpha^2} \right)^{1-n/2} - \left(\frac{1}{k} + \frac{1 - \cos \alpha \delta}{\alpha^2} \right)^{1-n/2} & \text{if } r < \delta, \\ 0 & \text{if } r > \delta. \end{cases}$$

Observe that the functions $r \rightarrow [1/k + (1 - \cos \alpha r)/\alpha^2]^{1-n/2}$ solve the Yamabe equation for $M = S_n$.

Obviously v_k have supports contained in $\overline{B(P, \delta)}$. We set

$$(26) \quad u_k = \frac{v_k}{|v_k|_{2^*}}.$$

In order to prove Theorem 2.2 we need the following lemma

LEMMA 2.3. *For $k \rightarrow +\infty$ we have the following asymptotic estimates*

$$(2.7) \quad \psi(u_k) = \gamma S + \sigma_k$$

where

$$(2.8) \quad \sigma_k = \begin{cases} O(k^{-2}) & \text{if } n > 6, \\ O(\lg k \cdot k^{-2}) & \text{if } n = 6, \\ O(k^{-(n-2)/2}) & \text{if } n = 3, 4, 5. \end{cases}$$

Moreover

$$(2.9) \quad |u_k|_2^2 = \begin{cases} \frac{\gamma}{2} \omega_n^{2/n} k^{-1} + o(k^{-1}) & \text{if } n \geq 5, \omega_n = \text{meas } S_n, \\ \frac{\gamma}{2} \omega_4^{1/2} \frac{\lg k}{k} + o(k^{-1} \lg k) & \text{if } n = 4, \\ \text{const } k^{-1/2} + o(k^{-1/2}) & \text{if } n = 3, \end{cases}$$

and the same asymptotic estimates hold for $|r|\nabla u_k|_2^2$ with different constants.

The proof of lemma 2.3 is quite technical and it will be sketched in Appendix.

PROOF OF THEOREM 2.2. Consider the sequence defined in (2.6). Obviously $|u_k|_{2^*} = 1$.

Moreover by lemma 2.3

$$\psi(u_k) \rightarrow \gamma S \quad \text{and} \quad |u_k|_2 \rightarrow 0.$$

Then $\{u_k\}$ is not precompact in H^1 . Therefore, in order to verify that $\psi|_V$ does not satisfy the (P-S) condition in γS , we need only to prove that

$$d\psi|_V(u_k) \rightarrow 0 \text{ strongly in } H^{-1}.$$

Now

$$\Delta u_k(r) = u_k''(r) + \frac{n-1}{r} u_k'(r) + u_k'(r) \frac{\partial}{\partial r} \lg \sqrt{|g|}$$

and there exists $A > 0$ such that

$$(2.10) \quad \left| \frac{\partial}{\partial r} \lg \sqrt{|g|} \right| < Ar.$$

By the above expression for $\Delta u_k(r)$, we have

$$(2.11) \quad \begin{aligned} -\gamma \Delta u_k + R(x) u_k - n(n-1) \omega_n^{2/n} \cdot u_k^{(n+2)/(n-2)} = \\ = Lu_k - n(n-1) \omega_n^{2/n} u_k^{(n+2)/(n-2)} + \\ + \gamma u_k' \frac{\partial}{\partial r} \left[\lg \left(\frac{\sin \alpha r}{r} \right)^{n-1} - \lg \sqrt{|g|} \right] + u_k [R(x) - \alpha^2 n(n-1)] = \chi_k \end{aligned}$$

where

$$(2.12) \quad Lu_k = -\gamma \left(u_k'' + \frac{n-1}{r} u_k' + u_k' \frac{\partial}{\partial r} \lg \left(\frac{\sin \alpha r}{r} \right)^{n-1} \right) + \alpha^2 n(n-1) u_k.$$

Moreover it can be verified that

$$(2.13) \quad \begin{aligned} -\gamma \Delta u_k + R(P) u_k = \frac{n(n-1)}{|v_k|_2^*} \left[\left(\frac{1}{k} + \frac{1 - \cos \alpha r}{\alpha^2} \right)^{-1-n/2} \cdot \right. \\ \left. \cdot \frac{2}{k} \left(1 + \frac{\alpha^2}{2k} \right) - \alpha^2 \left(\frac{1}{k} + \frac{1 - \cos \alpha \delta}{\alpha^2} \right)^{1-n/2} \right] \end{aligned}$$

using (A.10) in the Appendix

$$|v_k|_2^{4/(n-2)} = k \left[2^{-1+n/2} \omega_{n-1} I_n^{-1+n/2} + O \left(\frac{1}{k} \right) \right]^{2/n}.$$

Moreover

$$\omega_n = 2^{n-1} \omega_{n-1} I_n^{-1+n/2}.$$

where

$$I_n^{-1+n/2} = \int_0^{+\infty} (1+t)^{-n} t^{-1+n/2} dt.$$

Then

$$(2.14) \quad \frac{2}{k|v_k|_{2^*}} = \left(\frac{1}{|v_k|_{2^*}} \right)^{(n+2)/(n-2)} \cdot \left(\omega_n + O\left(\frac{1}{k}\right) \right)^{2/n}.$$

Inserting (2.14) into (2.13) and setting

$$v_k = \left(\frac{1}{k} + \frac{1 - \cos \alpha \delta}{\alpha^2} \right)^{1-n/2},$$

we get

$$(2.15) \quad \begin{aligned} Lu_k = n(n-1) \left(1 + \frac{\alpha^2}{2k} \right) \left(\omega_n + O\left(\frac{1}{k}\right) \right)^{2/n} & \left[u_k + \frac{v_k}{|v_k|_{2^*}} \right]^{(n+2)/(n-2)} + \\ & - \alpha^2 n(n-1) \frac{v_k}{|v_k|_{2^*}} \end{aligned}$$

by (2.15), (2.10), (2.11) and lemma 2.3 it is easy to verify that $\chi_k \rightarrow 0$ in $L^{2n/(n+2)} = (L^{2^*})' \hookrightarrow H^{-1}$; in fact

$$\begin{aligned} (R(x) - \alpha^2 n(n-1)) u_k &\rightarrow 0 \quad \text{in } L^2 \hookrightarrow L^{2n/(n+2)} \\ v_k/|v_k|_{2^*} &\rightarrow 0 \quad \text{in } L^\infty \text{ by (A.10)} \end{aligned}$$

and also

$$\begin{aligned} \int_M [(u_k + v_k)^{(n+1)/(n-2)} - u_k^{(n+1)/(n-2)}]^{2n/(n+2)} dM &\rightarrow 0, \\ \int_M \left| u'_k \cdot \frac{\partial}{\partial r} \left(\lg \left(\frac{\sin \alpha r}{r} \right)^{n-1} - \lg \sqrt{|g|} \right) \right|^2 dM &\rightarrow 0 \end{aligned}$$

Q.E.D.

Finally we can argue in an analogous manner if $R(P) < 0$ for any P .

PROOF OF THEOREM 0.2. We shall prove that for any $\varepsilon > 0$ the functional

$$\psi_\varepsilon(u) = \int_M [\gamma |\nabla u|^2 + (R(x) - \varepsilon) u^2] dM$$

has a minimum on the manifold

$$V = \{u \in H^1: |u|_{2^*} = 1\}.$$

By theorem 0.1 (see also remark 1.2) we need only to show that

$$(2.16) \quad \mu = \inf \psi_\varepsilon|_V < \gamma S.$$

If $R(x) \leq 0$ for any $x \in M$, (2.16) is obviously satisfied.

Suppose now that there exists $P \in M$ such that $R(P) > 0$ and consider the sequence $\{u_k\}$ defined by (2.6).

By lemma 2.3 we have

$$\psi_\varepsilon(u_k) = \psi(u_k) - \varepsilon |u_k|_2^2 = \begin{cases} \gamma S - \varepsilon \frac{\gamma}{2} \omega_n^{2/n} \frac{1}{k} + o\left(\frac{1}{k}\right) & \text{if } n \geq 5, \\ \gamma S - \varepsilon \frac{\gamma}{2} \omega^{\frac{1}{2}} \frac{\lg k}{k} + o(k^{-1} \lg k) & \text{if } n = 4. \end{cases}$$

Then we can choose $k \in \mathbf{N}$ s.t.

$$\inf \psi_\varepsilon|_V \leq \psi_\varepsilon(u_k) < \gamma S \quad \text{Q.E.D. .}$$

Appendix.

In proving Theorem 2.1 we have used the following result:

PROPOSITION A.1. *Let f be a C^1 functional on an Hilbert manifold V . Suppose that f is bounded from below and that it satisfies the Palais-Smale condition at $\mu = \inf f$, in the following sense:*

If $\{u_j\} \subset V$ s.t. as $j \rightarrow \infty$

$$(i) \quad df(u_j) \rightarrow 0$$

$$(ii) \quad f(u_j) \rightarrow \mu$$

then $\{u_j\}$ is precompact. Under the above assumption for every minimizing sequence $\{w_n\}$ for f we have $df(w_n) \rightarrow 0$.

The proof of proposition A.1 is deduced by following standard variational arguments. For completeness we shall give a sketch of the proof.

Since f satisfies the (P-S) condition at μ , the set

$$K_\mu = \{u \in V : f(u) = \mu, df(u) = 0\}$$

is compact.

Let $k \in \mathbf{N}$ and consider the neighborhood of K_μ defined by

$$N_k = \left\{ x \in V : d(x, K_\mu) < \frac{1}{k} \right\}.$$

In correspondence of N_k , there exists $\varepsilon > 0$ and an homeomorphism $\eta: V \rightarrow V$ s.t.

$$\eta(f^{-1}(]-\infty, \mu + \varepsilon]) \setminus N_k) \subset f^{-1}(]-\infty, \mu - \varepsilon]) = \emptyset.$$

Then

$$(A.0) \quad f^{-1}(]-\infty, \mu + \varepsilon]) \subset N_k.$$

Now consider $\{\omega_n\} \subset V$ s.t.

$$f(\omega_n) \rightarrow \mu.$$

Then by (A.0) there exists n_k s.t. for $n > n_k$

$$\omega_n \in N_k.$$

Therefore, since K_μ is compact there exists subsequence $\{\omega_{n_k}\}$ converging to $u \in K_\mu$, then $df(\omega_{n_k}) \rightarrow 0$. Q.E.D.

Let us give a sketch of the proof of Lemma 2.3. Set

$$S(r) = \{x \in M : d(x, P) = r\} = \partial B(P, r)$$

Then (cp. [1, lemma 1]) we have

$$(A.1) \quad \int_{S(r)} \sqrt{|g|} d\Omega = \omega_{n-1} \left\{ \left[1 + a \left(\frac{1 - \cos \alpha r}{\alpha^2} \right)^2 \right] \left(\frac{\sin \alpha r}{\alpha r} \right)^{n-1} + o(r^5) \right\}$$

where a is a suitable constant depending on the metric. Set

$$(A.2) \quad I_p^a = \int_0^{+\infty} (1+t)^{-p} t^a dt,$$

$$(A.3) \quad \nu_k = \left(\frac{1 - \cos \alpha \delta}{\alpha^2} + \frac{1}{k} \right)^{1-n/2}.$$

Then

$$(A.4) \quad |v_k|_2^2 = \int_0^\delta \left[\left(\frac{1}{k} + \frac{1 - \cos \alpha r}{\alpha^2} \right)^{1-n/2} - \nu_k \right]^2 r^{n-1} \int_{S(r)} \sqrt{|g|} d\Omega.$$

Inserting (A.1) in (A.4) we have

$$(A.5) \quad |v_k|_2^2 = \omega_{n-1} \int_0^\delta \left\{ \left[\left(\frac{1}{k} + t \right)^{1-n/2} - \nu_k \right]^2 \cdot (2t)^{(n-2)/2} \left(1 - \frac{\alpha^2 t}{2} \right)^{(n-2)/2} (1 + at)^2 + O(t^{(4+n)/2}) \right\} dt$$

where

$$t = \frac{1 - \cos \alpha r}{\alpha^2}, \quad \varrho = \frac{1 - \cos \alpha \delta}{\alpha^2}.$$

Then by using the Taylor formula for the factor

$$(2t)^{(n-2)/2} \left(1 - \frac{\alpha^2 t}{2} \right)^{(n-2)/2} (1 + at)^2$$

and observing that for p and q positives

$$\int_0^\varrho \left(\frac{1}{k} + t \right)^{-p} t^q dt = \begin{cases} I_p^a k^{p-q-1} + O(1) & \text{if } p - q - 1 > 0, \\ \log k + O(1) & \text{if } p - q - 1 = 0, \\ O(1) & \text{if } p - q - 1 < 0, \end{cases}$$

we get

$$(A.6) \quad |v_k|_2^2 = \begin{cases} 2^{(n-2)2} \omega_{n-1} [k^{(n-4)/2} I_{n-2}^{(n-2)/2} + \sigma_n^1] & \text{if } n \geq 5, \\ 2^{(n-2)/2} \omega_{n-1} \{\log k + O(1)\} & \text{if } n = 4, \\ O(1) & \text{if } n = 3, \end{cases}$$

where

$$(A.7) \quad \sigma_n^1 = \begin{cases} \frac{2-n}{4} \alpha^2 k^{(n-6)/2} I_{n-2}^{n/2} + o(k^{(n-6)/2}) & \text{if } n > 6, \\ \frac{2-n}{4} \alpha^2 \log k + o(\log k) & \text{if } n = 6, \\ O(1) & \text{if } n = 5. \end{cases}$$

Moreover it can be verified that the following asymptotic expansions hold:

$$(A.8) \quad |\nabla v_k|_2^2 = \left(\frac{n-2}{2}\right)^2 2^{n/2} \omega_{n-1} [I_n^{n/2} k^{(n-2)/2} + \sigma_n^2]$$

where

$$(A.9) \quad \sigma_n^2 = \begin{cases} -\frac{n\alpha^2}{4} \frac{n+2}{n-4} I_n^{n/2} k^{n/2-2} + o(k^{(n-4)/2}) & \text{if } n \geq 5, \\ -\frac{n\alpha^2}{4} \log k + O(1) & \text{if } n = 4, \\ O(1) & \text{if } n = 3, \end{cases}$$

$$(A.10) \quad |v_k|_{2^*}^{2^*} = 2^{(n-2)/2} \omega_{n-1} k^{n/2} I_n^{(n-2)/2} \left(1 - \frac{n\alpha^2}{4k}\right) + \sigma_n^3$$

where

$$(A.11) \quad \sigma_n^3 = \begin{cases} O(k^{n/2-2}) & \text{if } n \geq 6, \\ O(k) & \text{if } n = 5, \\ -4k\nu_k I_3^1 + O(\log k) & \text{if } n = 4, \\ -6k\nu_k I_{5/2}^{1/2} + O(\log k) & \text{if } n = 3, \end{cases}$$

$$(A.12) \quad \int_M Rv_k^2 dM = \begin{cases} \omega_{n-1} 2^{(n+2)/2} \frac{(n-2)(n-1)^2}{n-4} \alpha^2 I_n^{n/2} k^{(n-4)/2} + \sigma_n^4 & \text{if } n \geq 5, \\ \omega_3 2^3 \cdot 3 \cdot \alpha^2 \log k + O(1) & \text{if } n = 4, \\ O(1) & \text{if } n = 3, \end{cases}$$

where

$$(A.13) \quad \sigma_n^4 = \begin{cases} O(k^{(n-6)/2}) & \text{if } n > 6, \\ O(\log k) & \text{if } n = 6, \\ O(1) & \text{if } n = 5. \end{cases}$$

Moreover

$$(A.14) \quad \int_M r^2 |\nabla v_k|^2 dM = \begin{cases} 2^{(n+2)/2} \left(\frac{n-2}{2}\right)^2 \omega_{n-1} I_n^{(n+2)/2} k^{(n-4)/2} + \sigma_k^5 & \text{if } n \geq 5, \\ 2^3 \omega_3 \log k + O(1) & \text{if } n = 4, \\ O(1) & \text{if } n = 3, \end{cases}$$

where

$$(A.15) \quad \sigma_k^5 = \begin{cases} O(k^{(n-6)/2}) & \text{if } n > 6, \\ O(\log k) & \text{if } n = 6, \\ O(1) & \text{if } n = 5. \end{cases}$$

Then, since $u_k = v_k/|v_k|_2$, the asymptotic expansions (2.7), (2.9), are easily derived. Q.E.D.

Notes added in proofs. After submission of this paper we have known that R. Schoen (*Conformal deformation of a Riemannian metric to constant scalar curvature*, preprint) has proved that for a manifold conformally different from S^n , the conformal invariant μ is (strictly) less than γS ; then the Yamabe conjecture is positively solved.

REFERENCES

- [1] T. AUBIN, *Equations différentielles non linéaires et Problème de Yamabe concernant la courbure scalaire*, J. Math. Pure Appl., **55** (1976), pp. 269-296.
- [2] T. AUBIN, *Espace de Sobolev sur le variétés Riemanniennes*, Bull. Sc. Math., **100** (1976), pp. 149-173.
- [3] T. AUBIN, *Nonlinear Analysis on Manifolds. Monge-Ampère Equations*, Springer-Verlag (1982).

- [4] H. BREZIS - L. NIRENBERG, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math., **36** (1983), pp. 437-477.
- [5] G. CERAMI - D. FORTUNATO - M. STRUWE, *Bifurcation and multiplicity results for nonlinear elliptic problems involving critical Sobolev exponents*, Ann. Inst. H. Poincaré - Nonlinear Analysis, (1984), pp. 341-350.
- [6] R. PALAIS, *Ljusternik-Schnirelman theory on Banach manifold*, Topology, **5** (1966), pp. 115-132.
- [7] P. H. RABINOWITZ, *Variational methods for nonlinear eigenvalue problems*, Proceedings of C.I.M.E. (1974), pp. 139-195, G. Prodi Editor.
- [8] G. TALENTI, *Best constant in Sobolev inequality*, Ann. Math. Pura Appl., **110** (1976), pp. 353-372.
- [9] N. TRÜDINGER, *Remarks concerning the conformal deformation of Riemannian structures on compact manifolds*, Ann. Sc. Norm. Sup. Pisa, **22** (1968), pp. 265-274.
- [10] M. VAUGON, *Equations différentielles non linéaires sur les variétés riemanniennes compactes*, Bull. Sci. Math., II Sér. 107 (1983) pp. 371-391.
- [11] H. YAMABE, *On the deformation of Riemannian structures on compact manifolds*, Osaka Math. J., **12** (1960), pp. 21-37.

Manoscritto pervenuto in redazione il 10 aprile 1985.