



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Partial differential equations/Numerical analysis

On steady-state preserving spectral methods for homogeneous Boltzmann equations



Sur des méthodes spectrales préservant les équilibres de l'équation de Boltzmann homogène

Francis Filbet^a, Lorenzo Pareschi^b, Thomas Rey^c^a Université Lyon-1 & Inria, Institut Camille-Jordan 43, boulevard du 11-Novembre-1918, 69622 Villeurbanne cedex, France^b Mathematics and Computer Science Department, University of Ferrara, Italy^c Center of Scientific Computation and Mathematical Modeling (CSCAMM), The University of Maryland, College Park, MD 20742-4015, USA

ARTICLE INFO

Article history:

Received 8 August 2014

Accepted after revision 23 January 2015

Available online 13 February 2015

Presented by Olivier Pironneau

ABSTRACT

In this note, we present a general way to construct spectral methods for the collision operator of the Boltzmann equation that preserves exactly the Maxwellian steady state of the system. We show that the resulting method is able to approximate with spectral accuracy the solution uniformly in time.

© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Dans cette note, nous présentons une construction générale de méthodes spectrales pour l'opérateur de collision de l'équation de Boltzmann permettant de préserver exactement les états stationnaires maxwelliens de ce type d'équations. Cette nouvelle approche est basée sur une décomposition de type «micro-macro» de la solution de l'équation, tout en restant très proche d'une méthode spectrale plus classique. Nous montrons que les méthodes obtenues sont capables d'approcher avec une précision spectrale, uniformément en temps, la solution de l'équation considérée, et nous présentons leur efficacité dans un test numérique.

© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

Spectral methods for the numerical solution to the homogeneous Boltzmann equation have been proposed originally in [13] and [14]. Recently, their stability properties were studied in [5]. Related approaches, based on the use of the Fourier transform, have been introduced by [1] and [7].

In classical kinetic theory, the behavior of a rarefied gas of particles is described by the Boltzmann equation [3]:

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f = Q(f, f)$$

E-mail addresses: filbet@math.univ-lyon1.fr (F. Filbet), lorenzo.pareschi@unife.it (L. Pareschi), trey@cscamm.umd.edu (T. Rey).

<http://dx.doi.org/10.1016/j.crma.2015.01.015>

1631-073X/© 2015 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

where $f(t, x, v)$, $x \in \mathbb{R}^d$, $v \in \mathbb{R}^d$ ($d \geq 2$), is the time-dependent particle distribution function in the phase space and the collision operator \mathcal{Q} is defined by

$$\mathcal{Q}(f, f)(v) = \int_{\mathbb{R}^d} \int_{\mathbb{S}^{d-1}} B(\cos \theta, |v - v_*|) [f'_* f' - f_* f] d\omega dv_*. \quad (1)$$

Time and position act only as parameters in \mathcal{Q} and therefore will be omitted in its description. In (1) we used the short-hands $f = f(v)$, $f_* = f(v_*)$, $f' = f(v')$, $f'_* = f(v'_*)$. The velocities of the colliding pairs (v, v_*) and (v', v'_*) are related by

$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2} \sigma, \quad v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2} \sigma.$$

The collision kernel B is a non-negative function that only depends on $|v - v_*|$ and $\cos \theta = ((v - v_*) / |v - v_*|) \cdot \omega$.

In this note, we shall assume that the system considered is space homogeneous and then the particle distribution function f will only depend on time t and velocity v . The Boltzmann equation then becomes

$$\frac{\partial f}{\partial t} = \mathcal{Q}(f, f). \quad (2)$$

Boltzmann's collision operator has the fundamental properties of conserving mass, momentum and energy

$$\int_{v \in \mathbb{R}^d} \mathcal{Q}(f, f) \phi(v) dv = 0, \quad \phi(v) = 1, v, |v|^2$$

and satisfies the well-known Boltzmann's H -theorem

$$-\frac{d}{dt} \int_{v \in \mathbb{R}^d} f \log f dv = - \int_{\mathbb{R}^d} \mathcal{Q}(f, f) \log(f) dv \geq 0.$$

The functional $-\int f \log f$ is the entropy of the solution. Boltzmann H -theorem implies that any equilibrium distribution function has the form of a locally Maxwellian distribution

$$M(\rho, u, T)(v) = \frac{\rho}{(2\pi T)^{d/2}} \exp\left(-\frac{|u - v|^2}{2T}\right),$$

where ρ, u, T are the density, mean velocity and temperature of the gas:

$$\rho = \int_{\mathbb{R}^d} f(v) dv, \quad u = \frac{1}{\rho} \int_{\mathbb{R}^d} v f(v) dv, \quad T = \frac{1}{d\rho} \int_{\mathbb{R}^d} |u - v|^2 f(v) dv.$$

A major drawback of the classical spectral method is the lack of exact conservations and, as a consequence, the incapacity of the scheme to preserve the Maxwellian steady states of the system. In this paper, we overcome this drawback thanks to a new reformulation of the method that permits to preserve the spectral accuracy and to capture the long-time behavior of the system.

2. A steady-state-preserving spectral method

Although the method applies in principle to any collisional kinetic equation that possesses a local Maxwellian-type equilibrium, here we describe the method in the case of the Boltzmann operator.

2.1. A micro-macro decomposition

Let us start with the decomposition

$$f = M + g, \quad (3)$$

with M the local Maxwellian equilibrium and g such that $\int_{\mathbb{R}^d} g \phi dv = 0$, $\phi = 1, v, |v|^2$. When inserted into a Boltzmann-type collision operator, the decomposition (3) gives

$$\mathcal{Q}(f, f) = \mathcal{L}(M, g) + \mathcal{Q}(g, g), \quad (4)$$

where $\mathcal{L}(M, g) = \mathcal{Q}(g, M) + \mathcal{Q}(M, g)$ is a linear operator, and we used the fact that

$$\mathcal{Q}(M, M) = 0. \quad (5)$$

There are two major features in the decomposition (4):

- (1) it embeds the identity (5);
- (2) the steady state of (4) is given by $g = 0$.

This type of micro–macro decomposition has been used, e.g., in [9,11] to develop numerical methods that preserve asymptotic behaviors of some kinetic models. Reviews of such methods can be found in [8,4].

To illustrate our method, let us consider now the space homogeneous equation (2) that we rewrite using the micro–macro decomposition as

$$\begin{cases} \frac{\partial g}{\partial t} = \mathcal{L}(M, g) + \mathcal{Q}(g, g), \\ f = M + g. \end{cases} \tag{6}$$

2.2. Derivation of the spectral method

To simplify the notations, we derived the spectral method in the classical setting introduced in [14]; similarly, it can be extended to the representation used in [12] for the derivation of fast algorithms. Thus, we perform the usual periodization in a bounded domain of the operators \mathcal{L} and \mathcal{Q} and denote by \mathcal{L}^λ and \mathcal{Q}^λ the operators with cut-off on the relative velocity on $\mathcal{B}_0(2\lambda\pi)$.

Let us first set up the mathematical framework of our analysis. For any $t \geq 0$, $f_N(v, t)$ is a trigonometric polynomial of degree N in v , i.e. $f_N(t) \in \mathbb{P}^N$ where

$$\mathbb{P}^N = \text{span} \left\{ e^{ik \cdot v} \mid -N \leq k_j \leq N, j = 1, \dots, d \right\}.$$

Moreover, let $\mathcal{P}_N : L^2([-\pi, \pi]^3) \rightarrow \mathbb{P}^N$ be the orthogonal projection upon \mathbb{P}^N in the inner product of $L^2([-\pi, \pi]^3)$

$$\langle f - \mathcal{P}_N f, \phi \rangle = 0, \quad \forall \phi \in \mathbb{P}^N.$$

We denote the L^2 -norm by

$$\|f\|_2 = (\langle f, f \rangle)^{1/2}.$$

With this definition, $\mathcal{P}_N f = f_N$, where f_N is the truncated Fourier series of f given by

$$f_N(v) = \sum_{k=-N}^N \hat{f}_k e^{ik \cdot v},$$

with

$$\hat{f}_k = \frac{1}{(2\pi)^d} \int_{[-\pi, \pi]^d} f(v) e^{-ik \cdot v} dv.$$

We then write the Fourier–Galerkin approximation of the micro–macro equation (6):

$$\begin{cases} \frac{\partial g_N}{\partial t} = \mathcal{L}_N^\lambda(M_N, g_N) + \mathcal{Q}_N^\lambda(g_N, g_N), \\ f_N = M_N + g_N, \end{cases} \tag{7}$$

where $M_N = \mathcal{P}_N M$, $g_N = \mathcal{P}_N g$, $\mathcal{L}_N^\lambda(M_N, g_N) = \mathcal{P}_N \mathcal{L}^\lambda(M_N, g_N)$ and $\mathcal{Q}_N^\lambda(g_N, g_N) = \mathcal{P}_N \mathcal{Q}^\lambda(g_N, g_N)$. More precisely, using multi-index notations in the above expressions we have

$$\mathcal{Q}_N^\lambda(g_N, g_N) = \sum_{k=-N}^N \left(\sum_{\substack{l, m=-N \\ l+m=k}}^N \hat{g}_l \hat{g}_m \hat{\beta}(l, m) \right) e^{ik \cdot v}, \quad k = -N, \dots, N, \tag{8}$$

where the Boltzmann kernel modes $\hat{\beta}(l, m) = \hat{B}(l, m) - \hat{B}(m, m)$ are given by

$$\hat{B}(l, m) = \int_{\mathcal{B}_0(2\lambda\pi)} \int_{\mathbb{S}^{d-1}} B(\cos \theta, |q|) e^{-i(l \cdot q^+ + m \cdot q^-)} d\omega dq. \tag{9}$$

In this last identity, $q = v - v_*$ is the relative velocity and q^+, q^- are the following parameterizations of the post-collisional velocities:

$$q^+ = \frac{1}{2}(q + |q|\omega), \quad q^- = \frac{1}{2}(q - |q|\omega).$$

It is immediate to show that

Proposition 1. *The function $g_N \equiv 0$ is an admissible local equilibrium of the scheme (7) and therefore $f_N = M_N$ is a local equilibrium state.*

3. Spectral accuracy

In this section, we show that (7) is a spectrally accurate approximation to (6), provided that M_N is a spectrally accurate approximation of M . This is clearly guaranteed if initially the support is large enough. Note that due to space homogeneity, M does not change in time and so does M_N . It is interesting to observe that the only difference between scheme (7) and the usual spectral method developed on the original formulation

$$\begin{aligned} \frac{\partial f_N}{\partial t} &= \frac{\partial g_N}{\partial t} \\ &= \mathcal{Q}_N^\lambda(f_N, f_N) \\ &= \mathcal{L}_N^\lambda(M_N, g_N) + \mathcal{Q}_N^\lambda(g_N, g_N) + \mathcal{Q}_N^\lambda(M_N, M_N) \end{aligned} \quad (10)$$

is due to the constant (in time) term

$$\mathcal{Q}_N^\lambda(M_N, M_N) \neq 0, \quad (11)$$

which, as we will prove in the sequel, is spectrally small and is not present in (7).

If $f \in H_p^r([- \pi, \pi]^d)$, where $r \geq 0$ is an integer and $H_p^r([- \pi, \pi]^d)$ is the subspace of the Sobolev space $H^r([- \pi, \pi]^d)$, which consists of periodic functions, we have the following estimate for the spectral accuracy of the classical spectral method [14].

Theorem 1. *Let $f \in H_p^r([- \pi, \pi]^d)$, $r \geq 0$ then*

$$\|\mathcal{Q}^\lambda(f, f) - \mathcal{Q}_N^\lambda(f_N, f_N)\|_2 \leq \frac{C}{N^r} \left(\|f\|_{H_p^r} + \|\mathcal{Q}^\lambda(f_N, f_N)\|_{H_p^r} \right), \quad (12)$$

Therefore we have the following result for (11):

Lemma 1. *Let $M_N = \mathcal{P}_N M$ where M is a local Maxwellian. Then*

$$\|\mathcal{Q}_N^\lambda(M_N, M_N)\|_2 \leq \frac{C}{N^r} \left(\|M\|_{H_p^r} + \|\mathcal{Q}^\lambda(M_N, M_N)\|_{H_p^r} \right), \quad (13)$$

Merging the two results we obtain the spectral accuracy of the new steady-state formulation

Theorem 2. *Let $f \in H_p^r([- \pi, \pi]^3)$, $r \geq 0$ then*

$$\|\mathcal{Q}^\lambda(f, f) - \mathcal{L}_N^\lambda(M_N, g_N) - \mathcal{Q}_N^\lambda(g_N, g_N)\|_2 \leq \frac{C}{N^r} \left(\|f\|_{H_p^r} + \|M\|_{H_p^r} + \|\mathcal{Q}^\lambda(f_N, f_N)\|_{H_p^r} + \|\mathcal{Q}^\lambda(M_N, M_N)\|_{H_p^r} \right). \quad (14)$$

4. A numerical example

We present in this section a numerical example of our method for the space homogeneous Boltzmann equation in dimension 2, with Maxwell molecules:

$$B(\cos \theta, |v - v_*|) = \frac{1}{2\pi}.$$

We compare the classical spectral method to the new steady-state preserving one. For this, we use an exact solution to the homogeneous Boltzmann equation, the so called Bobylev–Krook–Wu solution [2,10]. It is given by

$$f_{\text{BKW}}(t, v) = \frac{\exp(-v^2/2S)}{2\pi S^2} \left[2S - 1 + \frac{1-S}{2S} v^2 \right]$$

with $S = S(t) = 1 - \exp(-t/8)/2$.

For the resolution of the Boltzmann equation (2), we use the fast spectral method of [6], with $N = 32$ half-modes in each direction of the box $[-V, V]^2$ for $V = 8$. We take $M = 8$ angular discretizations. Here, we insist on the fact that both




Fig. 1. (Color online.) Comparison of the classical spectral method with the steady-state preserving one: Relative entropy $\mathcal{H}(f|M)(t)$ (left) and L^2 error $\|f(t) - f_{\text{BKW}}(t)\|_2$ (right).




Fig. 2. (Color online.) Comparison of the classical spectral method with the steady-state-preserving one: Evolution of the temperature.

the classical and steady-state-preserving methods rely on the same numerical algorithm. The only difference between the two is the presence of the constant-over-time term (11). In particular, both methods have the same computational cost, namely $\mathcal{O}(MN \log_2 N)$.

Fig. 1 presents a comparison between both methods for the relative entropy of the solution f with respect to the global Maxwellian M

$$\mathcal{H}(f|M)(t) := \int_{\mathbb{R}^d} f(t, v) \log \left(\frac{f(t, v)}{M(v)} \right),$$

and the absolute L^2 error between the numerical solution $f(t, v)$ and the exact one $f_{\text{BKW}}(t, v)$. We observe that for both quantities, the behavior of the steady-state-preserving method is better than the classical one. In particular, the new method achieves a nice monotonous decay of the relative entropy, without the large time increase of the classical spectral method. This can be explained thanks to the fact that the equilibria of the classical spectral method are constants over the computational domain [5]. The behavior of the L^2 error is also monotonous for the steady-state preserving method, which is not the case for the classical one.

We then present in Fig. 2 a comparison between the temperatures of the solutions obtained with the classical and steady-state-preserving methods. As expected with such a few Fourier mode, the temperature of the classical spectral method is not preserved, but grows almost linearly because of the convergence toward the discrete equilibrium, a constant on the cube. In contrast, it seems that the steady-state-preserving spectral methods improve greatly this behavior: although not perfectly conserved, the temperature of the solution decays by roughly 0.1% and then stabilizes at a constant value.

Finally, as a last observation, we noticed that even if the positivity of the solution is not preserved by the new method, the number of nonpositive cells seems to be lower at a fixed time for the steady-state-preserving method, compared to the classical one. We recall that it was already proved in [5] that the number of nonpositive cells is asymptotically 0 for the classical spectral method (Theorem 3).

Theorem 3. (See Filbet, Mouhot [5].) Let f_N be the unique, global solution to the truncated, Fourier transformed Boltzmann equation, associated with a nonnegative initial datum $f_0 \in H^k_p([-\pi, \pi]^d)$ with $k > d/2$. Then the mass of the negative values of f_N can be made uniformly in time small as the number of Fourier modes $N \rightarrow \infty$.

Acknowledgement

The research of FF is partially supported by the Inria EPI Kaliffe and European Research Council ERC Starting Grant 2009, project 239983–NuSiKiMo. The research of TR is granted by the NSF Grants #1008397 and #1107444 (KI-Net) and ONR grant #000141210318. Part of this work was conducted during a KI-Net (NSF #1107444) meeting in North Carolina State University.

References

- [1] A. Bobylev, S. Rjasanow, Fast deterministic method of solving the Boltzmann equation for hard spheres, *Eur. J. Mech. B, Fluids* 18 (1999) 869–887.
- [2] A. Bobylev, Exact solutions of the Boltzmann equation, *Dokl. Akad. Nauk SSSR* 225 (1975) 1296–1299.
- [3] C. Cercignani, R. Illner, M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Springer-Verlag, 1994.
- [4] G. Dimarco, L. Pareschi, Numerical methods for kinetic equations, *Acta Numer.* 23 (2014) 369–520.
- [5] F. Filbet, C. Mouhot, Analysis of spectral methods for the homogeneous Boltzmann equation, *Trans. Amer. Math. Soc.* 363 (2011) 1947–1980.
- [6] F. Filbet, C. Mouhot, L. Pareschi, Solving the Boltzmann equation in $N \log_2 N$, *SIAM J. Sci. Comput.* 28 (3) (2007) 1029–1053.
- [7] I. Gamba, S. Tharkabhushanam, Spectral-Lagrangian methods for collisional models of non-equilibrium statistical states, *J. Comput. Phys.* 228 (6) (2009) 2012–2036.
- [8] S. Jin, Asymptotic preserving (AP) schemes for multiscale kinetic and hyperbolic equations: a review, in: *Lecture Notes for Summer School on “Methods and Models of Kinetic Theory” (M&MKT)*, Porto Ercole (Grosseto, Italy), June, 2010; *S. Jin, Riv. Mat. Univ. Parma* 3 (2012) 177–216.
- [9] S. Jin, Y. Shi, A micro–macro decomposition-based asymptotic-preserving scheme for the multispecies Boltzmann equation, *SIAM J. Sci. Comput.* 31 (6) (2009–2010) 4580–4606.
- [10] M. Krook, T. Wu, Exact solutions of the Boltzmann equation, *Phys. Fluids* 20 (10) (1977) 1589–1595.
- [11] M. Lemou, L. Mieussens, A new asymptotic preserving scheme based on micro–macro formulation for linear kinetic equations in the diffusion limit, *SIAM J. Sci. Comput.* 31 (10) (2008) 334–368.
- [12] C. Mouhot, L. Pareschi, Fast methods for the Boltzmann collision integral, *C. R. Acad. Sci. Paris, Ser. I* 339 (1) (2004) 71–76.
- [13] L. Pareschi, B. Perthame, A spectral method for the homogeneous Boltzmann equation, *Transp. Theory Stat. Phys.* 25 (1996) 369–383.
- [14] L. Pareschi, G. Russo, Numerical solution of the Boltzmann equation I. Spectrally accurate approximation of the collision operator, *SIAM J. Numer. Anal.* 37 (2000) 1217–1245.