quatrième série - tome 49

fascicule 4 juillet-août 2016

ANNALES SCIENTIFIQUES de L'ÉCOLE NORMALE SUPÉRIEURE

Frédéric CAMPANA & Andreas HÖRING & Thomas PETERNELL Abundance for Kähler threefolds

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / Editor-in-chief

Antoine CHAMBERT-LOIR

Publication fondée en 1864 par Louis Pasteur	
Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE	
de 1883 à 1888 par H. DEBRAY	
de 1889 à 1900 par C. HERMITE	
de 1901 à 1917 par G. Darboux	
de 1918 à 1941 par É. PICARD	
de 1942 à 1967 par P. MONTEL	

Comité de rédaction au 1 ^{er} janvier 2016		
N. Anantharaman	I. GALLAGHER	
P. Bernard	B. Kleiner	
E. Breuillard	E. Kowalski	
R. Cerf	M. Mustață	
A. Chambert-Loir	L. SALOFF-COSTE	

Rédaction / Editor

Annales Scientifiques de l'École Normale Supérieure, 45, rue d'Ulm, 75230 Paris Cedex 05, France. Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80. annales@ens.fr

Édition / Publication

Abonnements / Subscriptions

Société Mathématique de France Institut Henri Poincaré 11, rue Pierre et Marie Curie 75231 Paris Cedex 05 Tél. : (33) 01 44 27 67 99 Fax : (33) 01 40 46 90 96 Maison de la SMF Case 916 - Luminy 13288 Marseille Cedex 09 Fax : (33) 04 91 41 17 51 email : smf@smf.univ-mrs.fr

Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

© 2016 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris). *All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

ABUNDANCE FOR KÄHLER THREEFOLDS

BY Frédéric CAMPANA, Andreas HÖRING and Thomas PETERNELL

ABSTRACT. – Let X be a compact Kähler threefold with terminal singularities such that K_X is nef. We prove that K_X is semiample, i.e., some multiple mK_X is generated by global sections.

RÉSUMÉ. – Soit X une variété kählérienne compacte à singularités terminales. Si K_X est nef, nous montrons que K_X est semi-ample, c'est-à-dire qu'un multiple mK_X est engendré par ses sections globales.

1. Introduction

1.A. Main results

Since the 1990's, the minimal model program for smooth complex projective threefolds is complete: every such manifold X admits a birational model X', which is \mathbb{Q} -factorial with only terminal singularities such that either

- X' carries a Fano fibration, in particular, X' is uniruled, or
- the canonical bundle $K_{X'}$ is semi-ample, i.e., some positive multiple $mK_{X'}$ is generated by global sections.

There are basically two parts in the program: first to establish the existence of a model X' which is either a Mori fibre space or has nef canonical divisor, and then to show that nefness implies semi-ampleness. This second part, known as "abundance", is established by [53, 42, 49].

The aim of the present paper is to fully establish the minimal model program in the category of Kähler threefolds. The first part of the program, i.e., the existence of a bimeromorphic model X' which either is a Mori fibre space or has nef canonical divisor, was carried out in the papers [38] and [37]. Thus it remains to show that nefness of the canonical divisor implies semi-ampleness, i.e., abundance holds for Kähler threefolds:

THEOREM 1.1. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold with at most terminal singularities such that K_X is nef. Then K_X is semi-ample, that is some positive multiple mK_X is globally generated.

The paper [24] established the existence of some section in mK_X for non-algebraic minimal models, so the assumption above implies $\kappa(X) \geq 0$. In [60] abundance was shown for non-algebraic minimal models, excluding however the case when X has no non-constant meromorphic function. Our arguments do not use any information on the structure of X and work both for algebraic and non-algebraic Kähler threefolds.

As a corollary, we establish a longstanding conjecture [26] on Kähler threefolds:

THEOREM 1.2. – Let X be a smooth compact Kähler threefold. Assume that X is simple, i.e., there is no positive-dimensional proper subvariety through a very general point of X. Then there exists a bimeromorphic morphism $X \to T/G$ where T is a torus and G a finite group acting on T.

1.B. Outline of the paper

Let X be a normal compact \mathbb{Q} -factorial Kähler threefold X with only terminal singularities and nef canonical divisor K_X . Denote by $\kappa(X)$ its Kodaira dimension and by $\nu(X)$ the numerical dimension, which is defined as

$$\nu(X) := \max\{m \in \mathbb{N} \mid c_1(K_X)^m \neq 0\}.$$

Both invariants are subject to the inequality $\kappa(X) \leq \nu(X)$, with equality if K_X is semiample. Conversely, as Kawamata observed in [39, Thm. 6.1], in order to prove abundance, it is sufficient to prove equality: $\kappa(X) = \nu(X)$. By the base-point free theorem and an argument of Kawamata [39, Thm. 7.3] the main challenge is to rule out the possibility

$$\kappa(X) = 0 \text{ and } 0 < \nu(X) < 3.$$

Since we know that $\kappa(X) \ge 0$, there exists a positive number *m* and an effective divisor *D* such that $D \in |mK_X|$. A natural way to prove that $\kappa(X) \ge 1$ is to consider the restriction map

$$r: H^0(X, d(m+1)K_X) \to H^0(D, dK_D).$$

Arguing by induction on the dimension we want to prove that $H^0(D, dK_D) \neq 0$ for some $d \in \mathbb{N}$ and that some non-zero section $u \in H^0(D, dK_D)$ lifts via r to a global section $\tilde{u} \in H^0(X, d(m+1)K_X)$ on X. Since D might be very singular it is however not possible to analyse the divisor D directly. In order to circumvent this difficulty, Kawamata [42] developed the strategy, further explored in [49], to consider log pairs (X, B) with B = Supp D and to improve the singularities of this pair via certain birational transformations. This requires deep techniques of birational geometry of pairs within the theory of minimal models. In particular we have to run a log MMP for certain log pairs (X, Δ) .

Therefore the first part of the paper (Sections 3 and 4) establishes the foundations for a minimal model program for log pairs on Kähler threefolds. As a first step we prove the cone theorem for the dual Kähler cone:

4° SÉRIE – TOME 49 – 2016 – Nº 4

THEOREM 1.3. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold that is not uniruled. Let Δ be an effective \mathbb{Q} -divisor on X such that the pair (X, Δ) is dlt. Then there exists an at most countable family $(\Gamma_i)_{i \in I}$ of rational curves on X such that

$$0 < -(K_X + \Delta) \cdot \Gamma_i \le 4$$

and

$$\overline{NA}(X) = \overline{NA}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

The dual Kähler cone $\overline{NA}(X)$ replaces the Mori cone of curves $\overline{NE}(X)$ which is obviously too small in the non-algebraic setting (cf. [38, Sect. 1]). Our result is actually more general: first we prove a weak form of the cone theorem for lc pairs (X, Δ) such that X has rational singularities (cf. Theorem 4.1). Then we derive Theorem 1.3 on page 992 as a consequence of the weak cone theorem and some contraction results. The second step is to prove that the $(K_X + \Delta)$ -negative extremal rays appearing in the cone theorem can be contracted by a bimeromorphic morphism.

THEOREM 1.4. – Let X be a normal Q-factorial compact Kähler threefold that is not uniruled. Let Δ be an effective Q-divisor such that the pair (X, Δ) is dlt. Let R be a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$. If the ray R is divisorial with exceptional divisor S (cf. Definition 4.2), suppose that S has slc singularities.

Then the contraction of R exists in the Kähler category.

These two theorems generalize the analogous statements for terminal threefolds [38, Thm. 1.2, Thm. 1.3]. While the global strategy of the proofs is similar to [38], it is not an extension of earlier work: in this paper we have to deal with threefolds with non-isolated singularities. This leads to new geometric problems (cf. the proof of Theorem 4.2) and is more representative of the challenges that appear in higher dimension.

Based on these results we show in Section 6 how to replace the threefold X with some bimeromorphic model having a pluricanonical divisor $D \in |mK_X|$ such that D_{red} is not too singular. In this section we follow the arguments of [49, Ch. 13, 14], and we also address some technical points which do not appear in the literature. While these reduction steps assure the existence of some effective pluricanonical divisor on D_{red} , it is not obvious that it extends to a pluricanonical divisor on X (cf. [22] for recent progress in the projective case). Following a deformation argument of Miyaoka [54] [49, Ch. 11], the case $\kappa(X) = 0, \nu(X) = 1$ can be excluded without too much effort.

For the case $\nu(X) = 2$ the idea is to prove via a Riemann-Roch computation that $h^0(X, mK_X)$ grows linearly. Establishing the results necessary for this Riemann-Roch computation is the second part of the paper (Sections 7 and 8). First we extend Enoki's theorem [25] to show that the cotangent sheaf of a non-uniruled threefold with canonical singularities is generically nef, then we want to use this positivity result to prove that $(K_X + B) \cdot \hat{c}_2(X) \ge 0$ where $K_X + B$ is a nef log-canonical divisor and $\hat{c}_2(X)$ is the second Chern class of the Q-sheaf $\hat{\Omega}_X$. Since X is not smooth in codimension two, the proof of this inequality in the projective case ([42], [49, Ch. 14]) uses some involved computation for the second Todd class of the Q-sheaf $\hat{\Omega}_X$. We give a new argument which should be of

independent interest: let $X' \to X$ be a terminal modification of X. Since X' is smooth in codimension one, the Riemann-Roch computation on X' can be done using the classical second Chern class $c_2(X')$. Although the (log-)canonical divisor of X' is not necessarily nef, a generalization of Miyaoka's inequality for the second Chern class (cf. Theorem 7.2) allows us to conclude.

Acknowledgements. – We thank S. Boucksom and V. Lazić for some very helpful communications. We thank the Forschergruppe 790 "Classification of algebraic surfaces and compact complex manifolds" of the Deutsche Forschungsgemeinschaft for financial support. A. Höring was partially also supported by the A.N.R. project CLASS⁽¹⁾.

2. Notation and basic facts

We will use frequently standard terminology of the minimal model program (MMP) as explained in [51] or [21]. We will also use without further comment results which are stated for algebraic varieties in [51, 49] if their proof obviously works for complex spaces.

Recall that a normal complex space X is \mathbb{Q} -factorial if for every Weil divisor D there exists an integer $m \in \mathbb{N}$ such that $\mathcal{O}_X(mD)$ is a locally free sheaf, i.e., mD is a Cartier divisor. Since however the canonical sheaf $K_X = \omega_X$ need not be a \mathbb{Q} -Weil divisor, we include in the definition of \mathbb{Q} -factoriality also the condition that there is a number $m \in \mathbb{N}$ such that the coherent sheaf

$$(K_X^{\otimes m})^{**} = (\omega_X^{\otimes m})^{**}$$

is locally free. We shall write

$$mK_X = (K_X^{\otimes m})^*$$

for short.

DEFINITION 2.1. – Let X be a normal complex space. A boundary divisor is an effective \mathbb{Q} -divisor $\Delta = \sum_{i} a_i \Delta_i$ such that $0 \le a_i \le 1$.

Given a boundary divisor Δ , we refer to (X, Δ) as a "pair". For pairs (X, Δ) , the notions "lc" (log-canonical), "klt" (Kawamata log-terminal), "dlt" (divisorial log-terminal) and "terminal" are defined exactly as in the algebraic context; we refer to [51].

DEFINITION 2.2. – Let X be a normal Q-factorial compact Kähler space that is not uniruled, and let Δ be a boundary divisor on X such that the pair (X, Δ) is lc (resp. dlt, klt, terminal). Set $(X_0, \Delta_0) := (X, \Delta)$. A terminating $(K + \Delta)$ -MMP is a finite sequence of bimeromorphic maps

$$\varphi_i : (X_i, \Delta_i) \dashrightarrow (X_{i+1}, \Delta_{i+1} := (\varphi_i)_* \Delta_i)$$

where $i \in \{0, ..., n\}$ for some $n \in \mathbb{N}$ that has the following properties:

a) For every $i \in \{0, ..., n+1\}$, the complex space X_i is a normal \mathbb{Q} -factorial compact Kähler space and the pair (X_i, Δ_i) is lc (resp. dlt, klt, terminal).

(1) ANR-10-JCJC-0111

 $4^{\,e}\,S\acute{E}RIE-TOME\,49-2016-N^o\,4$

- b) For every $i \in \{0, ..., n\}$, the map φ_i is either the contraction of a $(K_{X_i} + \Delta_i)$ -negative extremal ray $R_i \in \overline{NA}(X_i)$ that is divisorial or the flip of a $K_{X_i} + \Delta_i$ -negative extremal ray $R_i \in \overline{NA}(X_i)$ that is small.
- c) The class $K_{X_{n+1}} + \Delta_{n+1}$ is nef.

We will abbreviate such a $(K + \Delta)$ *-MMP by*

$$(X, \Delta) \dashrightarrow (X', \Delta')$$

where $(X', \Delta') := (X_{n+1}, \Delta_{n+1})$ and the bimeromorphic map is the composition $\varphi_n \circ \cdots \circ \varphi_0$.

In the definition, $\overline{NA}(X)$ denotes the "dual Kähler cone", as defined in [38, 3.8]. We will review this notion in Definition 3.3 below.

2.A. Bimeromorphic models

Bimeromorphic models arising as partial resolutions of singularities play an important role in the recent development of the MMP for projective manifolds. The existence of these models is usually—in the algebraic setting, see [29, 10.4], [50, 3.1], [48, Ch. 1.4]—derived as a consequence of the existence and termination of some MMP for pairs, which is at this point not yet established in the Kähler category. We may nevertheless assume their existence since they are constructed by taking first a log-resolution $Y' \rightarrow X$ of the pair (X, Δ) and then running some MMP over X. Since we may choose the log-resolution to be a projective morphism, the usual relative versions of the cone and contraction theorem [56] [51, Ch. 3.6] together with termination results in dimension three allow to conclude. The Kähler property of Y follows again from [65, 1.3.1], since μ is a projective morphism. We will now apply this argument in two situations:

THEOREM 2.1. – Let X be a normal compact Kähler threefold with lc singularities, i.e., (X, 0)is lc. Then there exists a bimeromorphic morphism $\mu : Y \to X$ from a normal Q-factorial Kähler threefold Y with terminal singularities and a boundary divisor Δ_Y such that the pair (Y, Δ_Y) is lc and

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* K_X.$$

Proof. – Let $\pi : \hat{X} \to X$ be a resolution of singularities. Let $\hat{X} \dashrightarrow Y$ be a K_{\bullet} -MMP over X. The outcome of this MMP is a normal \mathbb{Q} -factorial threefold $\mu : Y \to X$ with terminal singularities such that K_Y is μ -nef. We write

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* K_X,$$

where the support of Δ_Y is contained in the μ -exceptional locus. By the negativity lemma [51, Lemma 3.39] we see that Δ_Y is effective and since X has lc singularities, Δ_Y is a boundary divisor. The pair (Y, Δ_Y) is lc since this property is invariant under crepant bimeromorphic morphisms [51, Lemma 2.30].

THEOREM 2.2. – Let X be a normal compact Kähler threefold, and let Δ be a boundary divisor on X such that the pair (X, Δ) is lc. Then (X, Δ) has a dlt model, i.e., there exists

a bimeromorphic morphism from a normal \mathbb{Q} -factorial Kähler threefold Y and a boundary divisor Δ_Y such that the pair (Y, Δ_Y) is dlt and

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^*(K_X + \Delta).$$

Proof. – The same reasons as in Theorem 2.1 apply also here; we refer to [50, 3.1] and [29, 10.4], based on [10, Thm. 1.2], which is a result on projective morphisms. The deduction from [10] is basically a computation of discrepancies including an application of the negativity lemma.

2.B. Adjunction

Let X be a normal complex space and let $S \subset X$ be a prime divisor that is \mathbb{Q} -Cartier. Let B be an effective \mathbb{Q} -divisor such that $K_X + B$ is \mathbb{Q} -Cartier and $S \not\subset \text{Supp}(B)$. Let $\nu : \tilde{S} \to S$ be the normalization, then Shokurov's different [62, Ch. 3] is a naturally defined effective \mathbb{Q} -divisor $B_{\tilde{S}}$ on \tilde{S} such that

(1)
$$K_{\tilde{S}} + B_{\tilde{S}} \sim_{\mathbb{Q}} \nu^* (K_X + S + B)|_{S}$$

From the construction in [62, Ch. 3] one sees immediately that

$$\operatorname{Supp} \nu_*(B_{\tilde{S}}) \subset S_{\operatorname{sing}} \cup (\operatorname{Supp} B \cap S) \cup (X_{\operatorname{sing}} \cap S)$$

Suppose now that dim X = 3 and let $\mu : \hat{S} \to \tilde{S}$ be the minimal resolution. Then we have

$$K_{\hat{S}} \sim_{\mathbb{Q}} \mu^* K_{\tilde{S}} - N_{\tilde{S}}$$

where N is an effective μ -exceptional divisor [61, 4.1]. Set $E := N + \mu^* B_{\tilde{S}}$, then E is a canonically defined effective \mathbb{Q} -divisor, such that

(2)
$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^* (K_X + S + B)|_{S},$$

where $\pi: \hat{S} \to S$ is the composition $\nu \circ \mu$. Since E is μ -exceptional, we obtain

$$\operatorname{Supp} \pi_*(E) \subset S_{\operatorname{sing}} \cup (\operatorname{Supp} B \cap S) \cup (X_{\operatorname{sing}} \cap S).$$

Let $C \subset S$ be a curve such that

$$C \not\subset S_{\operatorname{sing}} \cup (\operatorname{Supp} B \cap S) \cup (X_{\operatorname{sing}} \cap S)$$

Then the morphism π is an isomorphism in the generic point of C, and we can define the strict transform $\hat{C} \subset \hat{S}$ as the closure of $C \setminus S_{\text{sing}}$. By what precedes, $\hat{C} \not\subset E$, thus by the projection formula and (2), we deduce

(3)
$$K_{\hat{S}} \cdot \hat{C} \le (K_{\hat{S}} + E) \cdot \hat{C} = (K_X + S + B) \cdot C.$$

3. Singular Kähler spaces

3.A. Bott-Chern cohomology

In this section we review very briefly—following [38]—the cohomology groups that replace the Néron-Severi space $NS(X) \otimes \mathbb{R}$ from the projective setting. For details, we refer to [38] and the literature given there.

4° SÉRIE – TOME 49 – 2016 – Nº 4

We recall that $\mathscr{Q}_X^{p,q}$ denotes the sheaf of \mathscr{C}^{∞} -forms of type (p,q), and \mathscr{Q}_X^0 denotes the sheaf of \mathscr{C}^{∞} -functions; \mathscr{D}_X denotes the sheaf of distributions.

DEFINITION 3.1 ([14, Defn. 4.6.2]). – Let X be an irreducible reduced complex space. Let \mathcal{H}_X be the sheaf of real parts of holomorphic functions multiplied⁽²⁾ with i. A (1, 1)-form (resp. (1, 1)-current) with local potentials on X is a global section of the quotient sheaf $\mathcal{C}_X^0/\mathcal{H}_X$ (resp. $\mathcal{D}_X/\mathcal{H}_X$). We define the Bott-Chern cohomology

$$N^{1}(X) := H^{1,1}_{\mathrm{BC}}(X) := H^{1}(X, \mathcal{H}_{X}).$$

Using the exact sequence

$$0 \to \mathcal{H}_X \to \mathcal{A}_X^0 \to \mathcal{A}_X^0 / \mathcal{H}_X \to 0,$$

and the fact that \mathscr{Q}^0_X is acyclic, we obtain a surjective map

$$H^0(X, \mathscr{C}^0_X/\mathscr{H}_X) \to H^1(X, \mathscr{H}_X)$$

Thus we can see an element of the Bott-Chern cohomology group as a closed (1, 1)-form with local potentials modulo all the forms that are globally of the form $dd^{c}u$. Using the exact sequence

$$0 \to \mathcal{H}_X \to \mathcal{D}_X \to \mathcal{D}_X / \mathcal{H}_X \to 0,$$

we see that one obtains the same Bott-Chern group, if we consider (1, 1)-currents with local potentials.

Recall the following basic lemma [38, Lemma 3.3.].

LEMMA 3.1. – Let $\varphi : X \to Y$ be a proper bimeromorphic morphism between compact normal spaces in class \mathcal{C} with at most rational singularities. Then we have an injection

$$\varphi^*: H^1(Y, \mathcal{H}_Y) \hookrightarrow H^1(X, \mathcal{H}_X)$$

and

$$\operatorname{Im} \varphi^* = \{ \alpha \in H^1(X, \mathcal{H}_X) \mid \alpha \cdot C = 0 \quad \forall \ C \subset X \ curve \ s.t. \ \varphi(C) = pt \}.$$

Furthermore, let $\alpha \in H^1(X, \mathcal{H}_X) \subset H^2(X, \mathbb{R})$ be a class such that $\alpha = \varphi^*\beta$ with $\beta \in H^2(Y, \mathbb{R})$. Then there exists a smooth real closed (1, 1)-form with local potentials ω_Y on Y such that $\alpha = \varphi^*[\omega_Y]$.

Following [58], we define the notion of a pseudo-effective resp. nef class:

DEFINITION 3.2. – Let X be an irreducible reduced compact complex space and $u \in N^1(X)$.

- a) *u* is pseudo-effective if it can be represented by a current T which is locally of the form $T = i\partial\overline{\partial}\varphi$ with a psh function φ .
- b) u is nef if it can be represented by a form α with local potentials such that for some positive (1,1)-form ω on X and for all $\epsilon > 0$, there exists $f_{\epsilon} \in \mathcal{C}^{0}(X)$, such that

$$\alpha + i\partial\partial f_{\epsilon} \ge -\epsilon\omega.$$

⁽²⁾ We "twist" the definition from [14] in order to get a group that injects in $H^2(X, \mathbb{R})$ rather than $H^2(X, i\mathbb{R})$

DEFINITION 3.3. – Let X be a normal compact complex space in class C. We define $N_1(X)$ to be the vector space of real closed currents of bidimension (1, 1) modulo the following equivalence relation: $T_1 \equiv T_2$ if and only if

$$T_1(\eta) = T_2(\eta)$$

for all real closed (1,1)-forms η (with local potentials).

We furthermore define $\overline{NA}(X) \subset N_1(X)$ to be the closed cone generated by the classes of positive closed currents. The closed cone of curves is the subcone

$$\overline{NE}(X) \subset \overline{NA}(X)$$

of those positive closed currents arising as currents of integration over curves.

For the proof of the contraction Theorem 1.4 we will need the following statements, generalizing [38, Prop. 8.1].

PROPOSITION 3.1. – Let X be a normal Q-factorial compact Kähler space. Let Δ be a boundary divisor such that the pair (X, Δ) is dlt. Let $\mathbb{R}^+[\Gamma_i]$ be a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$. Suppose that there exists a bimeromorphic morphism $\varphi : X \to Y$ onto a normal complex space Y such that $-(K_X + \Delta)$ is φ -ample and a curve $C \subset X$ is contracted if and only if $[C] \in \mathbb{R}^+[\Gamma_i]$.

a) Then we have two exact sequences

(4)
$$0 \to H^2(Y, \mathbb{R}) \xrightarrow{\varphi^*} H^2(X, \mathbb{R}) \xrightarrow{\alpha \mapsto \alpha \cdot \Gamma_i} \mathbb{R} \to 0$$

(5)
$$0 \to N^1(Y) \xrightarrow{\varphi^*} N^1(X) \xrightarrow{[L] \mapsto L \cdot \Gamma_i} \mathbb{R} \to 0.$$

In particular we have $b_2(X) = b_2(Y) + 1$.

b) We have an exact sequence

(6)
$$0 \to \operatorname{Pic}(Y) \xrightarrow{\varphi^*} \operatorname{Pic}(X) \xrightarrow{[L] \mapsto L \cdot \Gamma_i} \mathbb{Z}.$$

- c) If the contraction is divisorial, the variety Y is \mathbb{Q} -factorial and its Picard number is $\rho(X) 1$. Moreover the pair $(Y, \varphi_* \Delta)$ is dlt
- d) If the contraction is small with flip $\varphi^+ : X^+ \to Y$, the complex space X^+ is \mathbb{Q} -factorial and its Picard number is $\rho(X)$. Moreover the pair $(X^+, (\varphi^+)^{-1}_* \circ \varphi_* \Delta)$ is dlt.

Proof. – Since $-(K_X + \Delta)$ is φ -ample, the morphism φ is projective. Since (X, Δ) is dlt and X is Q-factorial, the pair $(X, (1-\varepsilon)\Delta)$ is klt for small positive ε . Since $-(K_X+(1-\varepsilon)\Delta)$ is φ -ample, possibly after passing to a smaller ε , we obtain $R^q \varphi_*(\Theta_X) = 0$ for $q \ge 1$ by [2, Thm. 3.1]. Since X has rational singularities ([51, 5.22]), so does Y; moreover a) is proved by Lemma 3.1. The property b) follows from [51, Thm. 3.25(4)] and implies the properties c) and d) as in [51, Prop. 3.36, Prop. 3.37, Cor. 3.44].

LEMMA 3.2. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold, and let Δ be a boundary divisor such that the pair (X, Δ) is dlt. Let

$$\varphi: X \to Y \text{ resp. } \varphi: X \dashrightarrow X^+$$

be the divisorial contraction resp. the flip of a $K_X + \Delta$ -negative extremal ray $R \in \overline{NA}(X)$. Then the following hold:

4° SÉRIE – TOME 49 – 2016 – Nº 4

a) Let D be a nef divisor such that $D \cdot R = 0$. Then $\varphi_*(D)$ is nef and

$$\nu(D) = \nu(\varphi_*(D)).$$

Moreover if $S \subset X$ is a prime divisor that is not contracted by φ , then $D \cdot S \neq 0$ in $H^4(X, \mathbb{R})$ if and only if $\varphi_*(D) \cdot \varphi(S) \neq 0$ in $H^4(Y, \mathbb{R})$.

b) Let B be a boundary divisor such that the pair (X, B) is lc. If $(K_X + B) \cdot R = 0$, then the pair $(Y, \varphi_*(B))$ is lc.

Proof. – Set $L := \Theta_X(D)$. We first treat the case when φ is divisorial: by Proposition 3.1,b) we have $L \simeq \varphi^* M$ with M a line bundle on Y and obviously $M \simeq \Theta_Y(\varphi_* D)$. Being nef is invariant under pull-back [58] [38, Lemma 3.13], so M is nef. The numerical dimension is invariant since the pull-back commutes with the intersection product. If $S' = \varphi(S)$, then $L|_S = (\varphi|_S)^* (M|'_S)$, hence $L \cdot S \neq 0$ if and only if $M \cdot \varphi(S) \neq 0$ by the projection formula. For the property b) we refer to [51, Lemma 2.30].

In the flipping case, consider the small contraction $f : X \to Y$ associated to R and the flip $f^+ : X^+ \to Y$. For a) apply the considerations above first to f and then to f^+ . For b), apply [51, 2.30] to see that $(Y, f_*(B))$ is lc and then again to conclude that $(X^+, \varphi_*(B) = (f^+)_*^{-1} \circ f_*(B))$ is lc.

3.B. Kähler criteria

In this subsection we generalize some Kähler criteria given in [38] to threefolds with nonisolated singularities.

THEOREM 3.1. – Let X be a normal compact threefold in class C. Let $\eta \in \mathcal{A}^{1,1}(X)$ be a closed real (1,1)-form with local potentials such that $T(\eta) > 0$ for all $[T] \in \overline{NA}(X) \setminus 0$. Suppose that for every irreducible curve $C \subset X$ we have $[C] \neq 0$ in $N_1(X)$. Then $\{\eta\} \in N^1(X)$ is represented by a Kähler class, in particular X is Kähler.

Proof. – We slightly generalize the arguments of [38, Thm. 3.18] by removing the assumption that the singularities are isolated. If X is smooth, then [38, Thm. 3.18] applies, so that $\{\eta\}$ is a Kähler class.

We shall now reduce ourselves to this case by considering a resolution $\mu : \hat{X} \to X$ with \hat{X} a compact Kähler manifold such that a suitable exceptional divisor E of \hat{X} is μ -ample. As in [38], we argue that the class of an irreducible curve in \hat{X} does not vanish. Furthermore, we check as in Step 2 of the proof of [38, Thm. 3.18] that

$$T(\mu^*(\eta) - \frac{1}{n}E) > 0$$

for all classes $[T] \in \overline{NA}(\hat{X})$ if *n* is a sufficiently big positive integer. Hence, by the solution in the smooth case, we may choose a Kähler form $\hat{\eta}$ on \hat{X} in the class $\{\mu^*(\eta) - \frac{1}{n}E\}$. Consider the Kähler current

$$\{\mu_*(\hat{\eta})\} = \{\eta\}.$$

Let Z be an irreducible component of the Lelong level sets of $\mu_*(\hat{\eta})$; then Z is either a point or an irreducible curve. By our assumption, $\{\eta\}|Z$ is a Kähler class on Z. Note also that the

Kähler current $\mu_*(\hat{\eta})$ has local potentials, since by assumption η has local potentials. Thus we can apply [23, Prop. 3.3(iii)] to see that $\{\eta\}$ is a Kähler class on X.

COROLLARY 3.1. – Let X be a normal Q-factorial compact Kähler threefold that is not uniruled. Let Δ be an effective Q-divisor such that the pair (X, Δ) is dlt. Let $\mathbb{R}^+[\Gamma_i]$ be a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$.

Suppose that there exists a bimeromorphic morphism $\varphi : X \to Y$ such that $-(K_X + \Delta)$ is φ -ample and a curve $C \subset X$ is contracted if and only if $[C] \in \mathbb{R}^+[\Gamma_i]$. Then Y is a Kähler space.

Proof. – As in the proof of Proposition 3.1 we argue that Y has rational singularities. By assumption, $\mathbb{R}^+[\Gamma_i]$ is extremal in $\overline{NA}(X)$; denote by $\alpha \in N^1(X)$ a nef supporting class (cf. Proposition 4.3). Consequently

$$\alpha \cdot Z' > 0$$

for every class $Z' \in \overline{NA}(X) \setminus \mathbb{R}_0^+[\Gamma_i]$. By Proposition 3.1,a) there exists a $\beta \in N^1(Y)$ with $\alpha = \varphi^*(\beta)$ and by what precedes we have

$$\beta \cdot Z > 0$$

for any $Z \in \overline{\operatorname{NA}}(Y) \setminus 0$.

We claim that for any irreducible curve $C \subset Y$, there exists a curve $C' \subset X$ such that $C = \varphi(C')$. Since C' is not contracted by φ we have $\alpha \cdot C' > 0$, so $\beta \cdot C > 0$ by the projection formula. In particular, $[C] \neq 0$ in $N_1(X)$, so the statement follows from Theorem 3.1.

For the proof of the claim note first that if C is not contained in the image of the exceptional locus, then we can just take the strict transform $C' \subset X$. If C is in the image of the exceptional locus, then φ is divisorial and maps a surface S onto C. Since the morphism $\varphi|_S : S \to C$ is projective, the surface S is projective. Thus some multisection $C' \subset S$ has the required property.

4. MMP for pairs

In this section we prove several results on the MMP for pairs for Kähler threefolds. While we are not able to establish the log-MMP in full generality, the results are sufficient for the application to the abundance problem in Section 6. We start by proving the cone theorem and (parts of) the contraction theorem in the Subsections 4.A and 4.B, while Subsection 4.C merely collects the relevant results on existence of flips and termination. In order to simplify the statements we make the following

ASSUMPTION 4.1. – Let X be a normal Q-factorial compact Kähler threefold with rational singularities. Let $\Delta = \sum_i a_i \Delta_i$ be an effective Q-divisor on X such that the pair (X, Δ) is lc. We also assume that X is not uniruled, so by [38, Cor. 1.4] the Kodaira dimension $\kappa(\hat{X})$ of a desingularization \hat{X} is non-negative, hence $\kappa(X) \ge 0$. Thus we have

(7)
$$K_X \sim_{\mathbb{Q}} \sum \lambda_j S_j,$$

where the S_j are integral surfaces in X, the coefficients $\lambda_j \in \mathbb{Q}^+$.

4° SÉRIE - TOME 49 - 2016 - Nº 4

Note that in the situation of Theorem 1.3 and Theorem 1.4 the conditions above are satisfied: dlt pairs have rational singularities [51, 5.22].

4.A. Cone theorem

In this section we will prove a weak form of the cone theorem for non-uniruled lc pairs. The stronger cone Theorem 1.3 will then be a consequence of the contraction results in Subsection 4.B which are based on the weak cone theorem:

THEOREM 4.1. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold with rational singularities. Let Δ be a boundary divisor on X such that the pair (X, Δ) is lc. If X is not uniruled, there exists a positive integer $d \in \mathbb{N}$ and an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X such that

$$0 < -(K_X + \Delta) \cdot \Gamma_i \le d$$

and

$$\overline{NA}(X) = \overline{NA}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

We will prove this statement on page 985, but this will take some technical preparation.

LEMMA 4.1. – Under the Assumption 4.1, let $S \subset X$ be a prime divisor such that $(K_X + \Delta)|_S$ is not pseudoeffective. Then S is Moishezon and any desingularization \hat{S} is a uniruled projective surface.

Proof. – Let $\pi : \hat{S} \to S$ be the composition of the normalization followed by the minimal resolution of the normalized surface. Our goal is to show that there exists an effective \mathbb{Q} -divisor E such that $K_{\hat{S}} + E$ is not pseudoeffective, in particular we have $\kappa(\hat{S}) = -\infty$. It then follows from the Castelnuovo-Kodaira classification that \hat{S} is covered by rational curves, in particular it is a projective surface [6]. Thus S is Moishezon.

First case. Suppose that $K_X|_S$ is not pseudoeffective. – Since K_X is Q-effective this implies that S is one of the surfaces appearing in the decomposition (7). Up to renumbering we may suppose that $S = S_1$. Observe

$$S = S_1 = \frac{1}{\lambda_1} K_X - \sum_{j=2}^r \frac{\lambda_j}{\lambda_1} S_j.$$

Applying (2) with B = 0, there exists an effective Q-divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^* (K_X + S)|_S = \frac{(\lambda_1 + 1)}{\lambda_1} \pi^* K_X|_S - \sum_{j=2}^r \frac{\lambda_j}{\lambda_1} \pi^* S_j|_S.$$

By assumption $K_X|_S$ is not pseudoeffective, and $-\sum_{j=2}^r \frac{\lambda_j}{\lambda_1} \pi^* S_j|_S$ is anti-effective. Thus $K_{\hat{S}} + E$ is not pseudoeffective.

Second case. Suppose that $K_{X|_S}$ is pseudoeffective. – Then $S \subset \Delta$, hence, up to renumbering, we may suppose that $S = \Delta_1$. We claim that

$$(K_X + \Delta_1 + \sum_{i \ge 2} a_i \Delta_i)|_S$$

is not pseudoeffective. Otherwise, by $0 \le a_1 \le 1$ and the pseudo-effectivity of $K_X|_S$, the divisor

$$(K_X + \Delta)|_S = a_1(K_X + \Delta_1 + \sum_{i \ge 2} a_i \Delta_i)|_S + (1 - a_1)(K_X + \sum_{i \ge 2} a_i \Delta_i)|_S$$

is a convex combination of pseudoeffective classes, hence itself pseudoeffective, a contradiction. Applying (2) with $S = \Delta_1$ and $B = \sum_{i\geq 2} a_i \Delta_i$, there exists an effective \mathbb{Q} -divisor E such that

$$K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^* (K_X + \Delta + \sum_{i>2} a_i \Delta_i)|_S$$

Thus $K_{\hat{S}} + E$ is not pseudoeffective.

COROLLARY 4.1. – Under the Assumption 4.1, the divisor $K_X + \Delta$ is nef if and only if

$$(K_X + \Delta) \cdot C \ge 0$$

for every curve $C \subset X$.

Proof. – One implication is trivial. Suppose now that $K_X + \Delta$ is nef on all curves C. We will argue by contradiction and suppose that $K_X + \Delta$ is not nef. Since $K_X + \Delta$ is pseudo-effective and the restriction to every curve is nef, there exists by [12, Prop. 3.4] an irreducible surface $S \subset X$ such that $(K_X + \Delta)|_S$ is not pseudo-effective. By Lemma 4.1 a desingularization $\pi : \hat{S} \to S$ of the surface S is projective, so (by [13]) there exists a covering family of curves $C_t \subset S$ such that for the strict transforms

$$\pi^*(K_X + \Delta)|_S \cdot \hat{C}_t < 0.$$

Hence we obtain $(K_X + \Delta) \cdot C_t < 0$, a contradiction.

If X is a projective manifold, the cone theorem is a consequence of Mori's bend-and-break technique. We will now show that an analogue of this technique is available for threefolds that are lc pairs.

DEFINITION 4.1. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold, and let Δ be a boundary divisor. We say that $K_X + \Delta$ has the bend-and-break property if there exists a positive number $d = d_{K_X+\Delta} \in \mathbb{Q}^+$ such that the following holds: given any curve $C \subset X$ such that

$$-(K_X + \Delta) \cdot C > d_g$$

there exist non-zero effective 1-cycles C_1 and C_2 such that

$$[C] = [C_1] + [C_2].$$

PROPOSITION 4.1. – Under the Assumption 4.1, the divisor $K_X + \Delta$ has the bend-and-break property.

The proof of this result needs some preparation:

4° SÉRIE - TOME 49 - 2016 - Nº 4

LEMMA 4.2. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold with rational singularities, and let Δ be a boundary divisor on X. Let $\mu : Y \to X$ be a bimeromorphic morphism from a normal \mathbb{Q} -factorial threefold Y such that there exists a boundary divisor Δ_Y satisfying

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* (K_X + \Delta)$$

Suppose that $K_Y + \Delta_Y$ has the bend-and-break property for some integer $d_{K_Y+\Delta_Y}$. Then $K_X + \Delta$ has the bend-and-break property.

Proof. – Since dim X = 3, there are at most finitely many curves in the image of the μ -exceptional locus E. Hence

 $d := \max\{d_{K_Y + \Delta_Y}, -(K_X + \Delta) \cdot Z \mid Z \text{ an irreducible curve s.t. } Z \subset \mu(E)\}$

is a positive rational number. Let now $C \subset X$ be a curve such that

$$-(K_X + \Delta) \cdot C > d,$$

in particular $C \not\subset \mu(E)$. Thus the strict transform $\hat{C} \subset Y$ is well-defined and

$$-(K_Y + \Delta_Y) \cdot \hat{C} = -(K_X + \Delta) \cdot C > d \ge d_{K_Y + \Delta_Y}.$$

Consequently, there exist effective non-zero 1-cycles \hat{C}_1 and \hat{C}_2 on Y such that

$$[\hat{C}] = [\hat{C}_1] + [\hat{C}_2]$$

We claim that we can find a decomposition such that $\mu_*[\hat{C}_1]$ and $\mu_*[\hat{C}_2]$ are both non-zero. Since

$$[C] = \mu_*[\hat{C}] = \mu_*[\hat{C}_1] + \mu_*[\hat{C}_2]$$

this will finish the proof.

To prove the claim, fix a Kähler form ω_Y on Y such that $\omega_Y \cdot B \ge 1$ for every curve $B \subset Y$. We will prove the claim by induction on the degree $l := \omega_Y \cdot \hat{C}$. The start of the induction for l = 1 is trivial, since the class of a curve \hat{C} with $\omega_Y \cdot \hat{C} = 1$ does not decompose. For the induction step suppose that the claim holds for every curve $B \subset Y$ with $\omega_Y \cdot B < l$ and $-(K_Y + \Delta_Y) \cdot B > d$. Suppose that $l \le \omega_Y \cdot \hat{C} < l + 1$ and consider the decomposition

$$[\hat{C}] = [\hat{C}_1] + [\hat{C}_2].$$

If both $\mu_*[\hat{C}_1]$ and $\mu_*[\hat{C}_2]$ are non-zero, there is nothing to prove, so suppose that (up to renumbering) $\mu_*[\hat{C}_2] = 0$. Since \hat{C}_2 is effective, all the irreducible components of \hat{C}_2 are contracted by μ . Now $K_Y + \Delta_Y$ is μ -numerically trivial, so we get

$$-(K_Y + \Delta_Y) \cdot \hat{C}_1 = -(K_Y + \Delta_Y) \cdot \hat{C} > d.$$

Moreover since $\omega_Y \cdot \hat{C}_2 \ge 1$, we have $\omega_Y \cdot \hat{C}_1 < l$. Thus the induction hypothesis applies to C_1 , and since $[C] = \mu_*[\hat{C}] = \mu_*[\hat{C}_1]$ this proves the claim.

LEMMA 4.3. – In the situation of Assumption 4.1, suppose that K_X has the bend-and-break property for some integer d_0 . Then $K_X + \Delta$ has the bend-and-break property.

Proof. – Since X is a threefold, the set $(\text{Supp }\Delta)_{\text{sing}} \cup X_{\text{sing}}$ is a finite union of curves and points. Hence

 $d:=\max\{3,d_0,-(K_X+\Delta)\cdot Z\mid Z \text{ a curve s.t. } Z\subset (\operatorname{Supp}\Delta)_{\operatorname{sing}}\cup X_{\operatorname{sing}}\}$

is a positive rational number. Let now $C \subset X$ be a curve such that $-(K_X + \Delta) \cdot C > d$. If $\Delta \cdot C \ge 0$, then we have $-K_X \cdot C > d \ge d_0$, so we can apply the bend-and-break property for K_X . Therefore we may suppose

$$0 > \Delta \cdot C = \sum a_i \Delta_i \cdot C,$$

so, up to renumbering, we may suppose that $\Delta_1 \cdot C < 0$. Since $0 \le a_i \le 1$ (Δ is a boundary), this implies

$$-(K_X + \Delta_1 + \sum_{i \ge 2} a_i \Delta_i) \cdot C \ge -(K_X + \Delta) \cdot C > d.$$

Set $S := \Delta_1$ and $B := \sum_{i \ge 2} a_i \Delta_i$, and denote by $\pi : \hat{S} \to S$ the composition of the normalization and the minimal resolution. Note that by definition of d, the curve C is not contained in the set $S_{\text{sing}} \cup (\sum_{i \ge 2} \Delta_i \cap S) \cup (X_{\text{sing}} \cap S)$. Therefore the strict transform $\hat{C} \subset \hat{S}$ is well-defined and (3) yields

$$-K_{\hat{S}} \cdot \hat{C} \ge -(K_X + \Delta_1 + \sum_{i \ge 2} a_i \Delta_i) \cdot C > d.$$

Since $d \ge 3$, an application of [38, Lemma 5.5.b)] yields an effective 1-cycle $\sum_{k=1}^{m} \hat{C}_k$ in \hat{S} with $m \ge 2$ such that

$$\left[\sum_{k=1}^{m} \hat{C}_k\right] = \left[\hat{C}\right]$$

such that $K_{\hat{S}} \cdot \hat{C}_1 < 0$, and $K_{\hat{S}} \cdot \hat{C}_2 < 0$. Since $K_{\hat{S}}$ is π_1 -nef, we conclude $\pi_*[\hat{C}_1] \neq 0$ and $\pi_*[\hat{C}_2] \neq 0$. Thus

$$[C] = \pi_*[\hat{C}] = \sum_{k=1}^m \pi_*[\hat{C}_k]$$

and the first two terms of this sum are non-zero.

Proof of Proposition 4.1. – We will first prove the statement under some additional assumptions, then reduce the general case to this situation.

Step 1. Suppose that (X, Δ) is a lc pair and X has terminal singularities. – By [38, Cor. 5.7] we know that K_X has the bend-and-break property. Thus by Lemma 4.3 the divisor $K_X + \Delta$ has the bend-and-break property.

Step 2. Suppose that X has klt singularities, i.e., (X, 0) is klt. – By Theorem 2.1 there exists a bimeromorphic morphism $\mu : Y \to X$ from a normal complex Kähler space Y with terminal singularities and a boundary divisor Δ_Y such that (Y, Δ_Y) is lc and

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* K_X.$$

By Step 1 and Lemma 4.2 the divisor K_X has the bend-and-break property.

Step 3. Suppose that (X, Δ) is dlt. – Then (X, 0) is klt [51, 2.39, 2.41]. By Step 2 and Lemma 4.3 this implies that $K_X + \Delta$ has the bend-and-break property.

4° SÉRIE - TOME 49 - 2016 - Nº 4

Step 4. General case. – By Theorem 2.2 there exists a bimeromorphic morphism $\mu : Y \to X$ and a boundary divisor Δ_Y on Y such that (Y, Δ_Y) is dlt and

$$K_Y + \Delta_Y \sim_{\mathbb{Q}} \mu^* (K_X + \Delta)$$

By Step 3 and Lemma 4.2 this implies that $K_X + \Delta$ has the bend-and-break property. \Box

We can now prove the weak cone theorem for the classical Mori cone.

PROPOSITION 4.2. – Under the Assumption 4.1 there exists a number $d \in \mathbb{Q}^+$ and an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X such that

$$0 < -(K_X + \Delta) \cdot \Gamma_i \le d$$

and

$$\overline{NE}(X) = \overline{NE}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

Proof. – By Proposition 4.1 there exists a positive number $d \in \mathbb{Q}^+$ realizing the bendand-break property (cf. Definition 4.1) for $K_X + \Delta$. Since there are only countably many curve classes $[C] \subset \overline{NE}(X)$, we may choose a representative Γ_i for each class such that $0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$. We set

$$V := \overline{\operatorname{NE}}(X)_{(K_X + \Delta) \ge 0} + \sum_{0 < -(K_X + \Delta) \cdot \Gamma_i \le d} \mathbb{R}^+[\Gamma_i].$$

Fix a Kähler form η_X on X such that $\eta_X \cdot C \ge 1$ for every curve $C \subset X$. We need to prove that $\overline{NE}(X) = V$. By [38, Lemma 6.1]⁽³⁾ it is sufficient to show that

$$\overline{\operatorname{NE}}(X) = \overline{V},$$

i.e., the class [C] of every irreducible curve $C \subset X$ is contained in V. We will prove the statement by induction on the degree $l := \eta_X \cdot C$. The start of the induction for l = 0 being trivial, we suppose that we have shown the statement for all curves of degree at most l - 1 and let C be a curve such that

$$l-1 < \eta_X \cdot C \le l.$$

If $-(K_X + \Delta) \cdot C \leq d$ we are done. Otherwise, there exists by the bend-and-break property a decomposition

$$[C] = [C_1] + [C_2]$$

with C_1 and C_2 non-zero effective 1-cycles on X. Since $\eta_X \cdot C_i \ge 1$ for i = 1, 2 we have $\eta_X \cdot C_i \le l - 1$ for i = 1, 2. By induction both classes are in V, so [C] is in V. \Box

Proof of Theorem 4.1. – We follow the strategy of the proof of [38, Prop. 6.4]. By Proposition 4.2 there exists a number $d \in \mathbb{Q}^+$ and an at most countable family $(\Gamma_i)_{i \in I}$ of curves on X such that $0 < -(K_X + \Delta) \cdot \Gamma_i \leq d$ and

$$\overline{\operatorname{NE}}(X) = \overline{\operatorname{NE}}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

⁽³⁾ The statement in [38, Lemma 6.1] is for the canonical class K_X , but the proof does not use this hypothesis.

Set

$$V := \overline{\mathrm{NA}}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i].$$

By [38, Lemma 6.1] it is sufficient to show that $\overline{NA}(X) \subset \overline{V}$. Let $\pi : \hat{X} \to X$ be a desingularization, then by [38, Prop. 3.14] we have $\overline{NA}(X) = \pi_*(\overline{NA}(\hat{X}))$. Thus it is sufficient to prove that for a set of generators $\hat{\alpha}_i$ of $\overline{NA}(\hat{X})$, we have $\pi_*(\alpha_i) \in \overline{V}$. By [23, Cor. 0.3] the cone $\overline{NA}(\hat{X})$ is the closure of the convex cone generated by cohomology classes of the form $[\hat{\omega}]^2$, $[\hat{\omega}] \cdot [\hat{S}]$ and $[\hat{C}]$ where $\hat{\omega}$ is a Kähler form, \hat{S} a surface and \hat{C} a curve on \hat{X} . Let now $\hat{\alpha}$ be such a generator, then our goal is to show that the push-forward $\alpha := \pi_*(\hat{\alpha})$ of any of these three types is contained in \overline{V} .

First case. $\hat{\alpha} = [\hat{\omega}]^2$ with $\hat{\omega}$ a Kähler form. – Since $\pi^*(K_X + \Delta)$ is pseudoeffective, we have $\pi^*(K_X + \Delta) \cdot [\hat{\omega}]^2 \ge 0$, hence

$$(K_X + \Delta) \cdot \alpha = (K_X + \Delta) \cdot \pi_*(\hat{\alpha}) = \pi^*(K_X + \Delta) \cdot \hat{\alpha} \ge 0,$$

and thus $\alpha \in \overline{\operatorname{NA}}(X)_{(K_X + \Delta) \ge 0}$.

Second case. $\hat{\alpha} = [\hat{C}]$ with \hat{C} a curve. – Then set $C := \pi_*(\hat{C})$, so that $\alpha = [C]$. Since we have an inclusion

(8)
$$\overline{\operatorname{NE}}(X) = \overline{\operatorname{NE}}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i] \hookrightarrow \overline{\operatorname{NA}}(X)_{(K_X + \Delta) \ge 0} + \sum_{i \in I} \mathbb{R}^+[\Gamma_i],$$

and by hypothesis any curve class [C] is in the left hand side, we see that $[C] \in V$.

Third case. $\hat{\alpha} = [\hat{\omega}] \cdot [\hat{S}]$ with \hat{S} an irreducible surface and $\hat{\omega}$ a Kähler form. – If

 $\pi^*(K_X + \Delta) \cdot [\hat{\omega}] \cdot [\hat{S}] \ge 0,$

the class α is in $\overline{NA}(X)_{(K_X + \Delta) \ge 0}$. Suppose now that

$$\pi^*(K_X + \Delta) \cdot [\hat{\omega}] \cdot [\hat{S}] < 0.$$

Using the projection formula we see that $\pi(\hat{S})$ is not a point.

Case a). Suppose that $\pi(\hat{S})$ is a surface S. – Since $\pi^*(K_X + \Delta) \cdot [\hat{\omega}] \cdot [\hat{S}] < 0$, the restriction $\pi^*(K_X + \Delta)|_{\hat{S}}$ is not pseudoeffective. Thus the restriction $(K_X + \Delta)|_S$ is not pseudoeffective. Hence S is covered by rational curves by Lemma 4.1. Since $\hat{S} \to S$ is bimeromorphic, the same property holds for \hat{S} .

Let $\pi: \overline{S} \to \hat{S}$ be the composition of the normalization and the minimal resolution, then \overline{S} is a uniruled projective surface, in particular, $H^2(\overline{S}, \mathcal{O}_{\overline{S}}) = 0$. Hence the Chern class map

$$\operatorname{Pic} \bar{S} \to H^2(\bar{S}, \mathbb{Z})$$

is surjective, so $\pi^*(\hat{\omega}|_{\hat{S}})$, which is a real closed form of type (1, 1), is represented by a \mathbb{R} -divisor which is nef and big. As in the proof of [38, Prop. 6.4], this implies that

$$\hat{\alpha} = [\hat{\omega}] \cdot [\hat{S}] \in \overline{\mathrm{NE}}(\hat{X})$$

Thus we have

$$\alpha = \pi_*([\hat{\omega}] \cdot [\hat{S}]) \in \overline{\operatorname{NE}}(X),$$

so α is in the image of the inclusion (8).

4° SÉRIE - TOME 49 - 2016 - Nº 4

Case b). Suppose that $\pi(\hat{S})$ is a curve C. – We claim that there exists a number $\lambda \in \mathbb{R}$ such that

$$\alpha = \pi_*(\hat{\alpha}) = \lambda[C]$$

in $N_1(X)$, in particular α is in the image of the inclusion (8) which finishes the proof. By duality it is sufficient to prove that there exists a $\lambda \in \mathbb{R}$ such that for every class $H \in N^1(X)$ we have

$$H \cdot \pi_*(\hat{\alpha}) = \lambda (H \cdot C).$$

By the projection formula one has

$$H \cdot \pi_*(\hat{\alpha}) = \pi^* H \cdot \hat{\alpha} = \pi^* H \cdot [\hat{\omega}] \cdot [\hat{S}] = [(\pi^* H)_{|_{S}}] \cdot [\hat{\omega}_{|_{\hat{S}}}]$$

By definition of C we have a surjective map $\hat{S} \to C$, so $[(\pi^*H)|_S]$ is numerically equivalent to $(H \cdot C)[F]$ where F is a general fibre of $\hat{S} \to C$. Thus we see that

$$[(\pi^*H)|_{S}] \cdot [\hat{\omega}|_{\hat{S}}] = \lambda(H \cdot C)$$

where $\lambda := [\hat{\omega}|_{\hat{S}}] \cdot F$ does not depend on *H*.

4.B. The contraction theorem

Suppose that the Assumption 4.1 holds. For the whole subsection we fix $R := \mathbb{R}^+[\Gamma_{i_0}]$ a $(K_X + \Delta)$ -negative extremal ray in $\overline{NA}(X)$. The following proposition is a well-known consequence of the weak cone Theorem 4.1, cf. [38, Prop. 7.3] for details:

PROPOSITION 4.3. – There exists a nef class $\alpha \in N^1(X)$ such that

$$R = \{ z \in \overline{NA}(X) \mid \alpha \cdot z = 0 \},\$$

and such that, using the notation of Theorem 4.1, the class α is strictly positive on

$$\left(\overline{NA}(X)_{(K_X+\Delta)\geq 0}+\sum_{i\in I, i\neq i_0}\mathbb{R}^+[\Gamma_i]\right)\setminus\{0\}.$$

We call α a nef supporting class for the extremal ray R.

Note first that by hypothesis, the cohomology class $\alpha - (K_X + \Delta)$ is positive on the extremal ray R, moreover we know that α is positive on

$$\left(\overline{\mathrm{NA}}(X)_{(K_X+\Delta)\geq 0} + \sum_{i\in I, i\neq i_0} \mathbb{R}^+[\Gamma_i]\right) \setminus \{0\}.$$

Thus, up to replacing α by some positive multiple, we can suppose that $\alpha - (K_X + \Delta)$ is positive on $\overline{NA}(X) \setminus \{0\}$. Since X is a Kähler space, this implies by [38, Cor. 3.16] that

(9)
$$\omega := \alpha - (K_X + \Delta)$$

is a Kähler class. Let now $\pi : \hat{X} \to X$ be a desingularization. Since $K_X + \Delta$ is \mathbb{Q} -effective, this also holds for $\pi^*(K_X + \Delta)$. Thus the nef class $\pi^*\alpha$ is the sum of a \mathbb{Q} -effective class $\pi^*(K_X + \Delta)$ and the nef and big (i.e., contains a Kähler current, see [23, Thm. 0.5]) class $\pi^*\alpha$. Thus $\pi^*\alpha$ is nef and big and we have

(10)
$$\alpha^3 = (\pi^* \alpha)^3 > 0.$$

ANNALES SCIENTIFIQUES DE L'ÉCOLE NORMALE SUPÉRIEURE

We divide extremal rays into two classes, according to the deformation behavior of the curves they contain:

DEFINITION 4.2. – We say that the $(K_X + \Delta)$ -negative extremal ray R is small if every curve $C \subset X$ with $[C] \in R$ is very rigid in the sense of [38, Defn.4.3]. Otherwise we say that the extremal ray R is divisorial.

4.B.1. Small rays

THEOREM 4.2. – Under the Assumption 4.1, suppose that the extremal ray $R = \mathbb{R}^+[\Gamma_{i_0}]$ is small. Then the contraction of the ray R exists.

The proof requires a significant refinement of the argument in [38] since the description of the non-Kähler locus of α is much more delicate for non-isolated singularities. The following lemma is a key ingredient:

PROPOSITION 4.4. – Under the Assumption 4.1, suppose that the extremal ray $R = \mathbb{R}^+[\Gamma_i]$ is small. Let $S \subset X$ be an irreducible surface. Then we have $\alpha^2 \cdot S > 0$.

At this point we cannot yet exclude the possibility that there are infinitely many distinct curves $C \subset X$ such that $[C] \in R^{(4)}$. However by definition of a small ray no such curve (or its multiples) deforms. Since the irreducible components of the cycle space are countable we see that there are at most countably many curves $C \subset X$ such that $[C] \in R$.

Proof of Proposition 4.4. – If $\alpha_{|_{S}} = 0$ then (9) implies that

 $-(K_X + \Delta)|_S = \omega|_S.$

Thus the divisor $-(K_X + \Delta)|_S$ is ample, in particular S is projective and S is covered by curves. Since $\alpha|_S = 0$, the classes of all these curves are in R, a contradiction.

Arguing by contradiction we suppose now that $\alpha|_S \neq 0$ but $\alpha^2 \cdot S = 0$. Then we have

$$0 = \alpha^2 \cdot S = (K_X + \Delta) \cdot \alpha \cdot S + \omega \cdot \alpha \cdot S$$

and

$$\omega \cdot \alpha \cdot S = \omega_{|_{\mathcal{S}}} \cdot \alpha_{|_{\mathcal{S}}} > 0$$

since $\alpha_{|_{S}}$ is a non-zero nef class. Thus we obtain

(11)
$$(K_X + \Delta) \cdot \alpha \cdot S < 0.$$

In particular $(K_X + \Delta)|_S$ is not pseudoeffective, the class $\alpha|_S$ being nef.

Let $\pi : \hat{S} \to S$ be the composition of normalization and minimal resolution (cf. Subsection 2.B). We claim that there exists an effective \mathbb{Q} -divisor E on \hat{S} such that

(12)
$$(K_{\hat{S}} + E) \cdot \pi^*(\alpha|_{S}) < 0.$$

Assuming this for the time being, let us see how to derive a contradiction: note first that $K_{\hat{S}}$ is not pseudoeffective, so \hat{S} is uniruled and projective. In particular the nef class $\pi^*(\alpha|_{\alpha})$ is

⁽⁴⁾ If X has terminal singularities we can use some additional argument to obtain this property, cf. [38, Rem.7.2].

4° SÉRIE - TOME 49 - 2016 - Nº 4

represented by an \mathbb{R} -divisor. Fix an ample divisor A on \hat{S} . By [4, Thm. 1.3] we know that given $\varepsilon > 0$, there exists a decomposition

$$\pi^*(\alpha|_S) = C_{\varepsilon} + \sum \lambda_{i,\varepsilon} M_{i,\varepsilon}$$

where $\lambda_{i,\varepsilon} \ge 0$, $(K_{\hat{S}} + \varepsilon A) \cdot C_{\varepsilon} \ge 0$ and the $M_{i,\varepsilon}$ are movable curves. Since $M_{i,\varepsilon}$ belongs to an (uncountable) deformation family of curves we obtain

$$\pi^*(\alpha|_{\mathfrak{S}}) \cdot M_{i,\varepsilon} = \alpha \cdot \pi_*(M_{i,\varepsilon}) > 0.$$

Since $(\pi^*(\alpha|_S))^2 = 0$ this implies that $\pi^*(\alpha|_S) = C_{\varepsilon}$ for all $\varepsilon > 0$. Passing to the limit $\varepsilon \to 0$ we obtain $K_{\hat{S}} \cdot \pi^*(\alpha|_S) \ge 0$, a contradiction to (12).

Proof of the claim (12). - As in the proof of Lemma 4.1 we need a case distinction.

First case. Suppose that $K_X \cdot \alpha \cdot S < 0$. – Since K_X is Q-effective, S is one of the surfaces S_j in the decomposition (7) and $S \cdot \alpha \cdot S < 0$. In particular we have

(13)
$$(K_X + S) \cdot \alpha \cdot S < 0.$$

Applying (2) with B = 0 we obtain an effective \mathbb{Q} -divisor E on \hat{S} such that $K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^*(K_X + S)|_S$. Thus (12) follows from (13).

Second case. Suppose that $K_X \cdot \alpha \cdot S \ge 0$. – By (11) this implies that $\Delta \cdot \alpha \cdot S < 0$, so S is contained in the support of Δ ; up to renumbering we may suppose that $S = \Delta_1$. Since $\Delta_i \cdot \alpha \cdot S \ge 0$ for every $i \ge 2$, it follows $S \cdot \alpha \cdot S < 0$. Since $0 \le a_i \le 1$ we conclude

(14)
$$(K_X + S + \sum_{i \ge 2} a_i \Delta_i) \cdot \alpha \cdot S \le (K_X + \Delta) \cdot \alpha \cdot S < 0.$$

Applying (2) with $B = \sum_{i\geq 2} a_i \Delta_i$ we obtain an effective \mathbb{Q} -divisor E on \hat{S} such that $K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^* (K_X + S + \sum_{i\geq 2} a_i \Delta_i)|_S$. Thus (12) follows from (14).

Proof of Theorem 4.2. – Let $\pi : \hat{X} \to X$ be a desingularization. By (10) we have $(\pi^*\alpha)^3 > 0$, so we can apply the theorem of Collins and Tosatti [19, Thm. 1.1] to $\pi^*\alpha$: the non-Kähler locus $E_{nK}(\pi^*\alpha)$ is equal to the null-locus, i.e., we have

$$E_{nK}(\pi^*\alpha) = \bigcup_{\substack{(\pi^*\alpha) \mid d \text{im } Z = 0 \\ \alpha}} Z$$

where the union runs over all the subvarieties of \hat{X} . If $Z \subset X$ is a surface such that $(\pi^* \alpha)|_Z^2 = 0$ then it follows from the projection formula and Proposition 4.4 that dim $\pi(Z) \leq 1$. Thus we see that $E_{nK}(\pi^* \alpha)$ is a finite union of π -exceptional surfaces and curves. Since

$$E_{nK}(\alpha) \subset \pi(E_{nK}(\pi^*\alpha)),$$

and X_{sing} is a union of curves and points, we see that $E_{nK}(\alpha)$ is a union of curves and points. Clearly $E_{nK}(\alpha)$ contains all the curves $C \subset X$ such that $[C] \in R$, in particular

$$C := \bigcup_{C_l \subset X, [C_l] \in R} C_l$$

is a finite union of curves. Our goal is to show that the connected components of C can be contracted onto points. Since it is a-priori not clear that $C = E_{nK}(\alpha)$ we have to improve the construction from [38, Thm. 7.12]

By [12, Prop. 2.3]⁽⁵⁾ there exists a modification $\mu : \tilde{X} \to X$ and a Kähler class $\tilde{\alpha}$ on \tilde{X} such that $\mu_* \tilde{\alpha} = \alpha$. Since $\tilde{\alpha} - \mu^* \alpha$ is μ -nef, the negativity lemma [51, Lemma 3.39] [10, 3.6.2] implies that we have

(15)
$$\mu^* \alpha = \tilde{\alpha} + E$$

with *E* an effective μ -exceptional \mathbb{R} -divisor. Let now $C_l \subset X$ be an irreducible curve such that $[C_l] \in R$, and set D_l for the support of $\mu^{-1}(C_l)$. Then we have $\alpha|_{C_l} \equiv 0$, so by (15) we have

$$-E_{\mid D_{I}} = \tilde{\alpha}_{\mid D_{I}}$$

Thus $-E_{|_{D_l}}$ is ample. If $B \subset X$ is an arbitrary curve contained in the image of the exceptional locus, and D_B the support of $\mu^{-1}(B)$, we still have

$$-E_{\big|_{D_B}} = \tilde{\alpha}_{\big|_{D_B}} - (\mu^* \alpha)_{\big|_{D_B}}.$$

Thus we see that -E is μ -ample, in particular its support is equal to the μ -exceptional locus. Since ampleness is an open property we can find an effective \mathbb{Q} -divisor $E' \subset \tilde{X}$ that is μ -ample and such that $-E'|_{D_l}$ is ample for all l. Up to taking a positive multiple we can suppose that E' has integer coefficients. We set

$$\mathscr{K}_m := \mu_* \mathscr{O}_{\tilde{X}}(-mE'),$$

and we claim that there exists an $m \in \mathbb{N}$ such that the restriction $\mathcal{K}_m|_{C_l}$ is ample for all j. Note that this implies that the quotient $(\mathcal{K}_m/\mathcal{K}_m^2)|_{C_l}$ is ample for all j.

Step 2. Proof of the claim. – By relative Serre vanishing there exists an $m_0 \in \mathbb{N}$ such that

$$R^{i}\mu_{*}\mathcal{O}_{\tilde{X}}(-mE') = 0 \qquad \forall i > 0, m \ge m_{0}$$

and

(16)
$$R^{i}\mu_{*}\mathcal{O}_{\tilde{X}}(-D_{l}-mE')=0 \qquad \forall i>0, m\geq m_{0}$$

and all *l*. Moreover we know by [1, Thm. 3.1.] that there exists an $m_1 \in \mathbb{N}$ such that the direct image sheaf

$$(\mu|_{D_l})_*(\mathcal{O}_{\tilde{X}}(-mE')\otimes\mathcal{O}_{D_l})$$

is ample for all $m \ge m_1$ and l.

Fix now an $m \ge \max\{m_0, m_1\}$ and consider now the exact sequence

$$0 \to \mathcal{O}_{\tilde{X}}(-D_l - mE') \to \mathcal{O}_{\tilde{X}}(-mE') \to \mathcal{O}_{\tilde{X}}(-mE') \otimes \mathcal{O}_{D_l} \to 0.$$

Pushing the sequence down to X and using (16) we obtain an exact sequence

$$0 \to \mu_* \mathcal{O}_{\tilde{X}}(-D_l - mE') \to \mu_* \mathcal{O}_{\tilde{X}}(-mE') \to (\mu_{|_{D_l}})_* (\mathcal{O}_{\tilde{X}}(-mE') \otimes \mathcal{O}_{D_l}) \to 0.$$

⁽⁵⁾ Proposition 2.3. in [12] is for manifolds, but we can adapt the proof as follows: the non-Kähler locus of $\pi^* \alpha$ is the union of the π -exceptional locus and the strict transforms of the curves in the non-Kähler locus of α . By [11, Thm. 3.1.24] there exists a modification $\mu' : \tilde{X} \to \hat{X}$ and a Kähler class $\tilde{\alpha}$ on \tilde{X} such that $(\mu')^* \pi^* \alpha = \tilde{\alpha} + E$ where *E* is an effective \mathbb{R} -divisor. Then $\mu := \pi \circ \mu'$ has the stated properties.

4° SÉRIE – TOME 49 – 2016 – Nº 4

Since the inclusion $\mathcal{O}_{\tilde{X}}(-D_l - mE') \to \mathcal{O}_{\tilde{X}}(-mE')$ vanishes on D_l , its direct image

$$\mu_* \mathcal{O}_{\tilde{X}}(-D_l - mE') \to \mu_* \mathcal{O}_{\tilde{X}}(-mE')$$

vanishes on $C_l = \mu(D_l)$. In other words we have $\mathcal{O}_{\tilde{X}}(-D_l - mE') \subset \mathcal{J} \cdot \mathcal{O}_{\tilde{X}}(-mE')$ where \mathcal{J} is the full ideal sheaf of C_l . Thus we have an epimorphism

$$\mu_* \mathcal{O}_{\tilde{X}}(-mE') \otimes \mathcal{O}_{C_l} \to (\mu_{|_{D_l}})_* (\mathcal{O}_{\tilde{X}}(-mE') \otimes \mathcal{O}_{D_l}),$$

which is generically an isomorphism. In particular, $\mu_* \mathcal{O}_{\tilde{X}}(-mE') \otimes \mathcal{O}_{C_l}$ is ample. This proves the claim.

Step 3. Conclusion. – The ideal sheaf \mathcal{K}_m defines a 1-dimensional subspace $A \subset X$ such that $C \subset A$. Let $\mathscr{I} \subset \mathscr{O}_X$ be the largest ideal sheaf on X that coincides with \mathcal{K}_m in the generic point of every irreducible curve $C_l \subset C^{(6)}$ such that $[C_l] \in R$. For every curve C_l , the natural map

$$(\mathcal{K}_m/\mathcal{K}_m^2)|_{C_l} \to (\mathcal{I}/\mathcal{I}^2)|_{C_l}$$

is generically an isomorphism. Since $(\mathcal{K}_m/\mathcal{K}_m^2)|_{C_l}$ is ample and C_l is a curve, this implies that $(\mathscr{I}/\mathscr{I}^2)|_{C_l}$ is ample. Thus $\mathscr{I}/\mathscr{I}^2$ is ample on its support C. By [3, Cor. 3] [33] there exists a holomorphic map $\nu : X \to X'$ contracting each connected component of C onto a point.

4.B.2. Divisorial rays

NOTATION 4.1. – Under the Assumption 4.1, suppose that the extremal ray $R = \mathbb{R}^+[\Gamma_{i_0}]$ is divisorial. Since the divisor $K_X + \Delta$ is \mathbb{Q} -effective and $(K_X + \Delta) \cdot R < 0$ there exists an irreducible surface $S \subset X$ such that $S \cdot R < 0$. In particular any curve $C \subset X$ with $[C] \in R$ is contained in S and S is covered by these curves.

Let $\nu : \tilde{S} \to S \subset X$ be the normalization; then $\nu^*(\alpha|_S)$ is a nef class on \tilde{S} and we may consider the nef reduction

$$\tilde{f}: \tilde{S} \to \tilde{T}$$

with respect to $\nu^*(\alpha|_S)$ (cf. [7, Thm. 2.6] and [37, Thm. 3.19] for details on the Kähler case). Since S is covered by curves that are α -trivial, the surface \tilde{S} is covered by curves that are $\nu^*(\alpha|_S)$ -trivial. By definition of the nef reduction this implies

$$n(\alpha) := n(\nu^*(\alpha|_{c})) := \dim T \in \{0, 1\}.$$

LEMMA 4.4. – Under the Assumption 4.1, suppose that the extremal ray R is divisorial and $n(\alpha) = 0$. Then the surface S can be blown down to a point p: there exists a bimeromorphic morphism $\varphi : X \to Y$ to a normal compact threefold Y with dim $\varphi(S) = 0$ such that $\varphi|_{X \setminus S}$ is an isomorphism onto $Y \setminus \{p\}$.

Proof. – The proof is identical to the proof of [38, Cor. 7.7] which only uses that $S \cdot R < 0$.

⁽⁶⁾ The sheaf \mathscr{I} defines the scheme-theoretic image [36, II,Ex.3.11(d)] of the natural map $\bigcup_l C_{l,\text{gen}} \to X$ where we endowed the generic points $C_{l,\text{gen}} \subset A$ with the open subscheme structure.

The case $n(\alpha) = 1$ is much more subtle. If $S \subset \text{Supp }\Delta$, we can suppose up to renumbering that $S = \Delta_1$ and set $B = \sum_{i \geq 2} a_i \Delta_i$. If $S \not\subset \text{Supp }\Delta$ we simply set $B = \Delta$. In both cases we have by (1) that

(17)
$$K_{\tilde{S}} + B_{\tilde{S}} \sim_{\mathbb{Q}} \nu^* (K_X + S + B)|_S,$$

where $B_{\tilde{S}}$ is a canonically defined effective \mathbb{Q} -divisor. Let \tilde{C} be a general fibre of the nef reduction $\tilde{f}: \tilde{S} \to \tilde{T}$. Since $[\nu(\tilde{C})] \in R$ we have

$$(K_X + \Delta) \cdot \nu(\tilde{C}) < 0$$
 and $S \cdot \nu(\tilde{C}) < 0$

Since $0 \le a_1 \le 1$ this implies

$$(K_X + S + B) \cdot \nu(\tilde{C}) \le (K_X + \Delta) \cdot \nu(\tilde{C}) < 0.$$

Thus we have $(K_{\tilde{S}} + B_{\tilde{S}}) \cdot \tilde{C} < 0$, in particular $\tilde{C}^2 = 0$ implies that $\tilde{C} \simeq \mathbb{P}^1$. The normal surface \tilde{S} being smooth in a neighborhood of the fibre \tilde{C} , we conclude $K_{\tilde{S}} \cdot \tilde{C} = -2$. Since $B_{\tilde{S}} \cdot \tilde{C} \ge 0$ we arrive at

LEMMA 4.5. – Under the Assumption 4.1, suppose that the extremal ray R is divisorial and $n(\alpha) = 1$. Then the extremal ray R is represented by a rational curve C such that

$$-(K_X + \Delta) \cdot C \le 2.$$

This estimate allows us to complete the proof of the cone theorem:

Proof of Theorem 1.3. – The only statement that is not part of Theorem 4.1 is that in every $(K_X + \Delta)$ -negative extremal ray R_i we can find a rational curve Γ_i such that $\Gamma_i \in R_i$ and

$$0 < -(K_X + \Delta) \cdot \Gamma_i \le 4.$$

If the extremal ray Γ_i is divisorial with $n(\alpha) = 1$, we conclude by Lemma 4.5 (even if (X, Δ) is only lc). If the extremal ray Γ_i is small or divisorial with $n(\alpha) = 0$, the contraction exists by Theorem 4.2 and Lemma 4.4. Since (X, Δ) is dlt, we may simply apply [41] [21, Thm. 7.46].

REMARK 4.1. – In order to prove Theorem 1.3 in the lc case, it remains to prove the existence of rational curves if the ray is divisorial contracting a divisor S to a point or if the ray is small, with the bound d = 4 (cf. [28] for the projective case). The existence of rational curves in the first case just follows from the arguments preceding Lemma 4.5, which show that S is uniruled. The existence of rational curves in the small case requires some vanishing theorem which is not yet established in the Kähler case: if W is the union of the lc centers, then $H^1(W, \Theta_W) = 0$.

In order to prove the existence of the contraction in the case $n(\alpha) = 1$ we would like to construct a fibration $S \to T$ whose normalization is the nef reduction $\tilde{S} \to \tilde{T}$. At the moment we can only realize this strategy under an additional condition on the singularities of S. We will use the notion of a semi-log-canonical (slc) surface; see Section 5.

4° SÉRIE – TOME 49 – 2016 – Nº 4

PROPOSITION 4.5. – Under the Assumption 4.1, suppose that the extremal ray R is divisorial and $n(\alpha) = 1$. Suppose moreover S has slc singularities, i.e., (S,0) is slc. Then there exists a morphism with connected fibres $S \to T$ onto a curve T that contracts a curve $C \subset S$ if and only if $\alpha|_{S} \cdot C = 0$. Moreover the contraction of the extremal ray R exists.

Proof. – Since (S,0) is slc, it has normal crossing singularities in codimension one. The normalization map $\nu : \tilde{S} \to S$ is finite, so the general curve $\nu(\tilde{C})$ is contained in the locus where S has normal crossing singularities. If $\nu(\tilde{C})$ is in S_{nons} the proof of [38, Lemma 7.8] applies without changes, so suppose that this is not the case. Let

$$Z = \bigcup Z_k \subset S_{\text{sing}}$$

be the union of curves Z_k such that $\nu^{-1}(Z_k)$ meets the general curve \tilde{C} . For each k we set $\tilde{Z}_k := \nu^{-1}(Z_k)$ and set $\tilde{Z} := \bigcup \tilde{Z}_k$. Since S has normal crossings in the generic point of Z_k , the natural map $\tilde{Z}_k \to Z_k$ has degree two. Note now that every irreducible component of \tilde{Z} is $\nu^* \alpha$ -positive: otherwise \tilde{S} would be connected by $\nu^* \alpha$ -trivial curves, thus $n(\alpha) = 0$ in contradiction to our assumption. In particular every curve in \tilde{Z} dominates \tilde{T} .

Since \hat{Z} maps into the normal crossings locus of S we can decompose the different (17)

$$B_{\tilde{S}} = \tilde{Z} + R$$

where R is an effective \mathbb{Q} -divisor with no common component with \tilde{Z} . The intersection points of \tilde{Z} and \tilde{C} are contained in the smooth locus of \tilde{S} , so $\tilde{Z} \cdot \tilde{C}$ is a positive integer. On the other hand we know that $K_{\tilde{S}} \cdot \tilde{C} = -2$ and

$$-2 < (K_{\tilde{S}} + \tilde{Z}) \cdot \tilde{C} \le (K_{\tilde{S}} + \tilde{Z} + R) \cdot \tilde{C} < 0.$$

Thus $\tilde{Z} \cdot \tilde{C} = 1$. Since all the irreducible components of \tilde{Z} surject onto \tilde{T} , this shows that \tilde{Z} is irreducible. In particular Z itself is irreducible. Moreover we conclude that

- a) the curves Z and C meet in a unique point q; and
- b) we have $\nu^{-1}(q) = \{p_1, p_2\}$ with $p_1 \in \tilde{C}$, but $p_2 \notin \tilde{C}$.

Since \tilde{C} is general and ν is finite, the point p_2 lies on another general fibre \bar{C} such that $\nu(\tilde{C})$ and $\nu(\bar{C})$ meet in q. Let now T be the unique irreducible component of the cycle space $\operatorname{Chow}(S)$ such that the general point corresponds to the cycle $\nu(\tilde{C}) + \nu(\bar{C})$, and let $\Gamma \to T'$ be the semi-normalization [47, p.156] of the universal family over T. By construction we have a natural bimeromorphic morphism $\Gamma \to S$ and, by what precedes, this morphism is an isomorphism in the neighborhood of the general fibre F of $\Gamma \to T'$. Thus F defines a Cartier divisor on S such that $F^2 = 0$ and $\kappa(S, F) = 1$. Thus some positive multiple of F defines a morphism with connected fibres $S \to T$ and one easily checks that this morphism realises the nef reduction with respect to $\alpha|_{C}$. Now conclude as in [38, Cor. 7.9].

4.C. Running the MMP

We first collect a number of results which we could not find in this form in the literature and give an indication how to adapt the proof.

THEOREM 4.3 ([62], in the algebraic case [49]). – Let X be a normal Q-factorial compact Kähler threefold. Let Δ be a boundary divisor such that the pair (X, Δ) is dlt. Let $\mathbb{R}^+[\Gamma_i]$ be a $(K_X + \Delta)$ -negative extremal ray R in $\overline{NA}(X)$. Suppose that the contraction of $\varphi : X \to Y$ is small. Then the flip $\varphi^+ : X^+ \to Y$ exists. Moreover X^+ is a normal Q-factorial compact Kähler threefold and $(X^+, (\varphi^+)^{-1}_* \circ \varphi_* \Delta)$ is dlt.

Proof. – Since X is Q-factorial, the pair $(X, (1-\varepsilon)\Delta)$ is klt for every $0 < \varepsilon < 1$. Moreover for $0 < \varepsilon \ll 1$ the divisor $K_X + (1 - \varepsilon)\Delta$ is negative on the extremal ray R. Thus the flip $\varphi : X^+ \to Y$ exists by [62, Thm], see also [20] (and [49, 2.32] for passing to the limit). We alternatively may apply [49], where the existence of log flips is reduced to the existence of terminal flips, [55]. The existence of terminal flips is now a local analytic construction. The reduction to the terminal case as given in Sections 5, 6 and 8 in [49] works in the analytic setting as well⁽⁷⁾. By Corollary 3.1, the complex space Y is Kähler. Since the morphism φ^+ is projective, X^+ is again Kähler. For the remaining properties we refer to Proposition 3.1. □

While the existence of flips is a local property [51, 6.7], this is not necessarily the case for termination results. However in most cases we will only use special termination for dlt pairs which is much simpler to prove:

THEOREM 4.4 ([62, 49]). – Let X be a normal Q-factorial compact Kähler threefold. Let Δ be an effective reduced Weil divisor such that the pair (X, Δ) is dlt. Set $(X_0, \Delta_0) := (X, \Delta)$ and let

$$(\varphi_i : (X_i, \Delta_i) \dashrightarrow (X_{i+1}, \Delta_{i+1} := (\varphi_i)_* \Delta_i)_{i \in I}$$

be a sequence of $(K + \Delta)$ -flips where $I \subset \mathbb{N}$. Then for all $i \gg 0$ the flipping locus is disjoint from Δ_i .

In particular, if for every $i \in I$ the flipping locus of φ_i is contained in Δ_i , then the sequence of $(K + \Delta)$ -flips terminates.

Proof. – The proof of [27, Thm. 4.2.1] applies without changes: obviously the MMP exists for normal \mathbb{Q} -factorial compact Kähler *surfaces*, the discrepancy calculations are local properties, so they also apply in the non-algebraic setting.

THEOREM 4.5 ([43, 49]). – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold such that (X, 0) is klt. Set $X_0 := X$ and let

$$(\varphi_i: X_i \dashrightarrow X_{i+1})_{i \in I}$$

be a sequence of K-flips. Then I is finite, that is any sequence of K-flips terminates.

Proof. - The proof of [43, Thm. 1] is based on three tools:

- a) existence of terminal models for klt pairs [43, Thm. 5];
- b) crepant extraction of a unique divisor with nonpositive discrepancy [43, Lemma 6];
- c) discrepancy calculations.

⁽⁷⁾ It is important to notice that the construction is locally analytic near the contracted curves.

^{4°} SÉRIE - TOME 49 - 2016 - Nº 4

The first two tools are available in the Kähler setting since we can argue as in Subsection 2.A. Discrepancy calculations are local properties, so they also apply in the non-algebraic setting.

As an application we can run the MMP for certain dlt pairs:

THEOREM 4.6. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold that is not uniruled. Let $D \in |mK_X|$ be a pluricanonical divisor and set B := Supp D. Suppose that the pair (X, B) is dlt. Then there exists a terminating (K+B)-MMP, that is, there exists a bimeromorphic map

$$\varphi: (X, B) \dashrightarrow (X', B' := \varphi_*B)$$

which is a composition of K + B-negative divisorial contractions and flips such that X' is a normal \mathbb{Q} -factorial compact Kähler threefold, the pair (X', B') is dlt and $K_{X'} + B'$ is nef.

Proof. – Set $(X_0, B_0) := (X, B)$ and $D_0 := D$.

Step 1. Existence of the MMP. – If $K_{X_i} + B_i$ is nef, there is nothing to prove, so suppose that this is not the case. Then there exists by Theorem 1.3 a $K_{X_i} + B_i$ -negative extremal ray $R_i \in \overline{NA}(X_i)$. Since X_i is not uniruled, the extremal ray is divisorial or small.

If the extremal ray is small, the contraction $\psi_i : X_i \to Y_i$ exists by Theorem 4.2. By Theorem 4.3 the flip $\psi_i^+ : X_i^+ \to Y_i$ exists, and we denote by

$$\varphi_i : (X_i, B_i) \dashrightarrow (X_{i+1} := X_i^+, B_{i+1} := (\varphi_i)_* B_i)$$

the composition $(\psi_i^+)^{-1} \circ \psi_i$. By Theorem 4.3 we know that X_{i+1} is a normal Q-factorial Kähler space and (X_{i+1}, B_{i+1}) is dlt, so we can continue the MMP. Note finally that $D_{i+1} := (\varphi_i)_* D_i$ is a pluricanonical divisor such that $\operatorname{Supp} D_{i+1} = B_{i+1}$.

If the extremal ray is divisorial, let S_i be the unique surface such that $S_i \cdot R_i < 0$ (cf. Notation 4.1). Since $K_{X_i} + B_i$ is \mathbb{Q} -effective and represented by an effective \mathbb{Q} -divisor with support in B_i , the surface S_i must be an irreducible component of B_i . Since the pair (X_i, B_i) is dlt, any of the irreducible components of B_i is normal [51, 5.52]. Moreover, there exists a boundary divisor $B_{i,S}$ such that

$$K_{S_i} + B_{i,S} \sim_{\mathbb{Q}} K_{X_i} + B_i$$

and the pair $(S_i, B_{i,S})$ is slc [49, 16.9.1]. Thus by Proposition 4.5 the contraction

$$\varphi_i: (X_i, B_i) \to (X_{i+1}, B_{i+1}) := (\varphi_i)_* B_i$$

exists and X_{i+1} is a Kähler space by Corollary 3.1. Moreover by Proposition 3.1, (X_{i+1}, B_{i+1}) is dlt and X_{i+1} is Q-factorial, so we can continue the MMP. Note finally that $D_{i+1} := (\varphi_i)_* D_i$ is a pluricanonical divisor such that $\operatorname{Supp} D_{i+1} = B_{i+1}$.

Step 2. Termination. – For every $i \in I$ we have $\operatorname{Supp} D_i = B_i$, hence $K_{X_i} + B_i$ is \mathbb{Q} -effective and represented by an effective \mathbb{Q} -divisor with support in B_i . In particular the support of the extremal contraction is in B_i . Thus any sequence of (K+B)-flips terminates by Theorem 4.4.

5. Semi-log canonical Kähler surfaces

In this short section we gather the results concerning abundance in dimension 2 which will be used in the concluding sections. We refer to [49, Ch. 12] [48, Ch. 5] for basic definitions of the theory of semi-log canonical (slc) surfaces. All the statements in this section are shown for projective surfaces in [49, Ch. 12], we will see that the proofs also apply to our case with some minor modifications.

Let S be a reduced compact complex surface. We say that S is in class \mathcal{C} if the desingularizations of the irreducible components S_i of S are Kähler. The Kähler property is a bimeromorphic invariant of smooth compact surfaces, so this definition is independent of the choice of the desingularizations.

PROPOSITION 5.1. – Let S be an irreducible reduced compact complex surface in class C. Let Δ be an effective Q-Cartier divisor on S such that the pair (S, Δ) is slc. Suppose that $K_S + \Delta$ is nef and numerically trivial. Then $K_S + \Delta$ is torsion, i.e., there exists an $m \in \mathbb{N}$ such that

$$\mathcal{O}_S(m(K_S + \Delta)) \simeq \mathcal{O}_S.$$

Proof. – Let $\nu : \tilde{S} \to S$ be the normalization, and let $\tilde{\Delta}$ be the different (cf. Subsection 2.B), so that we have

$$K_{\tilde{S}} + \tilde{\Delta} \sim_{\mathbb{Q}} \nu^* (K_S + \Delta) \equiv 0,$$

and the pair $(\tilde{S}, \tilde{\Delta})$ is lc.

First case. $\tilde{\Delta} \neq 0$. – In this case the surface \tilde{S} is projective: indeed the anticanonical divisor $-K_S \equiv \Delta_S$ is pseudoeffective, so a K-MMP⁽⁸⁾ $\tilde{S} \rightarrow \bar{S}$ terminates with a Mori fibre space $\bar{S} \rightarrow W$. Since \bar{S} is a surface and $-K_{\bar{S}}$ is relatively ample we see that \bar{S} is projective. Since $\tilde{S} \rightarrow \bar{S}$ is a projective morphism, the surface \tilde{S} is projective. Thus the statements in [49, Ch. 12] apply.

Second case. $\tilde{\Delta} = 0$. – In this case S is normal, so if $\pi : \hat{S} \to S$ is the minimal resolution, then we have

$$\kappa(\hat{S}, K_{\hat{S}} + E) = \kappa(S, K_S),$$

where E is the canonical defined effective divisor such that $K_{\hat{S}} + E \sim_{\mathbb{Q}} \pi^* K_S \equiv 0$. Now if $E \neq 0$, then the smooth surface \hat{S} is uniruled, hence projective, and therefore $\kappa(\hat{S}, K_{\hat{S}} + E) = 0$ by [49, Thm. 12.1.1]. If E = 0, then \hat{S} is a compact Kähler surface with numerically trivial canonical bundle, so $\kappa(\hat{S}, K_{\hat{S}}) = 0$ by the Beauville-Bogomolov decomposition [8, Thm. 1].

LEMMA 5.1. – Let S be a normal compact Kähler surface, and let Δ be a boundary divisor on S. Suppose that the pair (S, Δ) is lc, that the class $K_S + \Delta$ is nef and that $\nu(K_S + \Delta) = 1$. Then $K_S + \Delta$ is semi-ample.

4° SÉRIE – TOME 49 – 2016 – Nº 4

⁽⁸⁾ The existence of an MMP for a compact Kähler surface with lc singularities can be established following our arguments in Section 4.

Proof. – Let $\mu : S' \to S$ be a dlt model, cf. Theorem 2.2, then (S', Δ') is dlt and $K_{S'} + \Delta' \sim_{\mathbb{Q}} \mu^*(K_S + \Delta)$. In particular $K_S + \Delta$ is semiample if and only if $K_{S'} + \Delta'$ is semiample. Thus we can assume without loss of generality that (S, Δ) is dlt.

Dlt singularities are rational [51, Thm. 5.22], so the surface S is projective if and only if some desingularization is projective [57]. Thus if the algebraic dimension a(S) is two, we conclude by [49, Thm. 12.1.1]. If a(S) = 1, let $f : S \to C$ be the algebraic reduction. The general fibre F is an elliptic curve, so $K_S|_F \simeq \mathcal{O}_F$. Moreover every curve in S is contracted by f, so $\Delta \cap F = \emptyset$. Since $K_S + \Delta$ is nef, this implies

$$K_S + \Delta \sim_{\mathbb{Q}} f^* A,$$

with A an ample \mathbb{Q} -divisor on C. Thus $K_S + \Delta$ is semiample. Finally we want to exclude the case a(S) = 0: note first that S contains only finitely many curves, so a desingularization $\hat{S} \to S$ is not uniruled. Thus we have

$$\kappa(K_S) \ge \kappa(K_{\hat{S}}) \ge 0,$$

in particular $K_S + \Delta$ is Q-linearly equivalent to an effective divisor. Yet the intersection form on a surface with a(S) = 0 is negative definite, hence $(K_S + \Delta)^2 < 0$. In particular $K_S + \Delta$ is not nef, a contradiction.

PROPOSITION 5.2. – Let S be a connected reduced compact complex surface in class C. Let Δ be an effective \mathbb{Q} -Cartier divisor on S such that the pair (S, Δ) is slc. Suppose that $K_S + \Delta$ is nef with numerical dimension $\nu(K_S + \Delta) \leq 1$. Suppose also that for every irreducible component $T \subset S$ the restriction $(K_S + \Delta)|_T$ is not zero.

Then the divisor $K_S + \Delta$ is semi-ample.

Proof. – Let $\nu : T \to S$ be the composition of a minimal semi-resolution (cf. [49, Defn.12.2.1, Prop. 12.2.3]) with the normalization. Since (S, Δ) is slc there exists for every irreducible component $T_i \subset T$ a boundary divisor Δ_i such that

$$K_{T_i} + \Delta_i \sim_{\mathbb{Q}} \nu^* (K_S + \Delta).$$

By our assumption $K_{T_i} + \Delta_i$ is nef of numerical dimension one, so $K_{T_i} + \Delta_i$ is semiample by Lemma 5.1. The whole point is now to prove that sufficiently many pluricanonical sections descend to S. However the proof of this descent theorem in [49, Ch. 12.4] is based on [49, Prop. 12.3.2, Thm. 12.3.4] which are statements about proper morphisms. Thus they apply in our setting.

THEOREM 5.1. – Let S be a compact connected reduced complex space of dimension two in class \mathcal{C} with slc singularities. Then the natural maps induced by $\mathbb{C} \subset \Theta_S$

$$H^p(S,\mathbb{C}) \to H^p(S,\mathcal{O}_S)$$

are surjective for every $p \in \mathbb{N}$.

Proof. – Simply note that the proof of Theorem 12.1.2 of [49] works practically word by word. The main ingredient [49, 12.2.8] is about germs of slc surfaces and works in our situation. \Box

6. Abundance: reduction steps

This section corresponds to the preparatory lemmas in [49, Ch. 13, Ch. 14]. While the statements and the basic strategy of proof are quite similar we have to be more careful since we do not have a full log-MMP at our disposal. Moreover we make several points more precise which allows us to conclude quicker in Section 8.

LEMMA 6.1. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold with terminal singularities. Suppose that K_X is nef and $\nu(X) > 0$. Then there exists a normal \mathbb{Q} -factorial compact Kähler threefold X' that is bimeromorphic to X and a $D' \in |mK_{X'}|$ with the following properties:

- a) Set B' := Supp D'. Then the pair (X', B') is dlt and $X' \setminus B'$ has terminal singularities.
- b) *The divisor* K_{X'} + B' *is nef and we have*(i) ν(X) = ν(K_{X'} + B'); and
 (ii) κ(X) = κ(K_{X'} + B').

The proof needs some technical preparation:

REMARK 6.1. – Let X be a normal Q-factorial compact Kähler space, and let $D \in |mK_X|$ be an effective pluricanonical divisor. If we set B = Supp D, then

$$\kappa(K_X) = \kappa(K_X + B).$$

Indeed, B being effective, the inequality $\kappa(K_X) \leq \kappa(K_X + B)$ is obvious. Yet we also have an inclusion of effective divisors $B \subset D$, so we get

$$\kappa(K_X + B) \le \kappa(K_X + D) = \kappa((m+1)K_X) = \kappa(X).$$

The following basic lemma has been shown in the algebraic case in [49, 13.2.4].

LEMMA 6.2. – Let $\psi : V \to W$ be a bimeromorphic proper Kähler morphism between normal complex spaces. Let D be an effective Q-Cartier Q-divisor that is ψ -nef. Then we have

$$\operatorname{Supp} \psi_*(D) = \psi(\operatorname{Supp} D),$$

i.e., the image of D has pure codimension one.

Proof. – The inclusion $\operatorname{Supp} \psi_*(D) \subset \psi(\operatorname{Supp} D)$ is trivial. Fix now a point $w \in \psi(\operatorname{Supp} D) \subset W$. Arguing by contradiction we assume that all the irreducible components of $\psi(\operatorname{Supp} D)$ passing through w have dimension at most dim W - 2. Thus, up to replacing W by an analytic neighborhood of w, we assume that $\psi(\operatorname{Supp} D)$ has dimension at most dim W - 2. Then $\psi_*(-D) = 0$ and -(-D) is ψ -nef, contradicting the negativity lemma, see e.g., [10, 3.6.2], unless D = 0.

The following crucial lemma seems to be well-known to experts and is used in several places without further mentioning, but we prefer to write it down in detail:

4° SÉRIE - TOME 49 - 2016 - Nº 4

LEMMA 6.3. – Let $X =: X_0$ be a normal Q-factorial compact Kähler space, and let $M =: M_0$ be an effective Cartier divisor on X that is nef. Let $N =: N_0$ be an effective Cartier divisor such that Supp M =Supp N, and let

$$(\varphi_i: X_i \dashrightarrow X_{i+1})_{i=0,\dots,n}$$

be a finite sequence of N-negative contractions, that is every X_i is a normal \mathbb{Q} -factorial compact Kähler space and φ_i is the divisorial contraction or flip of an extremal ray $R_i \in \overline{NA}(X_i)$ that is N_i -negative ⁽⁹⁾. Then for every i = 0, ..., n the N-MMP induces an isomorphism

$$(X_0 \setminus \operatorname{Supp} M_0) \simeq (X_{i+1} \setminus \operatorname{Supp} M_{i+1}).$$

REMARK. – The essence of Lemma 6.3 is the following. Given the flip $\varphi : X_i \longrightarrow X_{i+1}$ with contractions $f_i : X_i \longrightarrow Y_i$ and $f_i^+ : X_{i+1} \longrightarrow Y_i$, then the exceptional locus of f_i^+ is contained in M_{i+1} .

Proof. – For every i = 1, ..., n we set $M_{i+1} := (\varphi_i)_* M_i$. Moreover we denote by Γ_i the normalization of the graph of φ_i and by $p_i : \Gamma_i \to X_i$ and $q_i : \Gamma_i \to X_{i+1}$ the natural maps. We will construct inductively a sequence of normal compact Kähler spaces V_i admitting bimeromorphic morphisms $f_i : V_i \to X$ and $g_i : V_i \to X_{i+1}$ such that

- a) the support of $f_i^* M$ contains the g_i -exceptional locus;
- b) g_i factors through Γ_i , that is there exists a bimeromorphic map $\beta_i : V_i \to \Gamma_i$ such that $g_i = q_i \circ \beta_i$.

Assuming this for the time being, let us see how to conclude: arguing by induction we suppose that we have an isomorphism

$$(X_0 \setminus \operatorname{Supp} M_0) \simeq (X_i \setminus \operatorname{Supp} M_i).$$

Since Supp M_i = Supp N_i and φ_i is N_i -negative, the image of the p_i -exceptional locus is contained in M_i . Thus we are done if we show that the image of the q_i -exceptional locus is contained in M_{i+1} . Since $X_0 \dashrightarrow X_{i+1}$ does not extract a divisor we have

$$M_{i+1} = (g_i)_* f_i^* M.$$

Yet by the properties above f_i^*M contains the strict transforms of the q_i -exceptional divisors ⁽¹⁰⁾. Thus $g_i(\text{Supp } f_i^*M)$ contains the image of the q_i -exceptional locus. By Lemma 6.2 this implies that $\text{Supp}(g_i)_*f_i^*M$ contains the image of the q_i -exceptional locus.

Proof of the claim. – For the start of the induction we simply set $V_0 := \Gamma_0$. Indeed since Supp M_0 = Supp N_0 contains the exceptional locus of the extremal contraction, the divisor $p_0^*M_0$ contains the q_0 -exceptional locus.

For the induction step we make a case distinction: if φ_i is divisorial we simply set $V_i := V_{i-1}$ and note that M_i contains the exceptional divisor since $\operatorname{Supp} M_i = \operatorname{Supp} N_i$. If φ_i is a flip we define V_i as the normalization of the graph of the bimeromorphic map $V_{i-1} \dashrightarrow \Gamma_i$, and

⁽⁹⁾ As usual, we define inductively $N_{i+1} := (\varphi_i)_* N_i$, cf. [49, 2.26].

⁽¹⁰⁾ Note that the q_i -exceptional locus is divisorial since X_{i+1} is \mathbb{Q} -factorial.

denote by $\alpha_i : V_i \to V_{i-1}$ and $\beta_i : V_i \to \Gamma_i$ the natural maps. We set $f_i := f_{i-1} \circ \alpha_i$ and $g_i := q_i \circ \beta_i$ and summarize the situation in a commutative diagram:



All we have to show is that the support of f_i^*M contains the g_i -exceptional locus. Note that by our induction hypothesis f_{i-1}^*M contains the g_{i-1} -exceptional locus. Moreover we have

$$M_i = (g_{i-1})_* f_{i-1}^* M$$

and Supp M_i = Supp N_i . In particular $(g_{i-1})_* f_{i-1}^* M$ contains the image of the p_i -exceptional locus (which is of course equal to the q_i -exceptional locus). Using these two properties the claim follows by elementary set-theoretic computations.

Proof of Lemma 6.1. – Since $\kappa(X) \ge 0$ by [24], there exist an $m \in \mathbb{N}$ and an effective Cartier divisor $D \in |mK_X|$. Since $\nu(X) > 0$, the divisor D is not zero. Let

$$\mu: X_0 \to X$$

be a log-resolution of the divisor D in the following sense: the map μ is an isomorphism on $X \setminus D$, the support of $\mu^* D$ has simple normal crossings and X_0 is smooth in a neighborhood of $\mu^* D^{(11)}$. Thus if we set $B_0 := (\mu^* D)_{red}$, the pair (X_0, B_0) is dlt and $X_0 \setminus B_0$ has terminal singularities. Note also that since X has terminal singularities, we have

$$K_{X_0} \sim_{\mathbb{O}} \mu^* K_X + E$$

with E an effective μ -exceptional \mathbb{Q} -divisor. By construction

Supp
$$E \subset B_0$$
,

so the pluricanonical divisor

$$D_0 := \mu^* D + mE \in |mK_{X_0}|$$

satisfies Supp $D_0 = B_0$. By Remark 6.1 we have

$$\kappa(X) = \kappa(X_0) = \kappa(K_{X_0} + B_0).$$

By Theorem 4.6 there exists a terminating (K + B)-MMP

$$(\varphi_i: (X_i, B_i) \dashrightarrow (X_{i+1}, B_{i+1}))_{i=0,\dots,n}$$

where φ_i is the flip or divisorial contraction of a ray R_i . Then for every $i \in \{1, ..., n\}$

$$D_{i+1} := \varphi_* D_i \in |mK_{X_{i+1}}|$$

is a pluricanonical divisor such that Supp $D_{i+1} = B_{i+1}$.

 $^{(11)}$ This last condition can be satisfied since X is a terminal threefold, so it has only finitely many singular points.

4° SÉRIE - TOME 49 - 2016 - Nº 4

For every $0 \le i \le n$, if $C \subset X_i$ is a curve with class $[C] \in R_i$, then

$$(K_{X_i} + B_i) \cdot C = (\frac{1}{m}D_i + B_i) \cdot C < 0,$$

and since Supp $D_i = B_i$, the curve C is contained in B_i . Thus the exceptional locus of the contraction of the extremal ray R_i is contained in B_i . Now apply Lemma 6.3 to establish an isomorphism

$$\varphi_i|_{X_i \setminus B_i} : (X_i \setminus B_i) \to (X_{i+1} \setminus B_{i+1})$$

In conclusion we obtain that $X_{i+1} \setminus B_{i+1}$ has terminal singularities.

The Kodaira dimension of K + B being invariant under a (K + B)-MMP, so Remark 6.1 yields

$$\kappa(X) = \kappa(K_{X_0} + B_0) = \kappa(K_{X_{n+1}} + B_{n+1}).$$

Let now Γ be a desingularization of the graph of the bimeromorphic map $X_0 \dashrightarrow X_{n+1}$, and denote by $p: \Gamma \to X_0$ and $q: \Gamma \to X_{n+1}$ the natural projections. Then the effective divisors

$$(\mu \circ p)^* D \simeq (\mu \circ p)^* m K_X$$

and

$$q^*(D_{n+1} + mB_{n+1}) \simeq mq^*(K_{X_{n+1}} + B_{n+1})$$

are both nef and have the same support. As in [49, 11.3.3], arguing with a Kähler class, this implies

$$\nu(X) = \nu(K_{X_{n+1}} + B_{n+1}).$$

LEMMA 6.4. – Let X be a normal compact Kähler space of dimension $n \ge 2$, and let L be a nef \mathbb{Q} -Cartier \mathbb{Q} -divisor. Let $S \subset X$ be an effective \mathbb{Q} -Cartier \mathbb{Q} -divisor such that L - S is nef. Let $T \subset X$ be a prime divisor such that $L|_T \equiv 0$ and $T \not\subset \text{Supp } S$. Then

$$T \cap \operatorname{Supp} S = \emptyset.$$

Proof. – Indeed, if $T \cap \text{Supp } S \neq \emptyset$, then $-S|_T$ is a non-zero antieffective divisor. Thus the restriction $(L-S)|_T \equiv -S|_T$ is not nef, a contradiction.

LEMMA 6.5. – Let X be a normal \mathbb{Q} -factorial compact Kähler threefold. Suppose that there exists a $D \in |mK_X|$ with the following properties:

- Set B := Supp D. The pair (X, B) is dlt and $X \setminus B$ has terminal singularities.
- The divisor $K_X + B$ is nef.

Let $S \subset B$ be a non-zero effective Weil divisor such that the following holds: for every irreducible component $T \subset B - S$ we have

$$(K_X + B)|_{\mathcal{T}} \equiv 0.$$

Set $X_0 := X, D_0 := D$ and $B_0 := B$. Then there exists a finite sequence K + B - S-negative contractions

$$(\varphi_i: (X_i, B_i - S_i) \dashrightarrow (X_{i+1}, B_{i+1} - S_{i+1}))_{i=0,\dots,n}$$

of a ray $R_i \in \overline{NA}(X_i)$ such that the following properties hold:

a) The pair $(X_{i+1}, B_{i+1} - S_{i+1})$ is dlt.

b) Set $D_{i+1} := (\varphi_i)_* D_i$. Then the divisor $D_{i+1} \in |mK_{X_{i+1}}|$ satisfies $\operatorname{Supp} D_{i+1} = B_{i+1}$.

c) We have $(K_{X_i} + B_i) \cdot R_i = 0$ and the divisor $K_{X_{i+1}} + B_{i+1}$ is nef. For every irreducible component $T_{i+1} \subset B_{i+1} - S_{i+1}$ we have

$$(K_{X_{i+1}} + B_{i+1})|_{T_{i+1}} \equiv 0.$$

- d) X_{i+1} is \mathbb{Q} -factorial; the pair (X_{i+1}, B_{i+1}) is lc and $X_{i+1} \setminus B_{i+1}$ has terminal singularities.
- e) We have $(B_{n+1} S_{n+1}) \cap S_{n+1} = \emptyset$ and $S_{n+1} \neq 0$.

REMARK. – Note that we do not claim that $K_{X_{n+1}} + B_{n+1} - S_{n+1}$ is nef, in general this will not be true.

Proof. – If $(B - S) \cap S = \emptyset$, there is nothing to prove, so suppose $(B - S) \cap S \neq \emptyset$. Thus $K_X + B - S$ is not nef by Lemma 6.4. Note that $(X_0, B_0 - S_0)$ is dlt.

Step 1. Existence of the MMP. – We proceed by induction and assume that X_i , B_i and D_i are already constructed such that properties a), b) and d) hold at level *i*, moreover $K_{X_i} + B_i$ is nef and $K_{X_i} + B_i|_{T_i} \equiv 0$ for all irreducible components T_i of $B_i - S_i$. If there is no such T_i meeting S_i , then we stop and set n + 1 = i. Thus we may assume that $T_i \cap S_i \neq \emptyset$ for some component T_i . We first have to find the extremal ray R_i . Notice that the restriction

$$\left(K_{X_i} + B_i - S_i\right)_{|_{T_i}} \equiv -S_i|_{T_i}$$

is not nef, so by Corollary 4.1 there exists a curve $Z_i \subset T_i \subset (B_i - S_i)$ such that

$$(K_{X_i} + B_i - S_i) \cdot Z_i < 0.$$

The inclusion $Z_i \subset (B_i - S_i)$ yields $(K_{X_i} + B_i) \cdot Z_i = 0$ by c). By the cone Theorem 1.3 applied to the pair $(X_i, B_i - S_i)$, there exists a decomposition

(18)
$$Z_i = \sum \lambda_j C_{i,j} + M_i,$$

where $\lambda_j > 0$, the $C_{i,j}$ are irreducible curves generating a $(K_{X_i} + B_i - S_i)$ -negative extremal in $\overline{NA}(X_i)$ and $M_i \in \overline{NA}(X_i)$ such that

$$(K_{X_i} + B_i - S_i) \cdot M_i \ge 0.$$

Let R_i be the extremal ray generated by the curve $C_{i,1}$. Since $K_{X_i} + B_i$ is nef and $(K_{X_i} + B_i) \cdot Z_i = 0$, the decomposition (18) implies that $(K_{X_i} + B_i) \cdot R_i = 0$.

Since Supp $D_i = B_i$, the Q-Cartier divisor $K_{X_i} + B_i - S_i$ is Q-linearly equivalent to an effective divisor with support in B_i . In particular every curve $C \subset X_i$ such that $[C] \in R_i$ is contained in B_i . Since the pair (X_i, B_i) is lc this shows that the support of the extremal ray is contained in an irreducible surface with slc singularities, [49, 16.9]. Thus the contraction of R_i exists by Theorem 1.4. Moreover, if the ray is small, then the flip exists by Theorem 4.3. We denote by

$$\varphi_i: (X_i, B_i - S_i) \dashrightarrow (X_{i+1}, B_{i+1} - S_{i+1})$$

the flip or divisorial contraction of R_i . As in the proof of Lemma 6.1 we see that the properties a) and b) hold at level i + 1.

Since $(K_{X_i} + B_i) \cdot R_i = 0$, Lemma 3.2 proves that $K_{X_{i+1}} + B_{i+1}$ is nef, that the pair (X_{i+1}, B_{i+1}) is lc. and that for every irreducible component $T_{i+1} \subset B_{i+1} - S_{i+1}$, the restriction $(K_{X_{i+1}} + B_{i+1})|_T \equiv 0$. Finally, the divisor $K_{X_i} + B_i$ being nef and Q-linearly

4° SÉRIE - TOME 49 - 2016 - Nº 4

equivalent to an effective divisor with support in B_i , Lemma 6.3 implies that $(X_{i+1} \setminus B_{i+1}) \simeq (X_i \setminus B_i)$ has terminal singularities.

Step 2. Termination of the MMP. – By Theorem 4.4 we know that after finitely many steps the flipping locus is disjoint from $B_i - S_i$. Yet a K + B - S-negative extremal contraction that is disjoint from B - S is also a K-negative extremal contraction. Thus the sequence terminates by Theorem 4.5.

LEMMA 6.6. – In the situation of Lemma 6.5 suppose additionally that $\nu(K_X + B) = 1$, and let $S \subset B$ be an irreducible component.

Then there exists a normal \mathbb{Q} -factorial compact Kähler threefold X' that is bimeromorphic to X and has a $D' \in |mK_{X'}|$ with the following properties:

- a) Set B' := Supp D'. The pair (X', B') is lc and $X' \setminus B'$ has terminal singularities.
- b) The divisor $K_{X'} + B'$ is nef with $\nu(K_{X'} + B') = 1$. Moreover, $\kappa(K_X + B) = \kappa(K_{X'} + B')$.
- c) There exists an irreducible component $S' \subset B'$ that is a connected component of B'.
- d) The pair (X', B' S') is dlt.

Proof. – We first check that B - S satisfies the positivity condition in Lemma 6.5: indeed if T is any irreducible component of $\operatorname{Supp}(D)$, then $(K_X + B)|_T \equiv 0$, otherwise $\nu(K_X + B) \ge 2$. Using the notation of Lemma 6.5, set $X' = X_{n+1}$ etc. We know that S_{n+1} is a non-zero irreducible divisor that is disjoint from $B_{n+1} - S_{n+1}$. Thus the property c) holds. Since all the contractions in Lemma 6.5 are (K + B)-trivial, the Kodaira dimension is invariant:

$$\kappa(K_X + B) = \kappa(K_{X_{n+1}} + B_{n+1}).$$

The other properties were already shown in Lemma 6.5.

LEMMA 6.7. – Let X be a normal compact Kähler space with rational singularities of dimension $n \ge 2$. Let D be an effective Cartier divisor. Suppose that D is nef and $\nu(D) \ge 2$. Then the support of D is connected.

Proof. – It is sufficient to prove that $H^1(X, \mathcal{O}_X(-D)) = 0$. Since X has rational singularities we may replace it by a resolution of singularities without changing this cohomology group. Then we have

$$H^1(X, \mathcal{O}_X(-D)) \simeq H^{n-1}(X, K_X \otimes \mathcal{O}_X(D)) = 0$$

by [24, Thm. 0.1].

LEMMA 6.8. – In the situation of Lemma 6.5 suppose additionally that $\nu(K_X + B) = 2$. Then there exists a normal Q-factorial compact Kähler threefold X', bimeromorphic to X such that (X', 0) is klt and a divisor $D' \in |mK_{X'}|$ with the following properties:

- a) Set B' := Supp D'. The pair (X', B') is lc and $X' \setminus B'$ has terminal singularities.
- b) The divisor $K_{X'} + B'$ is nef with $\nu(K_{X'} + B') = 2$. Moreover, $\kappa(K_X + B) = \kappa(K_{X'} + B')$.
- c) For every irreducible component $T' \subset B'$ we have $(K_{X'} + B')|_{T'} \neq 0$.
- d) $(K_{X'} + B') \cdot K_{X'}^2 \ge 0.$

Proof. – The construction of the Kähler space X' proceeds in two steps.

ANNALES SCIENTIFIQUES DE L'ÉCOLE NORMALE SUPÉRIEURE

Part I. Eliminating the (K + B)-trivial components. – Let $S \subset B$ be the union of all the irreducible components $T \subset B$ such that $(K_X + B)|_T \neq 0$. Since $\nu(K_X + B) = 2$, the set S is not empty. Thus we can apply Lemma 6.5 to obtain a sequence of (K + B - S)-negative contractions

$$\psi: X \dashrightarrow X$$

where \hat{X} is a normal \mathbb{Q} -factorial compact Kähler threefold carrying a divisor $\hat{D} \in |mK_{\hat{X}}|$ subject to the following properties:

(α) Set $\hat{B} := \operatorname{Supp} \hat{D}$. The pair (\hat{X}, \hat{B}) is lc and $\hat{X} \setminus \hat{B}$ has terminal singularities.

(β) The divisor $K_{\hat{X}} + \hat{B}$ is nef with $\nu(K_{\hat{X}} + \hat{B}) = 2$. Moreover, $\kappa(K_X + B) = \kappa(K_{\hat{X}} + \hat{B})$.

- (γ) Set $\hat{S} := \psi_* S$. Then the pair $(\hat{X}, \hat{B} \hat{S})$ is dlt.
- (δ) supp $(\hat{B} \hat{S}) \cap \hat{S} = \emptyset$ and $\hat{S} \neq 0$.

Notice also that $K_{\hat{X}} + \hat{B}$ is Q-linearly equivalent to an effective divisor with support in \hat{B} . By Lemma 6.7 this implies that $\hat{B} = \hat{S} + (\hat{B} - \hat{S})$ is connected. Thus property (δ) implies that $\hat{B} - \hat{S} = 0$. Hence $(K_{\hat{X}} + \hat{B})|_T$ is non-zero for every irreducible component $T \subset \hat{B}$.

Part II. Eliminating K-negative curves. - Setting

$$X_0 := \hat{X}, \ B_0 := \hat{B}, \ D_0 := \hat{D},$$

we will next construct a finite sequence of K-negative contractions and flips

$$(\varphi_i: X_i \dashrightarrow X_{i+1})_{i=0,\dots,n}$$

of extremal rays $R_i \in \overline{NA}(X_i)$ such that the following properties hold:

- (1) X_{i+1} is \mathbb{Q} -factorial and $(X_{i+1}, 0)$ is klt.
- (2) Set $D_{i+1} := (\varphi_i)_* D_i$. Then $D_{i+1} \in |mK_{X_{i+1}}|$ satisfies $\text{Supp } D_{i+1} = B_{i+1}$.
- (3) We have $(K_{X_i} + B_i) \cdot R_i = 0$, the divisor $K_{X_{i+1}} + B_{i+1}$ is nef and $\nu(K_{X_{i+1}} + B_{i+1}) = 2$.
- (4) The pair (X_{i+1}, B_{i+1}) is lc and $X_{i+1} \setminus B_{i+1}$ has terminal singularities.
- (5) For every irreducible component $T \subset B_{i+1}$ we have $(K_{X_{i+1}} + B_{i+1})|_T \neq 0$.
- (6) $(K_{X_{n+1}} + B_{n+1}) \cdot K_{X_{n+1}}^2 \ge 0.$

Note first that $(\hat{X}, 0)$ is klt singularities since $(\hat{X}, \hat{B} - \hat{S}) = (\hat{X}, 0)$ is dlt [51, Prop. 2.41]. If $(K_{\hat{X}} + \hat{B}) \cdot K_{\hat{X}}^2 \ge 0$, there is nothing to prove, thus we may assume that

$$(K_{\hat{X}} + B) \cdot K_{\hat{X}}^2 < 0.$$

We therefore start by showing that $(K_{X_i} + B_i) \cdot K_{X_i}^2 < 0$ implies the existence of a K_{X_i} -negative contraction that is $(K_{X_i} + B_i)$ -trivial. Let $B_i = \sum B_{i,l}$ be the decomposition of B_i in its irreducible components. Since the pair (X_i, B_i) is lc, the surfaces $B_{i,l}$ have slc singularities. Moreover, by adjunction [49, Prop. 16.9], there exists a boundary divisor $\Delta_{i,l}$ such that

$$K_{B_{i,l}} + \Delta_{i,l} \sim_{\mathbb{Q}} (K_{X_i} + B_i)|_{B_{i,l}}$$

and the pair $(B_{i,l}, \Delta_{i,l})$ is slc. By Proposition 5.2 this implies that

$$(K_{X_i}+B_i)|_{B_i} \sim_{\mathbb{Q}} Z_{i,l}$$

with $Z_{i,l}$ an effective 1-cycle.

4° SÉRIE – TOME 49 – 2016 – Nº 4

Suppose that $K_{X_i|_{B_i,l}} \cdot Z_{i,l} \ge 0$ for all *l*. Since

$$K_{X_i} \sim_{\mathbb{Q}} \sum a_{i,l} B_{i,l}$$

with $a_{i,l} > 0$ we conclude

$$(K_{X_i} + B_i) \cdot K_{X_i}^2 = \sum a_{i,l} (K_{X_i} + B_i) \cdot K_{X_i} \cdot B_{i,l} = \sum a_{i,l} K_{X_i}|_{B_{i,l}} \cdot Z_{i,l} \ge 0,$$

a contradiction. Thus we can suppose (up to renumbering) that

$$K_{X_i|_{B_{i-1}}} \cdot Z_{i,1} < 0.$$

By the cone Theorem 1.3 applied to the pair $(X_i, 0)$, there exists a decomposition

(19)
$$Z_{i,1} = \sum \lambda_j C_{i,j} + M_i,$$

where $\lambda_j > 0$, the $C_{i,j}$ are irreducible curves generating a K_{X_i} -negative extremal in $\overline{NA}(X_i)$ and where $M_i \in \overline{NA}(X_i)$ is a class satisfying $K_{X_i} \cdot M_i \ge 0$.

Notice now the following: since $K_{X_i} + B_i$ is \mathbb{Q} -linearly equivalent to an effective divisor with support B_i and $(K_{X_i} + B_i)^3 = 0$ we have

$$(K_{X_i} + B_i)^2 \cdot B_{i,l} = 0$$

for all l. In particular $(K_{X_i} + B_i) \cdot Z_{i,1} = 0$. Since $K_{X_i} + B_i$ is nef we deduce from (19) that

(20)
$$(K_{X_i} + B_i) \cdot C_{i,1} = 0.$$

Let now R_i be the extremal ray generated by $C_{i,1}$. As in the proof of Lemma 6.5, the locus of R_i is contained in a surface with slc singularities (a component of B_i). Thus the contraction of R_i exists by Theorem 1.4 and, if the extremal ray is small, the flip exists by Theorem 4.3. We denote by

$$\varphi_i: (X_i, 0) \dashrightarrow (X_{i+1}, 0)$$

the flip or divisorial contraction of R_i . As in the proof of Lemma 6.1 we see that the properties (1) and (2) hold at level i + 1.

By the induction hypothesis, the divisor $K_{X_i} + B_i$ is nef of numerical dimension two and by (20) numerically trivial on the extremal ray R_i . Using Lemma 3.2 this implies that $K_{X_{i+1}} + B_{i+1}$ is nef of numerical dimension two, moreover the pair (X_{i+1}, B_{i+1}) is lc. Finally Lemma 6.3 implies that $(X_{i+1} \setminus B_{i+1}) \simeq (X_i \setminus B_i)$ has terminal singularities.

Part III In total we have constructed a sequence of K-negative contractions satisfying the properties (1)-(5). By Theorem 4.5 any such sequence terminates after n steps. By construction this yields

$$(K_{X_{n+1}} + B_{n+1}) \cdot K_{X_{n+1}}^2 \ge 0.$$

Since all the contractions are K + B-trivial, the Kodaira dimension is invariant, i.e., we have $\kappa(K_{\hat{X}} + \hat{B}) = \kappa(K_{X_{n+1}} + B_{n+1})$.

7. Positivity of cotangent sheaves

In this section we briefly review stability and Chern classes on singular complex spaces, then we prove the crucial Chern class inequality Theorem 7.2. For simplicity of notations we restrict ourselves to the case of threefolds with isolated singularities—which is all we need later—but all the statements can easily be adapted for spaces of arbitrary dimension that are smooth in codimension two.

DEFINITION 7.1. – Let X be a normal compact Kähler threefold with isolated singularities. Let \mathcal{F}_1 and \mathcal{F}_2 be coherent sheaves on X, and let $\pi : \hat{X} \to X$ be a log-resolution.

Then the Chern classes $c_i(\pi^*\mathcal{F}_1)$ and $c_i(\pi^*\mathcal{F}_2)$ are well-defined elements of $H^{2i}(\hat{X}, \mathbb{Z})$ [63], see also [34]. Thus for every $\alpha \in N^1(X)$ the intersection numbers

$$\pi^* \alpha \cdot c_1(\pi^* \mathcal{F}_1) \cdot c_1(\pi^* \mathcal{F}_2)$$
 and $\pi^* \alpha \cdot c_2(\pi^* \mathcal{F}_1)$

are well-defined, so by the duality $N_1(X) = N^1(X)^*$ we define

$$c_1(\mathcal{F}_1) \cdot c_1(\mathcal{F}_2) \in N_1(X) : \alpha \mapsto \pi^* \alpha \cdot c_1(\pi^* \mathcal{F}_1) \cdot c_1(\pi^* \mathcal{F}_2)$$

and

$$c_2(\mathcal{F}_1) \in N_1(X) : \alpha \mapsto \pi^* \alpha \cdot c_2(\pi^* \mathcal{F}_1).$$

LEMMA 7.1. – In the situation of Definition 7.1, the classes $c_1(\mathcal{F}_1) \cdot c_1(\mathcal{F}_2)$ and $c_2(\mathcal{F}_1)$ do not depend on the choice of the desingularization. Moreover if

 $0 \to \mathcal{F} \to \mathcal{G} \to \mathcal{Q} \to 0$

is an exact sequence of coherent sheaves, then we have the usual formula

$$c_2(\mathcal{G}) = c_2(\mathcal{F}) + c_2(\mathcal{Q}) + c_1(\mathcal{F}) \cdot c_1(\mathcal{Q}).$$

Proof. – In order to see for instance that the definition $c_2(\mathcal{F})$ does not depend on the resolution it suffices to consider the case where $\pi_1 : \hat{X}_1 \to X$ and $\pi_2 : \hat{X}_2 \to X$ are two log-resolutions with a factorization $q : X_2 \to X_1$. Then we want to show that

$$(q^*\pi_1^*\alpha) \cdot c_2(q^*\pi_1^*\mathcal{F}) = (\pi_1^*\alpha) \cdot c_2(\pi_1^*\mathcal{F}).$$

This follows from the general fact, that, given a holomorphic map $f : X \to Y$ between compact complex manifolds and \mathcal{G} a coherent sheaf on Y, then $c_j(f^*(\mathcal{G})) = f^*(c_j(\mathcal{G}))$, cf. [34].

Next consider a log-resolution $\pi : \hat{X} \to X$ with exceptional locus $D = \sum D_k$ and \mathcal{F} a coherent sheaf with support on D. Let $i^k : D_k \to X$ be the inclusion. Then by the Grothendieck-Riemann-Roch formula, see e.g., [34, Thm1.1] (and [32, Thm. 15.2] in the algebraic case),

$$\operatorname{ch}(i_*^k(\mathcal{F}_{\mid D_k}) = i_*^k(\operatorname{td}(N_{D_k/X})^{-1} \cdot \operatorname{ch}(\mathcal{F}_{\mid D_k})).$$

Thus if $\alpha \in N^1(X)$, then the projection formula gives

$$\pi^* \alpha \cdot c_2(i_*^k(\mathcal{F}_{|_{D_*}}) = 0.$$

Now an easy induction on the number of components of D shows that

(21)
$$\pi^*(\alpha) \cdot c_2(\mathcal{F}) = 0$$

4° SÉRIE – TOME 49 – 2016 – Nº 4

For the second statement we consider the pull-back

$$\pi^* \mathcal{F} \xrightarrow{\alpha} \pi^* \mathcal{G} \to \pi^* \mathcal{Q} \to 0$$

by a log-resolution $\pi : \hat{X} \to X$. In general the morphism α is not injective, but its kernel has support in the π -exceptional locus. Using the usual rules for computing Chern classes on compact manifolds and invoking (21), we have

$$\pi^* \alpha \cdot c_2(\pi^* \mathcal{F}) = \pi^* \alpha \cdot c_2(\pi^* \mathcal{F} / \ker \alpha)$$

and

$$\pi^* \alpha \cdot c_1(\pi^* \mathcal{G}) \cdot c_1(\pi^* \mathcal{Q}) = \pi^* \alpha \cdot c_1(\pi^* \mathcal{G} / \ker \alpha) \cdot c_1(\pi^* \mathcal{Q})$$

Thus the statement follows from the standard formula in the smooth case.

REMARK 7.1. – In the proof of Lemma 7.1 we have shown the following. Let $\pi : \hat{X} \to X$ be a log resolution of the Kähler threefold X with only isolated singularities. Let \mathcal{F} be a coherent sheaf on \hat{X} , supported on the exceptional divisor D of π and let $\alpha \in N^1(X)$. Then we have

$$\pi^*(\alpha) \cdot c_2(\mathcal{F}) = \pi^*(\alpha) \cdot c_1(\mathcal{F})^2 = 0.$$

Given a normal compact Kähler threefold X with isolated singularities and a torsion-free sheaf \mathcal{F} there is no obvious candidate for the first Chern class $c_1(\mathcal{F}) \in H^2(X, \mathbb{R})$. However we can define, as in the situation of Definition 7.1, for every $\alpha \in N^1(X)$ the intersection number $\alpha^2 \cdot c_1(\mathcal{F})$ by pulling-back to some log-resolution $\pi : \hat{X} \to X$:

$$\alpha^2 \cdot c_1(\mathcal{F}) := (\pi^*(\alpha))^2 \cdot c_1(\pi^*(\mathcal{F})).$$

DEFINITION 7.2. – Let X be a normal compact Kähler threefold with isolated singularities and let α be a nef class on X. We say that a non-zero torsion-free sheaf \mathcal{F} is α -semistable (resp. α -stable) if for every non-zero saturated subsheaf $\mathcal{E} \subset \mathcal{F}$ we have

$$\mu_{\alpha}(\mathcal{E}) := \frac{\alpha^2 \cdot c_1(\mathcal{E})}{\mathrm{rk}\mathcal{E}} \leq \frac{\alpha^2 \cdot c_1(\mathcal{F})}{\mathrm{rk}\mathcal{F}} =: \mu_{\alpha}(\mathcal{F}) \qquad (\textit{resp. } <).$$

PROPOSITION 7.1 (Harder-Narasimhan filtration). – Let X be a normal compact Kähler threefold with isolated singularities, and let α be a nef class on X. Let \mathcal{F} be a non-zero torsion-free coherent sheaf on X. Then there exists a filtration

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \cdots \subset \mathcal{F}_k = \mathcal{F}$$

such that for every $i \in \{1, ..., k\}$ the quotient $\mathcal{F}_i / \mathcal{F}_{i-1}$ is α -semistable and we have a strictly decreasing sequence of slopes

$$\mu_{\alpha}(\mathcal{F}_{i}/\mathcal{F}_{i-1}) > \mu_{\alpha}(\mathcal{F}_{i+1}/\mathcal{F}_{i}) \qquad \forall i \in 1, \dots, k-1.$$

Proof. – We proceed as in [44]; the main point is to show that there is a constant C such that

$$\mu_{\alpha}(\mathcal{F}) \le C$$

for all $\mathcal{G} \subset \mathcal{E}$. First we reduce to the smooth case: take a log resolution $\pi : \hat{X} \to X$ such that $\pi^*(\mathcal{E})^{**}$ is locally free and observe that $\mu_{\alpha}(\mathcal{G}) = \mu_{\pi^*(\alpha)}(\pi^*(\mathcal{G})/\text{tor}))$.

So we may assume X smooth and \mathcal{E} locally free. Then the proof of [44, Lemma 7.16] works. Once the boundedness is settled, the existence of the filtration is shown as in the classical case.

We shall also use the following elementary result, cf. [44, Thm. 7.18].

PROPOSITION 7.2 (Jordan-Hölder filtration). – Let X be a normal compact Kähler threefold with isolated singularities, and let α be a nef class on X. Let \mathcal{F} be a non-zero α -semi-stable torsion-free coherent sheaf on X. Then there exists a filtration

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \cdots \subset \mathcal{F}_k = \mathcal{F}$$

such that for every $i \in \{1, ..., k\}$ the quotient $\mathcal{F}_i / \mathcal{F}_{i-1}$ is α -stable.

DEFINITION 7.3. – Let X be a normal compact Kähler threefold, and let α be a nef class on X. A non-zero torsion-free coherent sheaf \mathcal{F} on X is α -generically nef if for every torsion-free quotient sheaf $\mathcal{F} \to Q \to 0$ we have

$$\alpha^2 \cdot c_1(Q) \ge 0.$$

REMARK 7.2. – In the situation of Definition 7.3, let

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \cdots \subset \mathcal{F}_k = \mathcal{F}$$

be the Harder-Narasimhan filtration with respect to α . If \mathcal{F} is α -generically nef, then by definition $\alpha^2 \cdot c_1(\mathcal{F}/\mathcal{F}_{k-1}) \geq 0$. Thus

$$\alpha^2 \cdot c_1(\mathcal{F}_i/\mathcal{F}_{i-1}) \ge 0$$

for every $i \in \{1, \ldots, k\}$.

We can now prove the Bogomolov inequality for stable sheaves on a singular space.

THEOREM 7.1. – Let X be a normal compact Kähler threefold with isolated singularities, and let α be a Kähler class on X. Let \mathcal{F} be an α -stable non-zero torsion-free coherent sheaf on X. Then we have

$$\alpha \cdot c_2(\mathcal{F}) \ge \left(\frac{r-1}{2r}\right) \alpha \cdot c_1^2(\mathcal{F}).$$

Proof. – We fix a log-resolution $\pi : \hat{X} \to X$.

Step 1. Suppose that \mathcal{F} is reflexive. – Set $\hat{\mathcal{F}} := (\pi^* \mathcal{F})^{**}$. Since \mathcal{F} is reflexive, the sheaves $\hat{\mathcal{F}}$ and $\pi^* \mathcal{F}$ coincide in the complement of the π -exceptional locus. In particular by Remark 7.1 one has

$$\alpha \cdot c_2(\mathcal{F}) = \pi^* \alpha \cdot c_2(\pi^* \mathcal{F}) = \pi^* \alpha \cdot c_2(\mathcal{F})$$

and

$$\alpha \cdot c_1^2(\mathcal{F}) = \pi^* \alpha \cdot c_1^2(\pi^* \mathcal{F}) = \pi^* \alpha \cdot c_1^2(\hat{\mathcal{F}}).$$

Thus it is sufficient to prove the inequality for $\hat{\mathcal{F}}$. Arguing exactly as in the proof of [24, Prop. 6.9] we see that $\hat{\mathcal{F}}$ is $\pi^* \alpha$ -stable. Since stability is an open property [17, Prop. 2.1] we obtain that $\hat{\mathcal{F}}$ is $(\pi^* \alpha + \varepsilon \omega)$ -stable where ω is a Kähler form on \hat{X} and $0 < \varepsilon \ll 1$. Thus [5] yields

$$(\pi^*\alpha + \varepsilon\omega) \cdot c_2(\hat{\mathcal{F}}) \ge \frac{r-1}{2r} (\pi^*\alpha + \varepsilon\omega) \cdot c_1^2(\hat{\mathcal{F}})$$

4° SÉRIE - TOME 49 - 2016 - Nº 4

for every $0 < \varepsilon \ll 1$. The claim follows by passing to the limit $\varepsilon \to 0$.

Step 2. Reduction to the case where \mathcal{F} is reflexive. – Since \mathcal{F} is torsion-free, the injection $i: \mathcal{F} \hookrightarrow \mathcal{F}^{**}$ is an isomorphism in codimension one. Thus the kernel and cokernel of

$$\pi^*(\mathcal{F}) \to \pi^*(\mathcal{F}^{**})$$

have support in the union of the π -exceptional locus and a set of dimension at most one. In particular $c_1(\pi^*\mathcal{F}) = c_1(\pi^*\mathcal{F}^{**}) + D$ with D a π -exceptional divisor, so the projection formula yields

$$\alpha \cdot c_1^2(\mathcal{F}) = \alpha \cdot c_1^2(\mathcal{F}^{**}).$$

Thus we are done if we prove that $\alpha \cdot c_2(\mathcal{F}^{**}) \ge \alpha \cdot c_2(\mathcal{F})$. By the second part of Lemma 7.1 it is sufficient to prove that

$$\alpha \cdot c_2(\mathcal{F}^{**}/\mathcal{F}) = \pi^*(\alpha) \cdot c_2(\pi^*(\mathcal{F}^{**}/\mathcal{F})) \ge 0.$$

Let S be the union of the 1-dimensional irreducible components of the support of $\mathcal{F}^{**}/\mathcal{F}$ and \hat{S} the strict transform in \hat{X} ; we may assume that the irreducible components \hat{S}_i of \hat{S} are smooth. Set

$$Q := (i_{\hat{S}})_* (\pi^* (\mathcal{F}^{**} / \mathcal{F})_{|_{\hat{S}}}).$$

Then by Remark 7.1 and Lemma 7.1, it suffices to show

$$\pi^*(\alpha) \cdot c_2(Q) \ge 0.$$

Yet Q has support on a set of dimension one, so the Grothendieck-Riemann-Roch formula yields

$$c_2(Q) = \sum a_i \hat{S}_i,$$

where $a_i \in \mathbb{N}_0$. Since α is nef, the statement follows.

LEMMA 7.2. – Let X be a compact Kähler threefold with isolated singularities. Let α be a nef class on X, and let \mathcal{F} be a coherent sheaf. Then we have

$$(\alpha^2 \cdot c_1(\mathcal{F}))^2 \ge (\alpha \cdot c_1^2(\mathcal{F})) \cdot \alpha^3.$$

Proof. – By our definition of the intersection numbers 7.1 we may suppose X smooth. Since α is a limit of Kähler classes, it is sufficient to prove the statement under the stronger hypothesis that α is a Kähler class. Now the inequality follows from the usual Hodge index theorem [66, Thm. 6.33] by an elementary computation.

THEOREM 7.2. – Let (X, ω) be a compact Kähler threefold with isolated singularities, and let \mathcal{F} be a non-zero reflexive coherent sheaf on X such that det \mathcal{F} is Q-Cartier. Suppose that there exists a pseudoeffective class $P \in N^1(X)$ such that

$$L := c_1(\mathcal{F}) + P$$

is a nef class. Suppose furthermore that for all $0 < \varepsilon \ll 1$ the sheaf \mathcal{F} is $(L + \varepsilon \omega)$ -generically nef. Then we have

$$L \cdot c_2(\mathcal{F}) \ge \frac{1}{2}(L \cdot c_1^2(\mathcal{F}) - L^3).$$

In particular, if $L \cdot c_1^2(\mathcal{F}) \geq 0$ and $L^3 = 0$, then

(22)
$$L \cdot c_2(\mathcal{F}) \ge 0.$$

REMARK. – This statement (and its proof) is a variation of [53, Thm. 6.1], [49, Prop. 10.12]. However the weaker assumptions will be crucial for the application in the proof of Theorem 8.2.

Proof. – Fix $0 < \varepsilon \ll 1$, and consider the Harder-Narasimhan filtration

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \cdots \subset \mathcal{F}_k = \mathcal{F}$$

for \mathcal{F} with respect to $L_{\varepsilon} := L + \varepsilon \omega$. Then by Lemma 7.1

(23)
$$L_{\varepsilon} \cdot c_2(\mathcal{F}) = L_{\varepsilon} \cdot \left(\sum_{1 \le i < j \le k} c_1(\mathcal{F}_i/\mathcal{F}_{i-1})c_1(\mathcal{F}_j/\mathcal{F}_{j-1}) + \sum_{i=1}^k c_2(\mathcal{F}_i/\mathcal{F}_{i-1}) \right).$$

Since \mathcal{F} is L_{ε} -generically nef, by Remark 7.2

(24)
$$L^{2}_{\varepsilon} \cdot c_{1}(\mathcal{F}_{i}/\mathcal{F}_{i-1}) \geq 0 \qquad \forall i \in \{1, \dots, k\}.$$

For every $i \in \{1, \ldots, k\}$, let

$$0 = \mathcal{F}_{i,0} \subset \mathcal{F}_{i,1} \subset \cdots \subset \mathcal{F}_{i,k_i} = \mathcal{F}_i / \mathcal{F}_{i-1}$$

be the Jordan-Hölder filtration of $\mathcal{F}_i/\mathcal{F}_{i-1}$ with respect to L_{ε} . Then we have

(25)
$$L_{\varepsilon} \cdot c_{2}(\mathcal{F}_{i}/\mathcal{F}_{i-1}) = L_{\varepsilon} \cdot \left(\sum_{1 \le p < q \le k_{i}} c_{1}(\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1})c_{1}(\mathcal{F}_{i,q}/\mathcal{F}_{i,q-1}) + \sum_{p=1}^{k_{i}} c_{2}(\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1}) \right).$$

Since $\mathcal{F}_i/\mathcal{F}_{i-1}$ is L_{ε} -semistable with non-negative slope by (24), we obtain

(26)
$$L^2_{\varepsilon} \cdot c_1(\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1}) \ge 0 \qquad \forall i \in \{1, \dots, k\}, p \in \{1, \dots, k_i\}.$$

Plugging the Equation (25) into the Equation (23) and following the lexicographic order, we rename the graded pieces $\mathcal{F}_{i,p}/\mathcal{F}_{i,p-1}$ into \mathcal{G}_l , where $l \in \{1, \ldots, n\}$. Thus

(27)
$$L_{\varepsilon} \cdot c_2(\mathcal{F}) = L_{\varepsilon} \cdot \left(\sum_{1 \le l < m \le n} c_1(\mathcal{G}_l) c_1(\mathcal{G}_m) + \sum_{l=1}^n c_2(\mathcal{G}_l) \right),$$

where the \mathcal{G}_l are non-zero torsion-free L_{ε} -stable sheaves. Moreover, by (26) we have

(28)
$$L^2_{\varepsilon} \cdot c_1(\mathcal{G}_l) \ge 0 \qquad \forall \ l \in \{1, \dots, n\}.$$

Since we renamed according to the lexicographic order, the sequence of slopes with respect to L_{ε} is (not necessarily strictly) decreasing. Thus, setting $r_l := \operatorname{rk} \mathcal{G}_l$ and

$$\alpha_l := \frac{L_{\varepsilon}^2 \cdot c_1(\mathcal{G}_l)}{r_l L_{\varepsilon}^3},$$

we obtain

(29)
$$\alpha_1 \ge \alpha_2 \ge \cdots \ge \alpha_n \ge 0.$$

 $4^{\,e}\,S\acute{E}RIE-TOME\,49-2016-N^o\,4$

By Theorem 7.1 we have $L_{\varepsilon} \cdot c_2(\mathcal{G}_l) \ge \left(\frac{r_l-1}{2r_l}\right) L_{\varepsilon} \cdot c_1^2(\mathcal{G}_l)$ for all $l \in \{1, \ldots, n\}$. Therefore using (27) we obtain

$$L_{\varepsilon} \cdot c_2(\mathcal{F}) = L_{\varepsilon} \cdot \left(\frac{1}{2}c_1(\mathcal{F})^2 + \sum_{l=1}^n c_2(\mathcal{G}_l) - \frac{1}{2}\sum_{l=1}^n c_1(\mathcal{G}_l)^2\right)$$
$$\geq L_{\varepsilon} \cdot \left(\frac{1}{2}c_1(\mathcal{F})^2 - \frac{1}{2r_l}\sum_{l=1}^n c_1(\mathcal{G}_l)^2\right).$$

By Lemma 7.2 we have

$$(L_{\varepsilon}^{2} \cdot c_{1}(\mathcal{G}_{l}))^{2} \geq (L_{\varepsilon} \cdot c_{1}^{2}(\mathcal{G}_{l})) \cdot L_{\varepsilon}^{3} \qquad \forall l \in \{1, \dots, n\},$$

so, using the coefficient α_l defined above, we get $L_{\varepsilon} \cdot c_1^2(\mathcal{G}_i) \leq \alpha_i^2 r_i^2 L_{\varepsilon}^3$. Putting this into the last inequality yields

(30)
$$L_{\varepsilon} \cdot c_2(\mathcal{F}) \ge L_{\varepsilon} \cdot \left(\frac{1}{2}c_1(\mathcal{F})^2 - \frac{1}{2}\sum_{l=1}^n \alpha_l^2 r_l L_{\varepsilon}^2\right)$$

(31)
$$= \frac{1}{2}L_{\varepsilon} \cdot \left((c_1(\mathscr{F})^2 - L_{\varepsilon}^2) + (1 - \sum_{l=1}^n \alpha_l^2 r_l) L_{\varepsilon}^2 \right)$$

We claim that

$$(1-\sum_{l=1}^n \alpha_l^2 r_l) \ge 0.$$

Assuming this for the time being, let us see how to conclude : since $L_{\varepsilon}^3 > 0$, the *claim* together with (31) yields

$$L_{\varepsilon} \cdot c_2(\mathcal{F}) \ge \frac{1}{2} L_{\varepsilon} \cdot \left(c_1(\mathcal{F})^2 - L_{\varepsilon}^2 \right)$$

Now we take the limit $\varepsilon \to 0$.

Proof of the claim. - First of all, (29) implies that

$$1 - \sum_{l=1}^{n} \alpha_l^2 r_l \ge 1 - \alpha_1 \sum_{l=1}^{n} r_l \alpha_l.$$

Now by definition of the numbers α_l we have

$$\sum_{l=1}^n r_l \alpha_l = \sum_{l=1}^n \frac{L_{\varepsilon}^2 \cdot c_1(\mathcal{G}_l)}{L_{\varepsilon}^3} = \frac{L_{\varepsilon}^2 \cdot c_1(\mathcal{F})}{L_{\varepsilon}^3}.$$

Yet by definition of L and L_{ε} ,

$$c_1(\mathcal{F}) = L - P = L_{\varepsilon} - P - \varepsilon \omega.$$

Since *P* is pseudoeffective and since ω is Kähler it follows that $L^2_{\varepsilon} \cdot c_1(\mathscr{G}) \leq L^3_{\varepsilon}$, hence $\sum_{l=1}^n r_l \alpha_l \leq 1$. Since $r_l \geq 1$ and $\alpha_l \geq 0$ for all $l \in \{1, \ldots, n\}$ this implies $\alpha_1 \leq 1$, proving the *claim*.

8. Abundance

In this section, we establish the abundance theorem for non-algebraic Kähler threefolds. The main difficulty is to show that if the numerical Kodaira dimension $\nu(X) = 1$ or $\nu(X) = 2$, then $\kappa(X) \ge 1$. This will be done in Theorems 8.1 and 8.2.

8.A. The case $\nu = 1$

THEOREM 8.1. – Let X be a normal Q-factorial compact Kähler threefold with at most terminal singularities such that K_X is nef. If $\nu(X) = 1$, then $\kappa(X) = 1$.

Proof. – Since $0 \le \kappa(X) \le \nu(X) = 1$, it is sufficient to prove that $\kappa(X) \ge 1$. By Lemma 6.1 and Lemma 6.6 we are reduced to proving the following statement:

Let X be a normal \mathbb{Q} -factorial compact Kähler threefold such that (X,0) is klt, carrying a divisor $D \in |mK_X|$ with the following properties:

- a) Set B := Supp D. The pair (X, B) is lc and $X \setminus B$ has terminal singularities.
- b) The divisor $K_X + B$ is nef and we have $\nu(K_X + B) = 1$.
- c) There exists an irreducible component $S \subset B$ that is a connected component.

Then $\kappa(X) \geq 1$.

We follow the argument in [49, Ch. 13]: by adjunction [49, 16.9.1] there exists a boundary divisor Δ on S such that (S, Δ) is slc and $K_S + \Delta$ is numerically trivial. By Proposition 5.1 the divisor $K_S + \Delta$ is torsion. Due to a covering trick of Miyaoka [49, 11.3.6] we may suppose after a finite cover, étale in codimension one, that

$$\omega_S = \mathcal{O}_S(K_S) \simeq \mathcal{O}_S \simeq \mathcal{O}_S(S).$$

Note that by [51, Prop. 5.20(4)] the klt property is preserved under a finite morphism which is étale in codimension one. In particular, X is Cohen-Macaulay. By [49, 11.3.7] we are finished if we prove that for every infinitesimal neighborhood S_n , the restriction

$$H^p(S_n, \mathcal{O}_{S_n}) \to H^p(S, \mathcal{O}_S)$$

is surjective for every $p \in \mathbb{N}$. Observe here that condition (3) in [49, 11.3.7] is satisfied by [49, 12.1.2], as explained in [49, p.158]. In fact, using the commutative diagram

and the isomorphism $H^p(S_n, \mathbb{C}) \simeq H^p(S, \mathbb{C})$ we see that it is sufficient to prove that

$$H^p(S,\mathbb{C}) \to H^p(S,\mathcal{O}_S)$$

is surjective. This is done in Theorem 5.1.

4° SÉRIE - TOME 49 - 2016 - Nº 4

8.B. The case $\nu = 2$

THEOREM 8.2. – Let X be a normal Q-factorial compact Kähler threefold with at most terminal singularities such that K_X is nef. If $\nu(X) = 2$, then $\kappa(X) \ge 1$.

The basic idea of the proof is the same as in the projective case [49, Sect. 14], however the computations get considerably simplified by our generalization of Miyaoka's Chern class inequality [53, Thm. 6.1]. Let us start by recalling the Riemann-Roch formula for terminal threefolds:

PROPOSITION 8.1. – Let X be a normal compact Kähler threefold with at most terminal singularities, and let L be a line bundle on X. Then we have

$$\chi(X,L) = \frac{L^3}{6} - \frac{1}{4}K_X \cdot L^2 + L \cdot \frac{K_X^2 + c_2(X)}{12} + \chi(X,\mathcal{O}_X),$$

where $L \cdot c_2(X) := \pi^* L \cdot c_2(\hat{X})$ with $\pi : \hat{X} \to X$ any resolution of singularities (cf. Definition 7.1).

Proof. – Let $\pi : \hat{X} \to X$ be a resolution of singularities, which is an isomorphism over the smooth locus of X. Since X has rational singularities, we have $\chi(X, L) = \chi(\hat{X}, \pi^*L)$. Riemann-Roch on the smooth Kähler threefold \hat{X} yields

(32)
$$\chi(\hat{X}, \pi^*L) = \frac{\pi^*L^3}{6} - \frac{1}{4}K_{\hat{X}} \cdot (\pi^*L)^2 + \pi^*L \cdot \frac{K_{\hat{X}}^2 + c_2(X)}{12} + \chi(\hat{X}, \theta_{\hat{X}}).$$

Now $\pi^* L^3 = L^3$, and, using again the rationality of the singularities of X, $\chi(\hat{X}, \theta_{\hat{X}}) = \chi(X, \theta_X)$. Since X is smooth in codimension two we may write

$$K_{\hat{X}} \sim_{\mathbb{Q}} \pi^* K_X + E$$

with E a divisor such that $\pi(E)$ is finite. In particular the projection formula gives $K_{\hat{X}} \cdot (\pi^*L)^2 = K_X \cdot L^2$ and $\pi^*L \cdot K_{\hat{X}}^2 = L \cdot K_X^2$. Thus (32) gives our claim.

We will also need the following Kähler version of Miyaoka's generic nefness theorem, due to Enoki in the smooth case:

PROPOSITION 8.2. – Let X be a normal compact Kähler space of dimension n with canonical singularities. Suppose that K_X is nef or $\kappa(X) \ge 0$. Then Ω_X is generically nef with respect to any nef class α , i.e., for every torsion-free quotient sheaf

$$\Omega_X \to \mathcal{Q} \to 0,$$

we have $\alpha^{n-1} \cdot c_1(Q) \ge 0$.

Proof. – Fix a nef class $\alpha \in N^1(X)$, and let $\Omega_X \to Q \to 0$ be a torsion-free quotient sheaf. Let $\pi : \hat{X} \to X$ be a desingularization by a compact Kähler manifold, and let \mathcal{K} be the kernel of the induced epimorphism

$$(\pi^*\Omega_X)/\text{torsion} \to (\pi^*Q)/\text{torsion} \to 0.$$

Using the injective map

$$(\pi^*\Omega_X)/\text{torsion} \hookrightarrow \Omega_{\hat{X}}$$

we may view \mathcal{K} as a subsheaf of $\Omega_{\hat{X}}$, and we denote by $\hat{\mathcal{K}}$ its saturation in $\Omega_{\hat{X}}$. Set

$$\hat{Q} := \Omega_{\hat{X}} / \hat{\mathcal{K}}_{\hat{X}}$$

then \hat{Q} is a torsion-free quotient of $\Omega_{\hat{X}}$ coinciding with $(\pi^* Q)/\text{torsion}$ in the complement of the exceptional locus. In particular we have

$$\alpha^{n-1} \cdot c_1(Q) = (\pi^* \alpha)^{n-1} \cdot c_1((\pi^* Q) / \text{torsion}) = (\pi^* \alpha)^{n-1} \cdot c_1(\hat{Q}).$$

We will now verify the conditions of [25, Thm. 1.4] in order to conclude that $(\pi^* \alpha)^{n-1} \cdot c_1(\hat{Q}) \ge 0$: since X has canonical singularities we have

$$K_{\hat{X}} = \pi^* K_X + E,$$

with *E* an effective, π -exceptional \mathbb{Q} -divisor.

Thus if K_X is nef the conditions of [25, Thm. 1.4] hold by setting $L = \pi^* K_X$ and D = E. If $\kappa(X) \ge 0$ we have $\pi^* K_X = F$ with F an effective Q-divisor, so the conditions are satisfied by setting L = 0 and D = F + E. In both cases Enoki's theorem tells us that

$$\omega^{n-1} \cdot c_1(\hat{Q}) \ge 0$$

for every Kähler form ω on \hat{X} . Since $\pi^* \alpha$ is nef, the statement follows by passing to the limit.

We will use the following Serre vanishing property:

LEMMA 8.1. – Let X be a normal Q-factorial compact Kähler threefold, and let B_1, \ldots, B_k be prime Weil divisor on X such that B_i is Cohen-Macaulay for every $i \in \{1, \ldots, k\}$. Let L be a nef Cartier divisor on X such that $L_{|B_i|} \not\equiv 0$ for every $i \in \{1, \ldots, k\}$. Let $Y \subset X$ be a subscheme such that $Y_{red} \subset \sum_{i=1}^k B_i$ and let \mathcal{F} be a coherent sheaf on Y. Then there exists a number $n_0 \in \mathbb{N}$ such that

$$H^2(Y, \mathcal{F} \otimes \mathcal{O}_Y(L^{\otimes n})) = 0$$

for every $n \ge n_0$.

Proof. – Note first that we may suppose that Y is defined by an ideal sheaf $\mathcal{O}_X(-\sum_{i=1}^k a_i B_i)$ with $a_i \in \mathbb{N}$. Indeed, at the general point of every surface B_i , the scheme Y is isomorphic to a scheme $\sum_{i=1}^k a_i B_i$ defined by the ideal sheaf $\mathcal{O}_X(-\sum_{i=1}^k a_i B_i)$. Thus if we consider the restriction map

$$\mathcal{F} \twoheadrightarrow \mathcal{F} \otimes \mathcal{O}_{\sum_{i=1}^{k} a_i B_i};$$

its kernel has support on a scheme of dimension at most one. Hence

$$H^{2}(Y, \mathcal{F} \otimes \mathcal{O}_{Y}(L^{\otimes n})) \simeq H^{2}(\sum_{i=1}^{k} a_{i}B_{i}, \mathcal{F} \otimes \mathcal{O}_{\sum_{i=1}^{k} a_{i}B_{i}}(L^{\otimes n}))$$

for every $n \in \mathbb{N}$. We will now argue by induction on $\sum_{i=1}^{k} a_i$. The start of the induction is the case where \mathcal{F} is a coherent sheaf on one of the surfaces B_i . Since B_i is Cohen-Macaulay, Serre duality gives

$$H^{2}(B_{i}, \mathcal{F} \otimes \mathcal{O}_{B_{i}}(L^{\otimes n})) \simeq \operatorname{Hom}(\mathcal{F} \otimes \mathcal{O}_{B_{i}}(L^{\otimes n}), \omega_{B_{i}}).$$

4° SÉRIE – TOME 49 – 2016 – Nº 4

Let $\nu : \tilde{B}_i \to B_i$ denote the normalization. Since $\mathcal{H}om(\mathcal{F}, \omega_{B_i})$ is torsion free, the natural map

$$\mathcal{H}om(\mathcal{F},\omega_{B_i}) \to \nu_*(\nu^*(\mathcal{H}om(\mathcal{F},\omega_{B_i})))$$

is injective. Since $\nu_*(\nu^*(\mathcal{H}om(\mathcal{F},\omega_{B_i})))$ and $\nu_*(\nu^*(\mathcal{H}om(\mathcal{F},\omega_{B_i})/Tor))$ coincide at the generic point of B_i and $\mathcal{H}om(\mathcal{F},\omega_{B_i})$ is torsion free, the map

$$\mathcal{H}om(\mathcal{F},\omega_{B_i}) \to \nu_*(\nu^*(\mathcal{H}om(\mathcal{F},\omega_{B_i})/Tor))$$

is injective, too. Thus it suffices to show that for any torsion-free sheaf \mathscr{A} on \tilde{B}_i one has

$$H^0(\tilde{B}_i, \mathscr{O} \otimes \mathcal{O}_{\tilde{B}_i}(\tilde{L}^{-\otimes n})) = 0$$

for $n \gg 0$ and $\tilde{L} := \nu^*(L_{|B_i})$. Passing to a desingularization, we may assume \tilde{B}_i smooth and φ locally free. Fix now a Kähler form ω on \tilde{B}_i , and let $\mathcal{G}_1 \subset \varphi$ be the first sheaf of the Harder-Narasimhan filtration with respect to ω . Since $L_{|B_i|}$ is a non-zero nef divisor we have $\tilde{L} \cdot \omega > 0$. Thus there exists a number $n_0 \in \mathbb{N}$ such that $\mathcal{G}_1 \otimes \mathcal{O}_{\tilde{B}_i}(\tilde{L}^{-\otimes n})$ has negative slope for all $n \geq n_0$. In particular $\varphi \otimes \mathcal{O}_{B_i}(L^{-\otimes n})$ has no global sections.

For the induction step we simply choose a surface B_j such that $a_j > 0$. Then the kernel of the restriction map

$$\mathcal{F} \twoheadrightarrow \mathcal{F} \otimes \mathcal{O}_{(\sum_{i=1}^{k} a_i B_i) - B_i}$$

is a sheaf \mathcal{G} with support on B_j . Thus by the induction hypothesis we know that for $n \gg 0$ the second cohomology vanishes for both $\mathcal{G} \otimes \mathcal{O}_{B_j}(L^{\otimes n})$ and $\mathcal{F} \otimes \mathcal{O}_{(\sum_{i=1}^k a_i B_i) - B_j}(L^{\otimes n})$. \Box

COROLLARY 8.1. – Let X be a normal Q-factorial compact Kähler threefold, and let L be a nef Cartier divisor on X. Let $D \in |L|$ be effective and set B := Supp D and B_1, \ldots, B_k for the irreducible components of B. Suppose that B_i is Cohen-Macaulay for every $i \in \{1, \ldots, k\}$. Suppose also that $L_{|B_i|} \not\equiv 0$ for every $i \in \{1, \ldots, k\}$. Then there exists a number $n_0 \in \mathbb{N}$ and constants $c_1, c_2 \in \mathbb{N}$ such that for all $n \ge n_0$:

$$\dim H^2(X, L^{\otimes n}) = c_1$$

and

$$\dim H^2(X, L^{\otimes n} \otimes \mathcal{O}_X(-B)) = c_2.$$

Proof. – For all $n \in \mathbb{N}$ we have an exact sequence

$$0 \to L^{\otimes n-1} \to L^{\otimes n} \to \mathcal{O}_D(L^{\otimes n}) \to 0.$$

By Lemma 8.1,

$$H^2(D, \mathcal{O}_D(L^{\otimes n})) = 0$$

for all $n \gg 0$. Thus the map $H^2(X, L^{\otimes n-1}) \to H^2(X, L^{\otimes n})$ is surjective for all $n \gg 0$. This shows the first statement.

For the second statement assume without loss of generality that $D - B \neq 0$. Note that D - B is an effective Weil divisor whose support is contained in B. For all $n \in \mathbb{N}$ we have an exact sequence

$$0 \to L^{\otimes n-1} \to L^{\otimes n-1} \otimes \mathcal{O}_X(D-B) \to \mathcal{O}_{D-B}(L^{\otimes n-1}) \to 0$$

Again by Lemma 8.1

$$H^2(D-B, \mathcal{O}_{D-B}(L^{\otimes n-1})) = 0$$

for all $n \gg 0$. Thus the map

$$H^2(X, L^{\otimes n-1}) \to H^2(X, L^{\otimes n-1} \otimes \mathcal{O}_X(D-B))$$

is surjective for $n \gg 0$ and by the first statement $H^2(X, L^{\otimes n-1})$ is constant for $n \gg 0$. We conclude by noting that

$$L^{\otimes n-1} \otimes \mathcal{O}_X(D-B) \simeq L^{\otimes n} \otimes \mathcal{O}_X(-B)$$

since both sheaves are reflexive and coincide on the smooth locus of X.

Proof of Theorem 8.2. – By Lemma 6.1 and Lemma 6.8 we are reduced to proving the following statement:

Let X be a normal \mathbb{Q} -factorial compact Kähler threefold such that (X,0) is klt, carrying a divisor $D \in |mK_X|$ with the following properties:

- a) Set B := Supp D. The pair (X, B) is lc and $X \setminus B$ has terminal singularities.
- b) The divisor $K_X + B$ is nef and we have $\nu(K_X + B) = 2$. Moreover we have $\kappa(X) = \kappa(K_X + B)$.
- c) For every irreducible component $T \subset B$ we have $(K_X + B)|_T \neq 0$.
- d) We have $(K_X + B) \cdot K_X^2 \ge 0$.

Then $\kappa(X) \geq 1$.

Step 1: Singularities of X. – We claim that there is a finite set $S \subset X$ such that $X \setminus S$ has canonical singularities. By hypothesis $X \setminus B$ has only terminal singularities, which are isolated. Thus it remains to consider the singular points of X which are contained in B. Taking a finite covering of X by analytic neighborhoods we see that it is sufficient to prove the claim for X a Stein variety. Thus we can take a hyperplane section H of X. Now for general H, the pair (H, B_H) is lc by [46, Prop. 7.7], so by [40, Thm. 9.6] every point $p \in B_H \subset H$ is a rational double point in H. Hence by [51, Thm. 5.34], the point p is a canonical singularity of X.

Step 2: A Chern class inequality. – Let $\mu : X' \to X$ be a terminal modification of X (cf. Theorem 2.1). Thus X' has only terminal singularities, and there exists an effective \mathbb{Q} -divisor Δ such that

$$K_{X'} + \Delta \sim_{\mathbb{O}} \mu^* K_X.$$

Let m be the Cartier index of $K_X + B$, then

$$L := m(K_X + B)$$

is Cartier, and we set $L' := \mu^*(L)$. We prove the basic Chern class inequality

(33)
$$L' \cdot (K_{X'}^2 + c_2(X')) \ge 0.$$

In fact, since X has only finitely many non-canonical points, $\mu(\Delta)$ is finite. Therefore the projection formula and our assumption d) yield

(34)
$$L' \cdot K_{X'}^2 = \mu^* L \cdot K_{X'}^2 = m(K_X + B) \cdot K_X^2 \ge 0.$$

By Proposition 8.2 the sheaf $\Omega_{X'}$ is generically nef. Since

$$K_{X'} + \Delta + \mu^* B \sim_{\mathbb{O}} \mu^* (K_X + B)$$

^{4°} SÉRIE – TOME 49 – 2016 – Nº 4

is nef, the conditions of Theorem 7.2 are satisfied for $\mathcal{F} := (\Omega_{X'})^{**}$ and $P := \Delta + \mu^* B$. Having in mind that $L^3 = m^3 (K_X + B)^3 = 0$ and using (34), Theorem 7.2 yields

$$L' \cdot c_2(X') \ge 0,$$

hence the Chern class inequality (33) is established.

Step 3: A Riemann-Roch computation. – Since $K_X + B$ is nef and \mathbb{Q} -linearly equivalent to an effective divisor with support B, the equality $(K_X + B)^3 = 0$ implies that

$$(35) (K_X+B)^2 \cdot T = 0$$

for every irreducible component $T \subset B$. Since K_X is \mathbb{Q} -linearly equivalent to an effective divisor with support B, we conclude

$$(K_X + B)^2 \cdot K_X = 0.$$

Since $\mu(\Delta)$ is finite the projection formula yields

$$K_{X'} \cdot (L')^2 = K_X \cdot L^2 - \Delta \cdot (\mu^* L)^2 = m^2 K_X \cdot (K_X + B)^2 = 0.$$

Thus Proposition 8.1 gives

$$\chi(X', \mathcal{O}_{X'}(nL')) = nL' \cdot \frac{K_{X'}^2 + c_2(X')}{12} + \chi(X', \mathcal{O}_{X'})$$

for all $n \in \mathbb{N}$. Thus (33) yields a constant k such that

 $\chi(X', \mathcal{O}_{X'}(nL')) \ge k$

for all $n \in \mathbb{N}$. Since X has rational singularities, we conclude that

$$\chi(X, \mathcal{O}_X(nL)) \ge k$$

for all $n \in \mathbb{N}$. Since $H^3(X, nL) = 0$ for $n \gg 0$ and dim $H^2(X, nL)$ is constant for $n \gg 0$ by Corollary 8.1 (note that the components of *B* are Cohen-Macaulay by [51, 5.25]), we arrive at

(36)
$$h^0(X, nL) \ge h^1(X, nL) + c$$

with some constant $c \in \mathbb{Z}$.

Step 4. A simple case. – Although not really necessary, it is instructive to give the simple concluding argument in the case of strict inequality in (33). Then the preceding computation yields that

$$h^0(X, nL) \ge h^1(X, nL) + Dn$$

with some positive constant D. Thus

$$\kappa(K_X + B) = \kappa(X, L) > 0,$$

so $\kappa(X) \ge 1$ by assumption b).

Step 5: Conclusion. – By (36) it suffices to show that $h^1(X, nL)$ grows at least linearly. We consider the exact sequence

(37)
$$0 \to \mathcal{O}_X(nL - B) \to \mathcal{O}_X(nL) \to \mathcal{O}_B(nL) \to 0.$$

By Corollary 8.1 we know that $h^2(X, \mathcal{O}_X(nL - B))$ is constant for $n \gg 0$. Taking cohomology of the exact sequence (37), it remains to show that $h^1(B, \mathcal{O}_B(nL))$ grows at least linearly. To this extent, we will prove that $\chi(B, \mathcal{O}_B(nL))$ is constant. Assuming this for the time being, let us see how to conclude: by Lemma 8.1 we have $H^2(B, \mathcal{O}_B(nL)) = 0$ for $n \gg 0$. Moreover by adjunction [49, 16.9.1] $\mathcal{O}_B(K_B) \simeq \mathcal{O}_B(K_X + B)$, so by Proposition 5.2

 $h^0(B, \mathcal{O}_B(nL)) = h^0(B, \mathcal{O}_B(nm(K_X + B))) = h^0(B, \mathcal{O}_B(nmK_B))$

grows linearly. Thus $h^1(B, \mathcal{O}_B(nL))$ grows linearly.

Proof of the claim. – By [52, Thm. 3.1] the Euler characteristic $\chi(B, \Theta_B(nL))$ on the slc surface *B* is computed by the usual Riemann-Roch formula⁽¹²⁾

$$\chi(B, \mathcal{O}_B(nL)) = \chi(B, \mathcal{O}_B) + \frac{1}{2}(nL_{|_B}) \cdot (nL_{|_B} - K_B).$$

Yet by (35) we have $0 = L^2 \cdot B = (L_{|_B})^2$. Since $\mathcal{O}_B(K_B) \simeq \mathcal{O}_B(K_X + B)$ is a multiple of $L_{|_B}$, this also implies that $L_{|_B} \cdot K_B = 0$.

8.C. Proof of Theorem 1.1

Proof. – By [24, Thm. 0.3] we have $\kappa(X) \ge 0$. If $\nu(X) = 3$, then K_X is big, hence X is Moishezon and therefore projective [57]. Thus the result follows from the base point free theorem. If $\kappa(X) = \nu(X) \le 2$ the statement follows from Kawamata's theorem [39, Thm. 1.1], [56, Thm. 5.5], [29, Sect. 4].

By Theorem 8.1 and Theorem 8.2 we are thus left to exclude the possibility that $\kappa(X) = 1$ and $\nu(X) = 2$. This is done exactly as in [39, Thm. 7.3].

9. Applications

In this concluding section we apply the MMP to explore the structure of non-algebraic compact Kähler threefolds X with Kodaira dimension $\kappa(X) \leq 0$.

9.A. Uniruled threefolds

THEOREM 9.1. – Let X be a smooth non-algebraic compact Kähler threefold with $\kappa(X) = -\infty$. Then X is bimeromorphic to a normal compact Kähler threefold X' with at most terminal singularities with the following properties. There exists a contraction

$$\varphi: X' \to Y$$

of an extremal ray in $\overline{NA}(X')$ such that

- (1) Y is a normal non-algebraic Kähler surface with only rational singularities.
- (2) There is a finite set $A \subset Y$ such that $\varphi | X' \setminus \varphi^{-1}(A) \to Y \setminus A$ is a conic bundle.

 $^{(12)}$ We refer to [52, Ch. 2.3] for the definition of the intersection product on the non-normal surface B.

4° SÉRIE - TOME 49 - 2016 - Nº 4

(3) φ realizes the MRC fibration of X', and $\kappa(\hat{Y}) \in \{0,1\}$ for any desingularization $\hat{Y} \to Y$.

Proof. – By [38, Cor. 1.4] the manifold X is uniruled. We first claim that "the" base B, chosen smooth, of the MRC-fibration $X \rightarrow B$ has dimension two. Indeed, if dim B = 1, then the MRC fibration is realized by a morphism $X \rightarrow B$ and the general fibre F is rationally connected, hence $H^2(F, \Theta_F) = 0$. This immediately implies $H^2(X, \Theta_X) = 0$, so X is projective by Kodaira's criterion. The same of course applies when dim B = 0 which is to say that X is rationally connected. Notice also that B is non-algebraic, otherwise X were algebraic.

By [37, Thm. 1.1] we can run a MMP which terminates with a Mori fibre space $\varphi : X' \to Y$. Moreover the properties (1) and (2) are shown in [37, Thm. 1.1] and [37, Rem.4.2]. We have seen above that the base of the MRC fibration has dimension two, so φ realizes this fibration for X'. In particular Y is not algebraic, so any desingularization \hat{Y} is not uniruled. Thus we have $\kappa(\hat{Y}) \ge 0$. The non-algebraicity of Y also yields $\kappa(\hat{Y}) \le 1$.

We describe the structure of X resp. X' more closely. If the algebraic dimension a(Y) = 1, we denote by $f: Y \to C$ the algebraic reduction.

COROLLARY 9.1. – In the setting of Theorem 9.1, the algebraic dimensions (a(X), a(Y)) can take the following values.

- (1) (0,0), and Y is bimeromorphic to a K3-surface or a torus.
- (2) (1,0), and X is bimeromorphic (up to an étale quotient possibly) of a product of a K3-surface or a tori with an elliptic curve.
- (3) (1,1), and $f \circ \varphi$ is the algebraic reduction of X'; the general fiber $f \circ \varphi$ being a ruled surface $\mathbb{P}(V)$ over a (possibly varying) elliptic curve E with $V = \Theta_E \oplus L$ and $L \equiv 0$ but not torsion or V the non-split extension of two trivial line bundles.
- (4) (2,1), and $X' \simeq Y \times \mathbb{P}_1$, possibly after a base change $\tilde{C} \to C$.

Proof. – Assertion (1), (2) and (3) are contained in [31, 14.1], [16, 9.1]. As to (4), since it is shown in [16, 9.1] that a more precise structure property holds: after passing to \tilde{X} and \tilde{Y} via a finite (ramified) cover $\tilde{C} \to C$, we obtain

$$\tilde{X} \simeq Z \times_{\tilde{C}} \tilde{Y},$$

where $Z \to \tilde{C}$ is bimeromorphic to $\mathbb{P}_1 \times \tilde{C}$, hence the claim.

EXAMPLE 9.1. – The cases (2) and (4) can obviously be realized by products, we give examples for the other cases.

a) Let Y be a K3 surface with Pic(Y) = 0 and set $X = \mathbb{P}(T_Y)$. Then X does not contain any divisor, hence a(X) = a(Y) = 0.

b) Let Y be a two-dimensional torus of algebraic dimension 1, and denote by F the general fibre of the algebraic reduction $f : Y \to C$. Let $\mathcal{L} \in \operatorname{Pic}(Y)$ be a line bundle such that $\mathcal{L}_{|_F}$ is numerically trivial but not torsion, and set $X := \mathbb{P}(\mathcal{O}_Y \oplus \mathcal{L})$. We claim that a(X) = a(Y) = 1: it suffices to prove that $\kappa(X, \mathcal{G}) \leq 1$ for any line bundle \mathcal{G} on X. Now any line bundle \mathcal{G} on X is of the form

$$\mathcal{G} = \mathcal{O}_{\mathbb{P}(\mathcal{O}_Y \oplus \mathcal{L})}(k) \otimes \pi^*(\mathcal{F})$$

ANNALES SCIENTIFIQUES DE L'ÉCOLE NORMALE SUPÉRIEURE

with a line bundle \mathcal{F} on Y. If $\kappa(\mathcal{G}) \geq 1$, then $k \geq 0$ and

$$h^{0}(X, \mathcal{G}^{\otimes m}) = h^{0}(Y, S^{km}(\mathcal{O}_{Y} \oplus \mathcal{L}) \otimes \mathcal{F}^{m}) = \sum_{l=0}^{km} h^{0}(Y, \mathcal{L}^{l} \otimes \mathcal{F}^{m}).$$

Since f is the algebraic reduction, the line bundle \mathscr{T} has degree 0 on F. Since $\mathscr{L}|_F$ is not torsion there exists a unique $l_0 \in \{0, \ldots, km\}$ such that $(\mathscr{L}^l \otimes \mathscr{T}^m)|_F \simeq \mathscr{O}_F$. Thus we have

$$h^0(X, \mathcal{G}^{\otimes m}) = h^0(Y, \mathcal{L}^{l_0} \otimes \mathcal{F}^m).$$

We immediately obtain $\kappa(X, \mathcal{G}) \leq 1$.

9.B. Threefolds with trivial canonical bundles

We will study the Albanese map for certain threefolds with terminal singularities (cf. [9, 2.4.1] for the existence of Albanese maps in the presence of rational singularities).

LEMMA 9.1. – Let X be a (non-algebraic) compact Kähler threefold with terminal singularities. If $\kappa(X) = 0$, then the Albanese map $\alpha : X \to A = Alb(X)$ is surjective with connected fibres. In particular we have $q(X) \leq 3$.

Proof. – Apply [64, Main Thm.I] to a desingularization $\hat{X} \to X$.

THEOREM 9.2. – Let X be a non-algebraic compact Kähler threefold with terminal singularities. If $K_X \equiv 0$, there exists a Galois cover $f : \tilde{X} \to X$ that is étale in codimension one such that either \tilde{X} is a torus or a product of an elliptic curve and a K3 surface.

Proof. – By Theorem 1.1, the divisor K_X is semi-ample. Thus we can choose $m \in \mathbb{N}$ minimal such that $\mathcal{O}_X(mK_X) = \mathcal{O}_X$. Let $X' \to X$ be the induced cyclic cover, then we have $\mathcal{O}_{X'}(K_{X'}) \simeq \mathcal{O}_{X'}$. In particular X' is Gorenstein with terminal singularities [51, Cor. 5.21]. Thus the Riemann-Roch Formula 8.1 gives $\chi(X', \mathcal{O}_{X'}) = 0$. Since $h^2(X', \mathcal{O}_{X'}) \geq 1$, the variety X' being non-algebraic with rational singularities only, we conclude that $h^1(X', \mathcal{O}_X) \geq 1$.

Now consider the Albanese map $\alpha : X' \to Alb(X') =: A$. By Lemma 9.1 the morphism α is surjective with connected fibres.

Case 1: q(X') = 3. – Then α is bimeromorphic. Since A is smooth, $K_{X'} \equiv E$ with E an effective divisor whose support equals the α -exceptional locus. Since $K_{X'} \equiv 0$ we conclude that $X' \simeq A$.

Case 2: q(X') = 2. – By [64, Main Thm.I,2)] there exists a finite set $S \subset A$ such that α is an elliptic bundle over $A \setminus S$. By [16, Prop. 6.7(i)] this implies that (after finite étale base change) the fibration α is bimeromorphic to a compact Kähler manifold X'' that is an elliptic bundle $X'' \to A$. Yet a Kähler manifold which is an elliptic bundle over a torus is an étale quotient of a torus [8], so after finite étale cover we have q(X') = q(X'') = 3. We conclude by applying Case 1.

Case 3: q(X') = 1. – By [64, Main Thm.I,3)] there exists an analytic fibre bundle $X^* \to A$ that is bimeromorphic to X'. Since the Kodaira dimension is an invariant of varieties with terminal singularities, we have $\kappa(X^*) = 0$. In particular the general fibre F of $X^* \to A$ is the blow-up of a torus or a K3 surface⁽¹³⁾. We next run a relative MMP over the elliptic curve A. Since $X^* \to A$ is a fibre bundle, every step of this MMP is the blow-up along an étale multisection of the fibration, so the outcome is an analytic fibre bundle $\bar{X} \to A$ such that the general fibre F is a torus or K3 surface. If F is covered by a torus, then \bar{X} is a torus after finite étale cover, see e.g., [8]. If F is a K3 surface, the fibre bundle trivializes after finite étale base change $A' \to A$ ([30, Cor. 4.10], cf. [18, Lemma 2.15] for more details). Thus we have $\bar{X} \simeq A' \times F$. In conclusion we see that (up to finite étale cover) we have a bimeromorphic map

$$\mu: X' \dashrightarrow \bar{X}$$

with \bar{X} a torus or a product $A' \times F$. Since both $K_{X'}$ and $K_{\bar{X}}$ are numerically trivial, we see that μ is an isomorphism in codimension one [35], [45, 4.3]. Moreover μ decomposes into a finite sequence of flops by [45, 4.9]. Note however that the last flop of this sequence yields a rational curve in \bar{X} that is very rigid (in the sense of [38, Defn. 4.3]). Since \bar{X} is a torus or a product $A' \times F$ such a curve does not exist on \bar{X} . Thus μ is an isomorphism.

Using the existence of minimal models for a smooth compact Kähler threefold, we deduce

COROLLARY 9.2. – Let X be a non-algebraic compact Kähler threefold with $\kappa(X) = 0$. There exists a finite cover which is bimeromorphic to a torus or a product of an elliptic curve and a K3 surface.

It was known since some time that Theorem 1.2 is a consequence of the existence of good minimal models [59, p.731]. We will now derive Theorem 1.2 from the more general Theorem 9.2:

Proof of Theorem 1.2. – Since X is not uniruled, K_X is pseudo-effective [15]. By [38, Thm. 1.1] there exists a minimal model $X \to X'$. Since X has algebraic dimension zero, we see that $\kappa(X) = 0$. Thus we have $\nu(X') = \kappa(X') = 0$ by Theorem 1.1, i.e., the canonical divisor $K_{X'}$ is numerically trivial. Since X (and hence X') is not covered by curves Theorem 9.2 yields that $X' \simeq T/G$ with T a torus and G a finite group. Since X (and hence X') is not covered by positive-dimensional subvarieties, the torus T has no positivedimensional subvarieties. In particular T/G has no positive-dimensional subvarieties, so $X \to T/G$ extends to a morphism.

BIBLIOGRAPHY

 V. ANCONA, Faisceaux amples sur les espaces analytiques, *Trans. Amer. Math. Soc.* 274 (1982), 89–100.

⁽¹³⁾ If F were bimeromorphic to an Enriques or bielliptic surface we would have $0 = H^2(X^*, \theta_{X^*}) = H^2(X', \theta_{X'})$, a contradiction.

- [2] V. ANCONA, Vanishing and nonvanishing theorems for numerically effective line bundles on complex spaces, Ann. Mat. Pura Appl. 149 (1987), 153–164.
- [3] V. ANCONA, V. V. TAN, On the blowing down problem in C-analytic geometry, J. reine angew. Math. 350 (1984), 178–182.
- [4] C. ARAUJO, The cone of pseudo-effective divisors of log varieties after Batyrev, *Math. Z.* 264 (2010), 179–193.
- [5] S. BANDO, Y.-T. SIU, Stable sheaves and Einstein-Hermitian metrics, in *Geometry and analysis on complex manifolds*, World Sci. Publ., River Edge, NJ, 1994, 39–50.
- [6] W. P. BARTH, K. HULEK, C. A. M. PETERS, A. VAN DE VEN, Compact complex surfaces, Ergebn. Math. Grenzg. 4, Springer, 2004.
- [7] T. BAUER, F. CAMPANA, T. ECKL, S. KEBEKUS, T. PETERNELL, S. RAMS, T. SZEM-BERG, L. WOTZLAW, A reduction map for nef line bundles, in *Complex geometry* (*Göttingen, 2000*), Springer, Berlin, 2002, 27–36.
- [8] A. BEAUVILLE, Variétés kählériennes dont la première classe de Chern est nulle, J. Differential Geom. 18 (1983), 755–782.
- [9] M. C. BELTRAMETTI, A. J. SOMMESE, The adjunction theory of complex projective varieties, de Gruyter Expositions in Mathematics 16, Walter de Gruyter & Co., Berlin, 1995.
- [10] C. BIRKAR, P. CASCINI, C. D. HACON, J. MCKERNAN, Existence of minimal models for varieties of log general type, J. Amer. Math. Soc. 23 (2010), 405–468.
- [11] S. BOUCKSOM, Cônes positifs des variétés complexes compactes, thèse de doctorat, Université de Grenoble, 2002.
- [12] S. BOUCKSOM, Divisorial Zariski decompositions on compact complex manifolds, Ann. Sci. École Norm. Sup. 37 (2004), 45–76.
- [13] S. BOUCKSOM, J.-P. DEMAILLY, M. PĂUN, T. PETERNELL, The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension, J. Algebraic Geom. 22 (2013), 201–248.
- [14] S. BOUCKSOM, P. EYSSIDIEUX, V. GUEDJ (eds.), An introduction to the Kähler-Ricci flow, Lecture Notes in Math. 2086, Springer, Cham, 2013.
- [15] M. BRUNELLA, A positivity property for foliations on compact Kähler manifolds, Internat. J. Math. 17 (2006), 35–43.
- [16] F. CAMPANA, T. PETERNELL, Complex threefolds with non-trivial holomorphic 2-forms, J. Algebraic Geom. 9 (2000), 223–264.
- [17] J. CAO, A remark on compact Kähler manifolds with nef anticanonical bundles and its applications, preprint arXiv:1305.4397.
- [18] B. CLAUDON, A. HÖRING, Compact Kähler manifolds with compactifiable universal cover, with an appendix by Frédéric Campana, *Bull. Soc. Math. France* 141 (2013), 355–375.
- [19] T. C. COLLINS, V. TOSATTI, Kähler currents and null loci, *Invent. math.* 202 (2015), 1167–1198.
- [20] A. CORTI (ed.), Flips for 3-folds and 4-folds, Oxford Lecture Series in Mathematics and its Applications 35, Oxford Univ. Press, Oxford, 2007.

 $4^{\,e}\,\,S\acute{E}RIE-TOME\,\,49-2016-N^o\,\,4$

- [21] O. DEBARRE, *Higher-dimensional algebraic geometry*, Universitext, Springer, New York, 2001.
- [22] J.-P. DEMAILLY, C. D. HACON, M. PĂUN, Extension theorems, non-vanishing and the existence of good minimal models, *Acta Math.* 210 (2013), 203–259.
- [23] J.-P. DEMAILLY, M. PĂUN, Numerical characterization of the Kähler cone of a compact Kähler manifold, Ann. of Math. 159 (2004), 1247–1274.
- [24] J.-P. DEMAILLY, T. PETERNELL, A Kawamata-Viehweg vanishing theorem on compact Kähler manifolds, J. Differential Geom. 63 (2003), 231–277.
- [25] I. ENOKI, Stability and negativity for tangent sheaves of minimal Kähler spaces, in *Geometry and analysis on manifolds (Katata/Kyoto, 1987)*, Lecture Notes in Math. 1339, Springer, Berlin, 1988, 118–126.
- [26] A. FUJIKI, On the structure of compact complex manifolds in C, in Algebraic varieties and analytic varieties (Tokyo, 1981), Adv. Stud. Pure Math. 1, North-Holland, Amsterdam, 1983, 231–302.
- [27] O. FUJINO, Special termination and reduction to pl flips, in *Flips for 3-folds and 4-folds*, Oxford Lecture Ser. Math. Appl. 35, Oxford Univ. Press, Oxford, 2007, 63–75.
- [28] O. FUJINO, Non-vanishing theorem for log canonical pairs, J. Algebraic Geom. 20 (2011), 771–783.
- [29] O. FUJINO, On Kawamata's theorem, in *Classification of algebraic varieties*, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2011, 305–315.
- [30] T. FUJITA, On Kähler fiber spaces over curves, J. Math. Soc. Japan 30 (1978), 779–794.
- [31] T. FUJITA, On polarized manifolds whose adjoint bundles are not semipositive, in *Algebraic geometry, Sendai, 1985*, Adv. Stud. Pure Math. 10, North-Holland, Amsterdam, 1987, 167–178.
- [32] W. FULTON, Intersection theory, second ed., Ergebn. Math. Grenzg. 2, Springer, Berlin, 1998.
- [33] H. GRAUERT, Über Modifikationen und exzeptionelle analytische Mengen, Math. Ann. 146 (1962), 331–368.
- [34] J. GRIVAUX, Chern classes in Deligne cohomology for coherent analytic sheaves, Math. Ann. 347 (2010), 249–284.
- [35] M. HANAMURA, On the birational automorphism groups of algebraic varieties, *Compositio Math.* 63 (1987), 123–142.
- [36] R. HARTSHORNE, *Algebraic geometry*, Graduate Texts in Math. 52, Springer, New York-Heidelberg, 1977.
- [37] A. HÖRING, T. PETERNELL, Mori fibre spaces for Kähler threefolds, J. Math. Sci. Univ. Tokyo 22 (2015), 219–246.
- [38] A. HÖRING, T. PETERNELL, Minimal models for Kähler threefolds, *Invent. math.* 203 (2016), 217–264.
- [39] Y. KAWAMATA, Pluricanonical systems on minimal algebraic varieties, *Invent. math.* 79 (1985), 567–588.
- [40] Y. KAWAMATA, Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces, Ann. of Math. 127 (1988), 93–163.

- [41] Y. KAWAMATA, On the length of an extremal rational curve, *Invent. math.* 105 (1991), 609–611.
- [42] Y. KAWAMATA, Abundance theorem for minimal threefolds, *Invent. math.* 108 (1992), 229–246.
- [43] Y. KAWAMATA, Termination of log flips for algebraic 3-folds, Internat. J. Math. 3 (1992), 653–659.
- [44] S. KOBAYASHI, Differential geometry of complex vector bundles, Publications of the Mathematical Society of Japan 15, Princeton Univ. Press, Princeton, NJ; Iwanami Shoten, Tokyo, 1987.
- [45] J. KOLLÁR, Flops, Nagoya Math. J. 113 (1989), 15-36.
- [46] J. KOLLÁR, Singularities of pairs, in *Algebraic geometry—Santa Cruz 1995*, Proc. Sympos. Pure Math. 62, Amer. Math. Soc., Providence, RI, 1997, 221–287.
- [47] J. KOLLÄR, Kodaira's canonical bundle formula and adjunction, in *Flips for 3-folds and 4-folds*, Oxford Lecture Ser. Math. Appl. 35, Oxford Univ. Press, 2007, 134–162.
- [48] J. KOLLÁR, Singularities of the minimal model program, Cambridge Tracts in Mathematics 200, Cambridge Univ. Press, Cambridge, 2013.
- [49] J. KOLLÁR et al., Flips and abundance for algebraic threefolds, Astérisque 211, Soc. Math. France, 1992.
- [50] J. KOLLÁR, S. J. KOVÁCS, Log canonical singularities are Du Bois, J. Amer. Math. Soc. 23 (2010), 791–813.
- [51] J. KOLLÁR, S. MORI, Birational geometry of algebraic varieties, Cambridge Tracts in Mathematics 134, Cambridge Univ. Press, Cambridge, 1998.
- [52] W. LIU, S. ROLLENSKE, Geography of Gorenstein stable log surfaces, Trans. Amer. Math. Soc. 368 (2016), 2563–2588.
- [53] Y. MIYAOKA, The Chern classes and Kodaira dimension of a minimal variety, in *Algebraic geometry, Sendai, 1985*, Adv. Stud. Pure Math. 10, North-Holland, Amsterdam, 1987, 449–476.
- [54] Y. MIYAOKA, Abundance conjecture for 3-folds: case $\nu = 1$, Compositio Math. 68 (1988), 203–220.
- [55] S. MORI, Flip theorem and the existence of minimal models for 3-folds, J. Amer. Math. Soc. 1 (1988), 117–253.
- [56] N. NAKAYAMA, The lower semicontinuity of the plurigenera of complex varieties, in *Algebraic geometry, Sendai, 1985*, Adv. Stud. Pure Math. 10, North-Holland, Amsterdam, 1987, 551–590.
- [57] Y. NAMIKAWA, Projectivity criterion of Moishezon spaces and density of projective symplectic varieties, *Internat. J. Math.* 13 (2002), 125–135.
- [58] M. PĂUN, Sur l'effectivité numérique des images inverses de fibrés en droites, Math. Ann. 310 (1998), 411–421.
- [59] T. PETERNELL, Towards a Mori theory on compact Kähler threefolds. II, Math. Ann. 311 (1998), 729–764.
- [60] T. PETERNELL, Towards a Mori theory on compact Kähler threefolds. III, Bull. Soc. Math. France 129 (2001), 339–356.

4° SÉRIE – TOME 49 – 2016 – N° 4

- [61] F. SAKAI, Weil divisors on normal surfaces, *Duke Math. J.* 51 (1984), 877–887.
- [62] V. V. SHOKUROV, Three-dimensional log perestroikas, *Izv. Ross. Akad. Nauk Ser. Mat.* 56 (1992), 105–203; English translation: *Russian Acad. Sci. Izv. Math.* 40 (1993), 95–202.
- [63] D. TOLEDO, Y. L. L. TONG, Green's theory of Chern classes and the Riemann-Roch formula, in *The Lefschetz centennial conference, Part I (Mexico City, 1984)*, Contemp. Math. 58, Amer. Math. Soc., Providence, RI, 1986, 261–275.
- [64] K. UENO, On compact analytic threefolds with nontrivial Albanese tori, Math. Ann. 278 (1987), 41–70.
- [65] J. VAROUCHAS, Kähler spaces and proper open morphisms, *Math. Ann.* 283 (1989), 13– 52.
- [66] C. VOISIN, *Théorie de Hodge et géométrie algébrique complexe*, Cours spécialisés **10**, Soc. Math. France, Paris, 2002.

(Manuscrit reçu le 20 avril 2015; accepté, après révision, le 22 septembre 2015.)

Frédéric CAMPANA Institut Élie Cartan Université Henri Poincaré BP 239 54506 Vandœuvre-lès-Nancy Cedex, France et Institut Universitaire de France E-mail: frederic.campana@univ-lorraine.fr

Andreas HÖRING Laboratoire de Mathématiques J.-A. Dieudonné UMR 7351 CNRS Université de Nice Sophia-Antipolis 06108 Nice Cedex 02, France E-mail: hoering@unice.fr

Thomas PETERNELL Mathematisches Institut Universität Bayreuth 95440 Bayreuth, Germany E-mail: thomas.peternell@uni-bayreuth.de