

quatrième série - tome 52 fascicule 4 juillet-août 2019

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Siddhartha SAHI & Hadi SALMASIAN

Quadratic Capelli operators and Okounkov polynomials

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Patrick BERNARD

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE

de 1883 à 1888 par H. DEBRAY

de 1889 à 1900 par C. HERMITE

de 1901 à 1917 par G. DARBOUX

de 1918 à 1941 par É. PICARD

de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} mars 2019

P. BERNARD

D. HARARI

S. BOUCKSOM

A. NEVES

R. CERF

J. SZEFTEL

G. CHENEVIER

S. VŨ NGỌC

Y. DE CORNULIER

A. WIENHARD

A. DUCROS

G. WILLIAMSON

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,

45, rue d'Ulm, 75230 Paris Cedex 05, France.

Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.

annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France

Case 916 - Luminy

13288 Marseille Cedex 09

Tél. : (33) 04 91 26 74 64

Fax : (33) 04 91 41 17 51

email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 420 euros.

Abonnement avec supplément papier :

Europe : 551 €. Hors Europe : 620 € (\$ 930). Vente au numéro : 77 €.

© 2019 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Stéphane Seuret

Périodicité : 6 n^{os} / an

QUADRATIC CAPELLI OPERATORS AND OKOUNKOV POLYNOMIALS

BY SIDDHARTHA SAHI AND HADI SALMASIAN

ABSTRACT. — Let Z be the symmetric cone of $r \times r$ positive definite Hermitian matrices over a real division algebra \mathbb{F} . Then Z admits a natural family of invariant differential operators—the *Capelli operators* C_λ —indexed by partitions λ of length at most r , whose eigenvalues are specializations of Knop-Sahi interpolation polynomials.

In this paper we consider a double fibration $Y \leftarrow X \rightarrow Z$ where Y is the Grassmanian of r -dimensional subspaces of \mathbb{F}^n with $n \geq 2r$. Using this we construct a family of invariant differential operators $D_{\lambda,s}$ on Y that we refer to as *quadratic Capelli operators*. Our main result shows that the eigenvalues of the $D_{\lambda,s}$ are specializations of Okounkov interpolation polynomials.

RÉSUMÉ. — Soit Z le cône symétrique de matrices de tailles $r \times r$ hermitiennes positives sur une algèbre de division réelle \mathbb{F} . Alors Z admet une famille naturelle d'opérateurs différentiels invariants — les *Opérateurs de Capelli* C_λ — indexés par des partitions λ de longueur au plus r , dont les valeurs propres sont des spécialisations de polynômes d'interpolation Knop-Sahi.

Dans cet article, nous considérons une double fibration $Y \leftarrow X \rightarrow Z$ où Y est la variété grassmannienne des sous-espaces de dimension r de \mathbb{F}^n avec $n \geq 2r$. En utilisant cela, nous construisons une famille d'opérateurs différentiels invariants $D_{\lambda,s}$ sur Y que nous appelons opérateurs de Capelli *quadratiques*. Notre résultat principal montre que les valeurs propres des $D_{\lambda,s}$ sont des spécialisations de polynômes d'interpolation Okounkov.

1. Introduction

Let $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ be a real division algebra. Fix integers r and n such that $1 \leq r \leq \frac{n}{2}$. Let Y be the Grassmannian of r -dimensional subspaces of \mathbb{F}^n , and let Z be the symmetric cone of $r \times r$ positive definite Hermitian \mathbb{F} -matrices. Then one has a double fibration

$$\begin{array}{ccc} & X & \\ \varphi \swarrow & & \searrow \psi \\ Y & & Z, \end{array}$$

where X is the space of $n \times r$ matrices of \mathbb{F} -rank r . For $x \in X$, $\varphi(x)$ is the column space (or range) of x , while $\psi(x) := x^\dagger x$, where x^\dagger denotes the \mathbb{F} -Hermitian adjoint of x .

One can give another description of the above structure in terms of the groups

$$G_m := \mathrm{GL}_m(\mathbb{F}), \quad K_m := \mathrm{U}_m(\mathbb{F}) := \left\{ g \in G_m : g^\dagger g = I_{m \times m} \right\}.$$

The groups K_n and G_r act on X by matrix multiplication on the left and right respectively, and the maps $X \xrightarrow{\psi} Z$ and $X \xrightarrow{\varphi} Y$ are simply the corresponding quotient maps. Moreover, $X \xrightarrow{\varphi} Y$ is a principal G_r -bundle, while $X \xrightarrow{\psi} Z$ is a fibration whose fibers are isomorphic to the Stiefel manifold K_n/K_{n-r} . Also, since the actions of K_n and G_r on X commute, it follows that G_r acts on Z , and K_n acts on Y . In fact, Y and Z are symmetric spaces for the latter actions. More precisely, we have

$$Y \simeq K_n/(K_r \times K_{n-r}) \quad \text{and} \quad Z \simeq G_r/K_r.$$

The cone Z is a symmetric space of type A , and admits an important basis of G_r -invariant differential operators C_λ , indexed by partitions $\lambda \in \mathcal{P}_r$, where

$$\mathcal{P}_r := \{(\lambda_1, \dots, \lambda_r) \in \mathbb{Z}^r : \lambda_1 \geq \dots \geq \lambda_r \geq 0\}.$$

The operators C_λ were first studied by the first author in [20], and were referred to as *Capelli operators*. It is known that the spectrum of C_λ is given by specialization of Knop-Sahi type A interpolation polynomials [20], [12], [30].

On the other hand, Y is a compact symmetric space of type BC . In this paper, we use the above double fibration to construct a family of K_n -invariant differential operators $D_{\lambda,s}$ on Y that correspond to the Capelli operators C_λ . We call the operators $D_{\lambda,s}$ the *quadratic Capelli operators* because they are obtained from C_λ by pullback of the quadratic map ψ . Our main result proves that the spectrum of $D_{\lambda,s}$ is given by specialization of the *Okounkov type BC* interpolation polynomials $P_\lambda(x; \tau, \alpha)$ (see [14, Sec. 5.3] and [19]).

To describe our main result precisely, we begin by introducing some notation. Set $K := K_n$ and $M := K_r \times K_{n-r} \subset K$, so that $Y \simeq K/M$. The group K acts by left translation on $C^\infty(Y)$, the space of complex-valued smooth functions on Y . The operators $D_{\lambda,s}$ leave the subspace $C^\infty(Y)_{K\text{-finite}}$ of K -finite vectors invariant. By standard results from the theory of compact symmetric spaces (for example, see [5, Chap. V]), $C^\infty(Y)_{K\text{-finite}}$ decomposes as a multiplicity-free direct sum of irreducible M -spherical K -modules, which are naturally parametrized by partitions $\mu \in \mathcal{P}_r$. Our next goal is to describe this parametrization. Let \mathfrak{k} and \mathfrak{m} denote the Lie algebras of K and M . Fix a Cartan decomposition $\mathfrak{k} = \mathfrak{m} \oplus \mathfrak{p}$. Let $\mathfrak{a} \subseteq \mathfrak{p}$ be a Cartan subspace, and let \mathfrak{h} be a Cartan subalgebra of \mathfrak{k} such that $\mathfrak{a} \subseteq \mathfrak{h}$. Then $\mathfrak{h} = \mathfrak{t} \oplus \mathfrak{a}$, where $\mathfrak{t} := \mathfrak{h} \cap \mathfrak{m}$. Set $\mathfrak{k}_{\mathbb{C}} := \mathfrak{k} \otimes_{\mathbb{R}} \mathbb{C}$, $\mathfrak{h}_{\mathbb{C}} := \mathfrak{h} \otimes_{\mathbb{R}} \mathbb{C}$, $\mathfrak{a}_{\mathbb{C}} := \mathfrak{a} \otimes_{\mathbb{R}} \mathbb{C}$, and $\mathfrak{t}_{\mathbb{C}} := \mathfrak{t} \otimes_{\mathbb{R}} \mathbb{C}$. The restricted root system $\Sigma := \Sigma(\mathfrak{k}_{\mathbb{C}}, \mathfrak{a}_{\mathbb{C}})$ is of type BC_r . We choose a positive system $\Sigma^+ \subset \Sigma$ and a basis e_1, \dots, e_r of $\mathfrak{a}_{\mathbb{C}}^*$ such that the multiplicity m_α of every $\alpha \in \Sigma^+$ is given in terms of n , r , and $d := \dim \mathbb{F}$ in Table 1 below.

α	$e_i, 1 \leq i \leq r$	$e_i \pm e_j, 1 \leq i < j \leq r$	$2e_i, 1 \leq i \leq r$
m_α	$d(n-2r)$	d	$d-1$

Table 1.

We also choose a positive system for the root system $\Delta := \Delta(\mathfrak{k}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ which is compatible with Σ^+ . Let $\tilde{\mu} \in \mathfrak{h}_{\mathbb{C}}^*$. By the Cartan-Helgason theorem, $\tilde{\mu}$ is the highest weight of an irreducible M -spherical K -module if and only if

$$(1) \quad \tilde{\mu}|_{\mathfrak{t}_{\mathbb{C}}} = 0 \text{ and } \tilde{\mu}|_{\mathfrak{a}_{\mathbb{C}}} = \sum_{i=1}^r 2\mu_i e_i, \text{ where } \mu := (\mu_1, \dots, \mu_r) \in \mathcal{P}_r.$$

REMARK 1.1. – Assume that $\mathbb{F} = \mathbb{R}$. Then K is disconnected, and if K° denotes the connected component of identity of K , then $M \cap K^\circ$ is also disconnected. Therefore the Cartan-Helgason theorem as stated for instance in [5, Cor. V.4.2] does not apply immediately to the case $\mathbb{F} = \mathbb{R}$. However, one can use the refinement of the Cartan-Helgason theorem for the pair $(K^\circ, M \cap K^\circ)$, given in [4, Sec. 12.3.2], as well as the description of irreducible representations of K in terms of irreducible representations of K° , given in [4, Thm 5.5.23], to obtain the condition (1).

From now on, we denote the M -spherical K -module with highest weight $\tilde{\mu}$ satisfying (1) by V_μ . Therefore as K -modules,

$$C^\infty(Y)_{K\text{-finite}} \simeq \bigoplus_{\mu \in \mathcal{P}_r} V_\mu.$$

The operator $D_{\lambda,s}$ acts on V_μ by the scalar

$$c_{\lambda,s}(\mu) := \text{HC}(D_{\lambda,s})(\tilde{\mu}|_{\mathfrak{a}_{\mathbb{C}}} + \rho),$$

where $\rho := \frac{1}{2} \sum_{\alpha \in \Sigma^+} \alpha$, $\tilde{\mu}$ is the highest weight of V_μ , and $\text{HC} : \mathbf{D}_K(Y) \rightarrow \mathcal{P}(\mathfrak{a}_{\mathbb{C}}^*)^W$ is the Harish-Chandra homomorphism from the algebra $\mathbf{D}_K(Y)$ of K -invariant differential operators on Y onto the algebra of polynomials on $\mathfrak{a}_{\mathbb{C}}^*$ that are invariant under the action of the restricted Weyl group W .

We now recall the definition of the Okounkov polynomials $P_\lambda(x; \tau, \alpha)$. Let $\mathbb{k} := \mathbb{C}(\tau, \alpha)$ denote the field of rational functions in τ and α . Let $\delta, \mathbb{1} \in \mathcal{P}_r$ and $\varrho_{\tau,\alpha} \in \mathbb{k}^r$ be defined by

$$(2) \quad \delta := (r - 1, \dots, 0), \quad \mathbb{1} := (1, \dots, 1), \quad \varrho_{\tau,\alpha} := \tau\delta + \alpha\mathbb{1}.$$

For $\lambda \in \mathcal{P}_r$, we define $|\lambda| := \sum_{i=1}^r \lambda_i$. Up to a scalar, $P_\lambda(x; \tau, \alpha) \in \mathbb{k}[x_1, \dots, x_r]$ is the unique polynomial of degree $2|\lambda|$, which is invariant under permutations and sign changes of x_1, \dots, x_r , and satisfies

$$P_\lambda(\mu + \varrho_{\tau,\alpha}; \tau, \alpha) = 0$$

for every $\mu \in \mathcal{P}_r$ such that $|\mu| \leq |\lambda|$ and $\mu \neq \lambda$ (for more details, see Section 4).

Recall that $d := \dim \mathbb{F}$. Let $\mathfrak{i} : \mathbb{C}^r \rightarrow \mathfrak{a}_{\mathbb{C}}^*$ be the linear map defined by $\mathfrak{i}(e^i) := 2e_i$ for $1 \leq i \leq r$, where e^1, \dots, e^r are the standard basis vectors of \mathbb{C}^r (therefore $\mathfrak{i}(\mathcal{P}_r)$ is the set of restrictions to $\mathfrak{a}_{\mathbb{C}}$ of highest weights of M -spherical K -modules). Set

$$\varrho := \mathfrak{i}^{-1}(\rho).$$

A simple calculation yields

$$(3) \quad \varrho = (\varrho_1, \dots, \varrho_r) \text{ where } \varrho_i := \frac{dn}{4} - \frac{1}{2} - \frac{d(i-1)}{2} \text{ for every } 1 \leq i \leq r.$$

We are now ready to state our main theorem.

THEOREM 1.2. – For every $\lambda, \mu \in \mathcal{P}_r$ and every $s \in \mathbb{C}$, the operator $D_{\lambda,s}$ acts on V_μ by the scalar

$$(4) \quad c_{\lambda,s}(\mu) := \gamma_\lambda P_\lambda \left(\mu + \varrho; \frac{d}{2}, s - \varrho_1 \right),$$

where γ_λ is a certain explicit constant, defined in (18) below.

We now briefly sketch the strategy behind the proof of Theorem 1.2. We first show that the equality (4) holds up to a scalar multiple. In view of the characterization of the Okounkov polynomials P_λ , it suffices to prove that $c_{\lambda,s}(x - \varrho)$ is a polynomial in x and s which has the same degree and vanishing property as $P_\lambda(x; \tau, \alpha)$ for $\tau := \frac{d}{2}$ and $\alpha := s - \varrho_1$ (see Proposition 4.3 and Proposition 5.6). It will be seen that verifying the pertinent vanishing property can be reduced to a slightly weaker one, that is, to show that the operators $D_\lambda := D_{\lambda,0}$ vanish on certain $V_\mu \subset C^\infty(Y)_{K\text{-finite}}$. Up to this point, the strategy is the same as the one in the case of Knop-Sahi type A polynomials. However, the proof of the vanishing property of the operators D_λ is more subtle than the type A case, in that it does *not* follow from a direct reasoning that is based on orders of differential operators. In addition, the ρ -shift of the symmetric space Y and $\varrho_{\tau,\alpha}$ are not identical. Rather, they are related to each other as in Remark 5.5.

To overcome these difficulties, we need to use the fact that $c_{\lambda,s}(x - \varrho)$ is symmetric under permutations and sign changes of its variables, and therefore $c_{\lambda,s}(x - \varrho) = 0$ for $x := (x_1, \dots, x_r)$ if and only if $c_{\lambda,s}(\tilde{x} - \varrho) = 0$ for $\tilde{x} := (-x_r, \dots, -x_1)$. The latter observation results in an equivalent form of the vanishing condition for $c_{\lambda,s}$, which is verified in the proof of Proposition 5.6 using differential operator techniques and branching rules. The branching rule from G_n to K_n that we need in the proof of the vanishing property of D_λ is the Littlewood-Richardson rule for $\mathbb{F} = \mathbb{C}$ and the Littlewood Restriction theorem for $\mathbb{F} = \mathbb{R}, \mathbb{H}$. As a result, some of our proofs are divided into two cases, but they lead to uniform statements.

The last step in the proof of our main result is to determine the scalar γ_λ that relates the two sides of (4). To this end, we use the fact that the top degree homogeneous term of $P_\lambda(x; \tau, \alpha)$ is a Jack polynomial. Using a trick which relies on an identity for Jack polynomials (see [28, Prop. 2.3]), calculation of γ_λ for general λ is reduced to the case where λ corresponds to the 1×1 Young diagram. Even this special case requires a rather intricate computation to express the corresponding quadratic Capelli operator in terms of the Casimir operators of G_n and K_n . This is carried out in the appendix.

We now describe the relation between Theorem 1.2 and earlier results on invariant differential operators on symmetric spaces. The Capelli operators C_λ were originally studied in connection to the famous Capelli identity, which has also been considerably generalized by Howe and Umeda [9] from the viewpoint of multiplicity-free actions, and by Kostant and the first author [15], [16] from the viewpoint of Jordan algebras. For $\mathbb{F} = \mathbb{R}$ and $\lambda = \mathbb{1}$, the operator $D_{\mathbb{1},-1}$ was first considered by Howe and Lee in [7], who computed its spectrum for $r = 2$ and asked for the determination of the spectrum for general r . This was solved by the first author more generally for $D_{m\mathbb{1},s}$ and $\mathbb{F} = \mathbb{R}$, [22], and subsequently by Zhang and the first author [25] for arbitrary \mathbb{F} , where a connection with the Radon transform was also established. The explicit form of the answer in [25] gave us the first hint that the general

situation might have something to do with the Okounkov polynomials. The result of [25] on the spectrum of $D_{m_1,s}$ is indeed a special case of our Theorem 1.2.

Finally, we say a few words about prospects for future research that emerge from this work. Quite recently, Zhang and the first author established another link between Okounkov interpolation polynomials and the spectrum of Shimura operators on Hermitian symmetric spaces [26]. It would be interesting to understand the connection between our main result and the results of [26]. Also, in view of our recent work [23] on the Capelli eigenvalue problem in the case of the supersymmetric pairs $(\mathfrak{gl}(m|m') \times \mathfrak{gl}(m|m'), \mathfrak{gl}(m|m'))$ and $(\mathfrak{gl}(m|2m'), \mathfrak{osp}(m|2m'))$, it is natural to ask whether Theorem 1.2 can also be extended to the setting of Lie superalgebras. This is likely to involve the deformed BC interpolation polynomials of [27]. We remark that in forthcoming papers [1] and [24], we extend the results of [23] to the setting of multiplicity-free actions obtained from Jordan superalgebras. Another interesting problem is to extend the Littlewood Restriction theorem (see Proposition 2.12) to the super setting, namely to $(\mathfrak{gl}(m|2m'), \mathfrak{osp}(m|2m'))$. We are planning to study these problems in the near future.

Acknowledgement. – The authors thank Kyo Nishiyama and Nolan Wallach for helpful e-mail correspondences. The research of Siddhartha Sahi was partially supported by a Simons Foundation grant (509766) and the research of Hadi Salmasian by an NSERC Discovery Grant (RGPIN-2013-355464). Part of this work was carried out during the Workshop on Hecke Algebras and Lie Theory held at the University of Ottawa during May 12–15, 2016. The authors thank the National Science Foundation (DMS-162350), the Fields Institute, and the University of Ottawa for funding this workshop.

2. Parametrization of representations by partitions

In this article, we will need various parametrizations of finite dimensional representations of G_n , K_n , and G_r by partitions. Instead of working with representations of these real Lie groups, it will be more convenient to work with representations of their complexifications.

Let $W := \text{Mat}_{n \times r}(\mathbb{F})$ denote the space of $n \times r$ matrices with entries in \mathbb{F} . Furthermore, set

$$A := \{x \in \text{Mat}_{r \times r}(\mathbb{F}) : x^\dagger = x\}.$$

Then $X \subset W$ and $Z := \{w^\dagger w : w \in X\} \subset A$ are open. The $G_n \times G_r$ -action on X is the restriction of the $G_n \times G_r$ -action on W given by

$$(g_1, g_2) \cdot w := g_1 w g_2^{-1} \text{ for } (g_1, g_2) \in G_n \times G_r, w \in W.$$

The G_r -action on Z is the restriction of the G_r -action on A given by

$$g \cdot x := (g^\dagger)^{-1} x g^{-1} \text{ for } g \in G_r, x \in A.$$

Let \mathbf{G}_n and \mathbf{G}_r denote the complexifications of the real Lie groups G_n and G_r . Similarly, let $\mathbf{K}_n \subset \mathbf{G}_n$ denote the complexification of K_n .

REMARK 2.1. – From now on, we need to fix an embedding of quaternionic matrices into complex matrices twice as big. For integers $k, m \geq 1$, let $x := a + b\mathbf{j} \in \text{Mat}_{k \times m}(\mathbb{H})$, where $a, b \in \text{Mat}_{k \times m}(\mathbb{C})$. We set

$$\check{x} := \begin{bmatrix} a & -\bar{b} \\ b & \bar{a} \end{bmatrix} \in \text{Mat}_{2k \times 2m}(\mathbb{C}).$$

REMARK 2.2. – The matrix realizations of the embeddings $\mathbf{i}_n : G_n \hookrightarrow \mathbf{G}_n$ and $\mathbf{i}_r : G_r \hookrightarrow \mathbf{G}_r$ are as follows. If $\mathbb{F} = \mathbb{R}$, then $\mathbf{G}_n \simeq \text{GL}_n(\mathbb{C})$ and $\mathbf{i}_n : \text{GL}_n(\mathbb{R}) \rightarrow \text{GL}_n(\mathbb{C})$ is the obvious map. If $\mathbb{F} = \mathbb{C}$, then $\mathbf{G}_n \simeq \text{GL}_n(\mathbb{C}) \times \text{GL}_n(\mathbb{C})$ and

$$\mathbf{i}_n : \text{GL}_n(\mathbb{C}) \rightarrow \text{GL}_n(\mathbb{C}) \times \text{GL}_n(\mathbb{C})$$

is the map $g \mapsto ((g^*)^{-1}, g)$, where $g^* := \bar{g}^T$ is the adjoint of g . If $\mathbb{F} = \mathbb{H}$, then $\mathbf{G}_n \simeq \text{GL}_{2n}(\mathbb{C})$ and $\mathbf{i}_n : \text{GL}_n(\mathbb{H}) \rightarrow \text{GL}_{2n}(\mathbb{C})$ is the map $g \mapsto \check{g}$. The matrix realization of \mathbf{K}_n as a subgroup of \mathbf{G}_n is as follows. If $\mathbb{F} = \mathbb{R}$, then $\mathbf{K}_n = \{g \in \text{GL}_n(\mathbb{C}) : g^T g = I_{n \times n}\}$. If $\mathbb{F} = \mathbb{C}$, then $\mathbf{K}_n = \{(g, g) : g \in \text{GL}_n(\mathbb{C})\}$. Finally, if $\mathbb{F} = \mathbb{H}$, then

$$\mathbf{K}_n = \{g \in \text{GL}_{2n}(\mathbb{C}) : g^T J_n g = J_n\},$$

where

$$(5) \quad J_n := \begin{bmatrix} 0 & I_{n \times n} \\ -I_{n \times n} & 0 \end{bmatrix}.$$

The definition of the embedding $\mathbf{i}_r : G_r \rightarrow \mathbf{G}_r$ is similar to that of \mathbf{i}_n . To help the reader, we summarize the information about \mathbf{G}_n and \mathbf{K}_n in Table 2 below.

\mathbb{F}	\mathbf{G}_n	\mathbf{K}_n	Realization of \mathbf{K}_n in \mathbf{G}_n
\mathbb{R}	$\text{GL}_n(\mathbb{C})$	$\text{O}_n(\mathbb{C})$	$\{g \in \text{GL}_n(\mathbb{C}) : g^T g = I_{n \times n}\}$
\mathbb{C}	$\text{GL}_n(\mathbb{C}) \times \text{GL}_n(\mathbb{C})$	$\text{GL}_n(\mathbb{C})$	$\{(g, g) : g \in \text{GL}_n(\mathbb{C})\}$
\mathbb{H}	$\text{GL}_{2n}(\mathbb{C})$	$\text{Sp}_{2n}(\mathbb{C})$	$\{g \in \text{GL}_{2n}(\mathbb{C}) : g^T J_n g = J_n\}$

Table 2.

Set $W_{\mathbb{C}} := W \otimes_{\mathbb{R}} \mathbb{C}$ and $A_{\mathbb{C}} := A \otimes_{\mathbb{R}} \mathbb{C}$. The map $X \xrightarrow{\psi} Z$ has a unique holomorphic extension $W_{\mathbb{C}} \xrightarrow{\psi} A_{\mathbb{C}}$. For an explicit description of $A_{\mathbb{C}}$, $W_{\mathbb{C}}$, and the map $\psi : W_{\mathbb{C}} \rightarrow A_{\mathbb{C}}$, see Table 3 below.

\mathbb{F}	$A_{\mathbb{C}}$	$W_{\mathbb{C}}$	ψ
\mathbb{R}	$\text{Sym}_{r \times r}(\mathbb{C})$	$\text{Mat}_{n \times r}(\mathbb{C})$	$x \mapsto x^T x$
\mathbb{C}	$\text{Mat}_{r \times r}(\mathbb{C})$	$\text{Mat}_{n \times r}(\mathbb{C}) \oplus \text{Mat}_{n \times r}(\mathbb{C})$	$(x_1, x_2) \mapsto x_1^T x_2$
\mathbb{H}	$\text{Skew}_{2r \times 2r}(\mathbb{C})$	$\text{Mat}_{2n \times 2r}(\mathbb{C})$	$x \mapsto -x^T J_n x$

Table 3.

In Table 3, $\text{Sym}_{r \times r}(\mathbb{C})$ denotes the space of complex symmetric $r \times r$ matrices and $\text{Skew}_{2r \times 2r}(\mathbb{C})$ denotes the space of complex skew symmetric $2r \times 2r$ matrices.

REMARK 2.3. – The matrix realizations of the maps $A \rightarrow A_{\mathbb{C}}$ and $W \rightarrow W_{\mathbb{C}}$ are as follows. For $\mathbb{F} = \mathbb{R}$ and $\mathbb{F} = \mathbb{C}$, the map $A \hookrightarrow A_{\mathbb{C}}$ is the obvious embedding, and for $\mathbb{F} = \mathbb{H}$, it is the map $a \mapsto -J_r a$, where J_r is defined similar to (5). In fact the map $A \hookrightarrow A_{\mathbb{C}}$ is related to realization of Euclidean Jordan algebras (see [2, Sec. VIII.5]). As for $W \hookrightarrow W_{\mathbb{C}}$, it is the obvious embedding if $\mathbb{F} = \mathbb{R}$, the map $w \mapsto (\bar{w}, w)$ if $\mathbb{F} = \mathbb{C}$, and the map $w \mapsto \check{w}$ if $\mathbb{F} = \mathbb{H}$.

The action of G_r on A extends uniquely to a holomorphic action of \mathbf{G}_r on $A_{\mathbb{C}}$. Similarly, the action of $G_n \times G_r$ on W extends uniquely to a holomorphic action of $\mathbf{G}_n \times \mathbf{G}_r$ on $W_{\mathbb{C}}$. These holomorphic actions are explicitly described in Table 4 below.

\mathbb{F}	$\mathbf{G}_r \circ A_{\mathbb{C}}$	$(\mathbf{G}_n \times \mathbf{G}_r) \circ W_{\mathbb{C}}$
\mathbb{R}	$S^2(\mathbb{C}^r)^*$	$\mathbb{C}^n \otimes (\mathbb{C}^r)^*$
\mathbb{C}	$\mathbb{C}^r \otimes (\mathbb{C}^r)^*$	$((\mathbb{C}^n)^* \otimes \mathbb{C}^r) \oplus (\mathbb{C}^n \otimes (\mathbb{C}^r)^*)$
\mathbb{H}	$\Lambda^2(\mathbb{C}^{2r})^*$	$\mathbb{C}^{2n} \otimes (\mathbb{C}^{2r})^*$

Table 4.

DEFINITION 2.4. – For every integer $m \geq 1$, let $H_m \subset GL_m(\mathbb{C})$ denote the standard Cartan subgroup of diagonal matrices, and let $B_m \subset GL_m(\mathbb{C})$ denote the standard Borel subgroup of upper triangular matrices. Let $\varepsilon_1, \dots, \varepsilon_m$ denote the standard generators of the weight lattice of $GL_m(\mathbb{C})$. For every $\lambda := (\lambda_1, \dots, \lambda_m) \in \mathbb{Z}^m$ such that $\lambda_1 \geq \dots \geq \lambda_m$, we denote the $GL_m(\mathbb{C})$ -module with B_m -highest weight $\sum_{i=1}^m \lambda_i \varepsilon_i$ by M_{λ} .

DEFINITION 2.5. – The standard Borel subgroup of \mathbf{G}_m will be denoted by \mathbf{B}_m . In cases $\mathbb{F} = \mathbb{R}$, $\mathbb{F} = \mathbb{C}$, and $\mathbb{F} = \mathbb{H}$, the group $\mathbf{B}_m \subset \mathbf{G}_m$ equals B_m , $B_m \times B_m$, and B_{2m} . The standard Cartan subgroup of \mathbf{G}_m will be denoted by \mathbf{H}_m .

REMARK 2.6. – For every integer $m \geq 1$, we define

$$\mathcal{P}_m := \{(\lambda_1, \dots, \lambda_m) \in \mathbb{Z}^m : \lambda_1 \geq \dots \geq \lambda_m \geq 0\}.$$

From now on, we will denote the length (that is, the number of nonzero parts) of a partition $\lambda \in \mathcal{P}_m$ by $\ell(\lambda)$. For two partitions $\lambda \in \mathcal{P}_m$ and $\mu \in \mathcal{P}_k$, where $k, m \geq 1$, we write $\lambda \subseteq \mu$ if and only if $\ell(\lambda) \leq \ell(\mu)$ and $\lambda_i \leq \mu_i$ for every $1 \leq i \leq \ell(\lambda)$.

Let $\mathcal{P}(A)$ and $\mathcal{P}(W)$ denote the \mathbb{C} -algebras of polynomials on $A_{\mathbb{C}}$ and $W_{\mathbb{C}}$. The canonical G_r -action on $\mathcal{P}(A)$, given by $g \cdot f(a) := f(g^{-1} \cdot a)$ for $g \in G_r$, $f \in \mathcal{P}(A)$, and $a \in A$, extends uniquely to a holomorphic \mathbf{G}_r -action on $\mathcal{P}(A)$. Similarly, the canonical $G_n \times G_r$ -action on $\mathcal{P}(W)$ extends uniquely to a holomorphic $\mathbf{G}_n \times \mathbf{G}_r$ -action on $\mathcal{P}(W)$. The pullback

$$(6) \quad \psi^* : \mathcal{P}(A) \rightarrow \mathcal{P}(W), \quad f \mapsto f \circ \psi$$

is a \mathbf{G}_r -equivariant embedding of \mathbb{C} -algebras. The image of ψ^* is precisely described by the First Fundamental Theorem of invariant theory [4, Sec. 5.2.1]. In particular, $\psi^*(\mathcal{P}(A)) = \mathcal{P}(W)^{\mathbf{K}_n}$.

By classical invariant theory (for example see [4]), $\mathcal{P}(A)$ decomposes into a direct sum of irreducible \mathbf{G}_r -modules which are naturally parametrized by partitions $\lambda \in \mathcal{P}_r$. Thus,

$$(7) \quad \mathcal{P}(A) \simeq \bigoplus_{\lambda \in \mathcal{P}_r} F_\lambda,$$

where F_λ is the irreducible \mathbf{G}_r -module corresponding to $\lambda \in \mathcal{P}_r$. In fact $F_\lambda \simeq M_\lambda^* \otimes M_\lambda$ if $\mathbb{F} = \mathbb{C}$, and $F_\lambda \simeq M_{\lambda^\bullet}$ if $\mathbb{F} = \mathbb{R}, \mathbb{H}$, where $\lambda^\bullet := (2\lambda_1, \dots, 2\lambda_r) \in \mathcal{P}_r$ if $\mathbb{F} = \mathbb{R}$, and $\lambda^\bullet := (\lambda_1, \lambda_1, \dots, \lambda_r, \lambda_r) \in \mathcal{P}_{2r}$ if $\mathbb{F} = \mathbb{H}$.

The map (6) is \mathbf{G}_r -equivariant, and therefore F_λ occurs as a \mathbf{G}_r -submodule of $\mathcal{P}(W)$ for every $\lambda \in \mathcal{P}_r$. Therefore by the well known $(\mathrm{GL}_n, \mathrm{GL}_r)$ duality (see [6, Sec. 2.1] or [4, Sec. 5.6.2]), for every $\lambda \in \mathcal{P}_r$ there exists a unique irreducible \mathbf{G}_n -module E_λ such that $E_\lambda \otimes F_\lambda$ occurs in $\mathcal{P}(W)$ as a $\mathbf{G}_n \times \mathbf{G}_r$ -submodule.

REMARK 2.7. – Let $\mathcal{P}_{\mathbf{K}_n}(W)$ denote the direct sum of irreducible \mathbf{K}_n -spherical \mathbf{G}_n -submodules of $\mathcal{P}(W)$. Then indeed

$$\mathcal{P}_{\mathbf{K}_n}(W) \simeq \bigoplus_{\lambda \in \mathcal{P}_r} E_\lambda \otimes F_\lambda \text{ as } \mathbf{G}_n \times \mathbf{G}_r\text{-modules.}$$

REMARK 2.8. – For every integer $l \geq 1$, let $S_{l \times l}$ be the $l \times l$ matrix with 1's in $(i, l - i + 1)$ -entry for every $1 \leq i \leq l$, and with 0's elsewhere. Consider $g_\circ \in \mathbf{G}_n$ defined as follows. If $\mathbb{F} = \mathbb{R}$, then we set

$$g_\circ := \begin{cases} \begin{bmatrix} \frac{1+i}{2} I_{l \times l} & \frac{1-i}{2} S_{l \times l} \\ \frac{1-i}{2} S_{l \times l} & \frac{1+i}{2} I_{l \times l} \end{bmatrix} & \text{for } n = 2l, \text{ and } \\ \begin{bmatrix} \frac{1+i}{2} I_{l \times l} & 0_{l \times 1} & \frac{1-i}{2} S_{l \times l} \\ 0_{1 \times l} & 1 & 0_{1 \times l} \\ \frac{1-i}{2} S_{l \times l} & 0_{l \times 1} & \frac{1+i}{2} I_{l \times l} \end{bmatrix} & \text{for } n = 2l + 1. \end{cases}$$

If $\mathbb{F} = \mathbb{C}$, then we set g_\circ equal to the identity element of \mathbf{G}_n . Finally, if $\mathbb{F} = \mathbb{H}$, then we set

$$g_\circ := \begin{bmatrix} S_{n \times n} & 0 \\ 0 & I_{n \times n} \end{bmatrix}.$$

We remark that when $\mathbb{F} = \mathbb{R}, \mathbb{H}$, the map $\mathbf{K}_n \rightarrow \mathbf{G}_n$, $g \mapsto g_\circ g g_\circ^{-1}$ is the embedding $\mathrm{O}_n(\mathbb{C}) \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ or $\mathrm{Sp}_{2n}(\mathbb{C}) \hookrightarrow \mathrm{GL}_{2n}(\mathbb{C})$ that is given in [4, Sec. 2.1.2]. In particular, in all cases $(g_\circ^{-1} \mathbf{H}_n g_\circ) \cap \mathbf{K}_n$ and $(g_\circ^{-1} \mathbf{B}_n g_\circ) \cap \mathbf{K}_n$ are Cartan and Borel subalgebras of \mathbf{K}_n .

Set $\mathbf{K} := \mathbf{K}_n$, and let $\mathbf{M} \subset \mathbf{K}$ denote the complexification of \mathbf{M} . The M -spherical K -module V_μ is naturally also an \mathbf{M} -spherical \mathbf{K} -module. By comparing the calculation of highest weights of \mathbf{M} -spherical \mathbf{K} -modules in [4, Sec. 12.3.2] (the pertinent cases are types BDI, AIII, and CII) with the parametrization of representations of \mathbf{K} by partitions that uses generalized Schur-Weyl duality (see [4, Thm 10.2.9] and [4, Thm 10.2.12]), it follows that for every $\mu \in \mathcal{P}_r$, the module V_μ is isomorphic to the \mathbf{K} -submodule of E_μ^* generated by $g_\circ^{-1} \cdot v_\lambda$, where v_λ denotes the \mathbf{B}_n -highest weight of E_μ^* .

DEFINITION 2.9. – Let G be a group, and let E and F be G -modules. We set

$$[E : F]_G := \dim \mathrm{Hom}_G(E, F).$$

REMARK 2.10. – In the following, we will need the Littlewood-Richardson Rule, which we now recall (for a more elaborate reference, see for example [18, Sec. I.9]). For a semi-standard skew tableau T , the word $w(T)$ corresponding to T is defined as the sequence of integers obtained by reading the contents of boxes of T from right to left and from top to bottom. A word $w_1 \cdots w_k$ in letters $\{1, \dots, N\}$ is called a lattice permutation word if for every $1 \leq i \leq k$ and every $1 \leq j \leq N - 1$, the number of occurrences of j in $w_1 \dots w_i$ is greater than or equal to the number of occurrences of $j + 1$. Now let $\lambda, \mu, \nu \in \mathcal{P}_m$, where $m \geq 1$. The Littlewood-Richardson Rule states that $[M_\nu : M_\lambda \otimes M_\mu]_{\text{GL}_m(\mathbb{C})}$ is equal to the number of tableaux T of shape $\nu \setminus \mu$ and weight λ such that $w(T)$ is a lattice permutation word. In particular, if $[M_\nu : M_\lambda \otimes M_\mu]_{\text{GL}_m(\mathbb{C})} \neq 0$, then $\mu, \lambda \subseteq \nu$.

LEMMA 2.11. – Let $\lambda, \mu \in \mathcal{P}_r$. Then $[V_\mu : E_\lambda]_{\mathbf{K}_n} = [V_\mu : E_\lambda^*]_{\mathbf{K}_n}$.

Proof. – It is enough to show that $E_\lambda \simeq E_\lambda^*$ as \mathbf{K}_n -modules. Let $\theta_n : \mathbf{G}_n \rightarrow \mathbf{G}_n$ denote the automorphism of \mathbf{G}_n that is obtained by holomorphic extension of the Cartan involution $g \mapsto (g^\dagger)^{-1}$ of G_n . Let $E_\lambda^{\theta_n}$ be the \mathbf{G}_n -module that results from twisting E_λ by θ_n . Then $E_\lambda^{\theta_n} \simeq E_\lambda^*$ as \mathbf{G}_n -modules. Moreover, since θ_n fixes \mathbf{K}_n pointwise, $E_\lambda \simeq E_\lambda^{\theta_n}$ as \mathbf{K}_n -modules. The \mathbf{K}_n -module isomorphism $E_\lambda \simeq E_\lambda^*$ now follows immediately. \square

When $\mathbb{F} = \mathbb{C}$, branching from \mathbf{G}_n to \mathbf{K}_n is described by the Littlewood-Richardson Rule. The next proposition is a branching from \mathbf{G}_n to \mathbf{K}_n when $\mathbb{F} \neq \mathbb{C}$.

PROPOSITION 2.12. – Assume that $\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{H}$, and let $\lambda, \mu \in \mathcal{P}_r$. Then

$$(8) \quad [V_\mu : E_\lambda]_{\mathbf{K}_n} = \sum_{\xi \in \mathcal{P}_r} [E_\lambda : E_\mu \otimes E_\xi]_{\mathbf{G}_n}.$$

Proof. – The statement follows as a special case of the Littlewood Restriction theorem, which was first proved in [17] (see also [13], [3], and [8, Sec. 1.3]). We now outline the calculations that are needed to deduce the proposition from the Littlewood Restriction theorem. We remark that the \mathbf{G}_n -modules that appear in the statement of the Littlewood Restriction theorem are polynomial representations, whereas the E_λ are indeed contragredients of polynomial representations. To get around this issue, we note that we can replace the left hand side of (8) by $[V_\mu, E_\lambda^*]_{\mathbf{K}_n}$ (see Lemma 2.11) and the terms of the right-hand side by $[E_\lambda^* : E_\mu^* \otimes E_\xi^*]_{\mathbf{G}_n}$. Using Remark 2.8 we can write the highest weight of V_μ explicitly, and then we can determine the partition that corresponds to V_μ in the parametrization of [4, Thm 10.2.9] and [4, Thm 10.2.12]. If $[E_\lambda : E_\mu \otimes M_\xi]_{\mathbf{G}_n} \neq 0$ for some $\xi \in \mathcal{P}_n$, then Remark 2.10 implies that $\ell(\xi) \leq r$ when $\mathbb{F} = \mathbb{R}$, and $\ell(\xi) \leq 2r$ when $\mathbb{F} = \mathbb{H}$. By putting all of these facts together, we can verify that the statement of the proposition is a special case of the Littlewood Restriction theorem. \square

3. The quadratic Capelli operators

In this section we define the differential operators $D_{\lambda,s}$. Let $\mathcal{D}(A) := \bigoplus_{m=0}^\infty \mathcal{D}^m(A)$ denote the \mathbb{C} -algebra of constant coefficient differential operators on A , endowed with the usual \mathbb{Z} -grading. We define $\mathcal{D}(W) := \bigoplus_{m=0}^\infty \mathcal{D}^m(W)$ similarly. There are natural G_r -actions

on $\mathcal{D}(A)$ and $\mathcal{D}(W)$, and as in Section 2, these actions extend uniquely to holomorphic \mathbf{G}_r -actions on the same vector spaces. Furthermore, the canonical isomorphisms

$$(9) \quad \mathcal{D}^m(A) \simeq \mathcal{P}^m(A)^* \text{ and } \mathcal{D}^m(W) \simeq \mathcal{P}^m(W)^*$$

are \mathbf{G}_r -equivariant.

Let $\mathcal{P}\mathcal{D}(A)$ and $\mathcal{P}\mathcal{D}(W)$ denote the algebras of polynomial coefficient differential operators on A and W . The multiplication map results in isomorphisms of vector spaces

$$(10) \quad \mathcal{P}(A) \otimes \mathcal{D}(A) \simeq \mathcal{P}\mathcal{D}(A) \text{ and } \mathcal{P}(W) \otimes \mathcal{D}(W) \simeq \mathcal{P}\mathcal{D}(W).$$

From now on, we set

$$\theta : G_r \rightarrow G_r, \theta(g) := (g^\dagger)^{-1}.$$

DEFINITION 3.1. – We define bilinear forms $(\cdot, \cdot)_W : W \times W \rightarrow \mathbb{R}$ and $(\cdot, \cdot)_A : A \times A \rightarrow \mathbb{R}$ by

$$(x, y)_W := \Re(\text{tr}(x^\dagger y)) \text{ and } (x, y)_A := \Re(\text{tr}(xy)).$$

The bilinear form $(\cdot, \cdot)_W$ is K_n -invariant and θ -invariant, that is,

$$(kx, ky)_W = (x, y)_W \text{ and } (g \cdot x, y)_W = (x, \theta(g)^{-1} \cdot y)_W \text{ for } x, y \in W, k \in K_n, \text{ and } g \in G_r.$$

Similarly, the bilinear form $(\cdot, \cdot)_A$ is θ -invariant, that is, $(g \cdot x, y)_A = (x, \theta(g)^{-1} \cdot y)_A$ for $x, y \in A$ and $g \in G_r$. The bilinear forms $(\cdot, \cdot)_W$ and $(\cdot, \cdot)_A$ yield canonical isomorphisms

$$(11) \quad \iota_W : W \rightarrow W^* \text{ and } \iota_A : A \rightarrow A^*.$$

These maps extend to \mathbf{G}_r -equivariant isomorphisms of \mathbb{C} -algebras

$$(12) \quad \iota_W : \mathcal{D}(W) \simeq \mathcal{P}(W) \text{ and } \iota_A : \mathcal{D}(A) \simeq \mathcal{P}(A).$$

Since ι_W is also K_n -equivariant, it restricts to an isomorphism $\mathcal{P}(W)^{K_n} \rightarrow \mathcal{D}(W)^{K_n}$. Consequently, we obtain a \mathbf{G}_r -equivariant isomorphism of \mathbb{C} -algebras

$$(\iota_W)^{-1} \circ \psi^* \circ \iota_A : \mathcal{D}(A) \rightarrow \mathcal{D}(W)^{K_n}.$$

Set $\check{\psi} := (\iota_W)^{-1} \circ \psi^* \circ \iota_A$. From (10) it follows that the map

$$(13) \quad \iota : \mathcal{P}\mathcal{D}(A) \hookrightarrow \mathcal{P}\mathcal{D}(W)^{K_n}, \iota := \check{\psi} \otimes \psi^*$$

is an embedding of \mathbf{G}_r -modules. From (7) and (9) it follows that $\mathcal{D}(A) \simeq \bigoplus_{\lambda \in \mathcal{P}_r} F_\lambda^*$, so that

$$\mathcal{P}\mathcal{D}(A) \simeq \mathcal{P}(A) \otimes \mathcal{D}(A) \simeq \bigoplus_{\lambda, \mu \in \mathcal{P}_r} F_\lambda \otimes F_\mu^* \simeq \bigoplus_{\lambda, \mu \in \mathcal{P}_r} \text{Hom}_{\mathbb{C}}(F_\mu, F_\lambda).$$

By Schur's Lemma $[F_\mu : F_\lambda]_{\mathbf{G}_r} \leq 1$, and equality occurs if and only if $\lambda = \mu$. Thus,

$$(14) \quad \mathcal{P}\mathcal{D}(A)^{\mathbf{G}_r} \simeq \bigoplus_{\lambda, \mu \in \mathcal{P}_r} \text{Hom}_{\mathbf{G}_r}(F_\mu, F_\lambda) \simeq \bigoplus_{\lambda \in \mathcal{P}_r} \mathbb{C}I_\lambda,$$

where I_λ denotes the identity element of $\text{Hom}_{\mathbb{C}}(F_\lambda, F_\lambda)$. Let $C_\lambda \in \mathcal{P}\mathcal{D}(A)^{\mathbf{G}_r}$ be the differential operator that corresponds to I_λ by the isomorphism (14). We now set

$$(15) \quad \tilde{D}_\lambda := \iota(C_\lambda) \in \mathcal{P}\mathcal{D}(W)^{K_n \times G_r},$$

where ι is the map defined in (13).

Set $\Psi(x) := \det(x^\dagger x)$ for $x \in W$ (in the case $\mathbb{F} = \mathbb{H}$ we define $\det(z) := \det(\check{z})$ for $z \in \text{Mat}_{r \times r}(\mathbb{H})$, where \check{z} is as in Remark 2.1). For every $\lambda \in \mathcal{P}_r$ and every $s \in \mathbb{C}$, let $\widetilde{D}_{\lambda,s}$ be the differential operator on X defined by

$$\widetilde{D}_{\lambda,s} := \Psi^s \widetilde{D}_\lambda \Psi^{-s}.$$

DEFINITION 3.2. – For $\lambda \in \mathcal{P}_r$ and $s \in \mathbb{C}$, we set $(D_{\lambda,s} f)(\varphi(x)) := \widetilde{D}_{\lambda,s}(f \circ \varphi)(x)$ for every $f \in C^\infty(Y)$ and every $x \in X$, where $X \xrightarrow{\varphi} Y$ is the map defined in Section 1.

Since $D_{\lambda,s}$ does not increase supports, by Peetre’s Theorem [5, Thm II.1.4] it is a differential operator on Y . From K_n -invariance of $\widetilde{D}_{\lambda,s}$ it follows that $D_{\lambda,s}$ is also K_n -invariant.

4. The polynomials $P_\lambda(x; \tau, \alpha)$

In this section we review the definition and properties of the polynomials $P_\lambda(x; \tau, \alpha)$.

DEFINITION 4.1. – Let $\lambda := (\lambda_1, \dots, \lambda_r) \in \mathbb{Z}^r$. A Laurent polynomial $f(x_1, \dots, x_r)$ in variables x_1, \dots, x_r is called λ -monic if the coefficient of $x_1^{\lambda_1} \dots x_r^{\lambda_r}$ in $f(x_1, \dots, x_r)$ is equal to 1.

Recall that the Weyl group W of type BC_r is a semidirect product $W := S_r \ltimes \{\pm 1\}^r$, where S_r denotes the symmetric group on r letters. In [19], Okounkov defined a family of Laurent polynomials

$$P_\lambda^{\text{ip}}(x; q, t, a) \in \mathbb{C}(q, t, a)[x_1^{\pm 1}, \dots, x_r^{\pm 1}],$$

parametrized by partitions $\lambda \in \mathcal{P}_r$ (we use the notation of [14, Sec. 5]). Every P_λ^{ip} is the unique λ -monic Laurent polynomial of degree $|\lambda|$ that is invariant under the action of W on the x_i ’s by permutations and inversions, and satisfies the vanishing condition

$$P_\lambda^{\text{ip}}(aq^\mu t^\delta; q, t, a) = 0 \text{ unless } \lambda \subseteq \mu,$$

where $\delta := (r - 1, \dots, 0)$ and $\mu \in \mathcal{P}_r$ (see [14, Sec. 5.3]). Here as usual we define $q^\mu t^\delta := (q^{\mu_1} t^{\delta_1}, \dots, q^{\mu_r} t^{\delta_r})$. The polynomials $P_\lambda^{\text{ip}}(x; q, t, a)$ are analogs (for the BC_r -type root system) of the q -deformed interpolation Macdonald polynomials defined by Knop [11] and Sahi [21].

By taking the $q \rightarrow 1$ limit of P_λ^{ip} (see [14, Def. 7.1]), one obtains a polynomial

$$P_\lambda(x; \tau, \alpha) \in \mathbb{C}(\tau, \alpha)[x_1, \dots, x_r].$$

More precisely,

$$P_\lambda(x; \tau, \alpha) := \lim_{q \uparrow 1} (1 - q)^{-2|\lambda|} P_\lambda^{\text{ip}}(q^x; q, q^\tau, q^\alpha),$$

where $|\lambda| := \sum_i \lambda_i$. From the symmetry property of P_λ^{ip} it follows that $P_\lambda(x; \tau, \alpha)$ is invariant under permutations and sign changes of x_1, \dots, x_r .

DEFINITION 4.2. – A polynomial in variables x_1, \dots, x_r is called *even-symmetric* if it is invariant under permutations and sign changes of the x_i ’s.

A combinatorial formula for $P_\lambda(x; \tau, \alpha)$ is given in [14, Sec. 7]. To recall this formula, we need some terminology. Every partition λ can be represented by a Young diagram consisting of boxes $b := b(i, j)$, where

$$(i, j) \in \{(p, q) \in \mathbb{Z}^2 : 1 \leq p \leq \ell(\lambda) \text{ and } 1 \leq q \leq \lambda_i\}.$$

The arm length and leg length of a box $b := b(i, j)$ in the Young diagram of λ are $a_\lambda(b) := \lambda_i - j$ and $l_\lambda(b) := |\{k > i : \lambda_k \geq j\}|$. We also set $a'_\lambda(b) := j - 1$ (arm co-length) and $l'_\lambda(b) := i - 1$ (leg co-length). By a reverse tableau of shape λ with entries in $\{1, \dots, r\}$ we mean a filling of the Young diagram that corresponds to λ , with weakly decreasing rows and strongly decreasing columns. For a reverse tableau T of shape λ and an integer $k \in \{0, \dots, r\}$, let $\lambda^{(k)} \subseteq \lambda$ be the partition corresponding to the boxes $b \in \lambda$ that satisfy $T(b) > k$. Thus for $1 \leq k \leq n$, $\lambda^{(k-1)} \setminus \lambda^{(k)}$ is the horizontal strip consisting of the boxes that contain k . Finally, for two partitions $\nu \subseteq \mu$, we define $(R \setminus C)_{\mu \setminus \nu}$ to be the set of boxes which are in a row of μ intersecting with $\mu \setminus \nu$, but not in a column of μ intersecting with $\mu \setminus \nu$. Set

$$b_\mu(b; \tau) := \frac{a_\mu(b) + \tau(l_\mu(b) + 1)}{a_\mu(b) + \tau l_\mu(b) + 1} \quad \text{and} \quad \psi_T(\tau) := \prod_{i=1}^r \prod_{b \in (R \setminus C)_{\lambda^{(i-1)} \setminus \lambda^{(i)}}} \frac{b_{\lambda^{(i)}}(b; \tau)}{b_{\lambda^{(i-1)}}(b; \tau)}.$$

Then

$$(16) \quad P_\lambda(x; \tau, \alpha) = \sum_T \psi_T(\tau) \prod_{b \in \lambda} \left(x_{T(b)}^2 - (a'_\lambda(b) + \tau(n - T(b) - l'_\lambda(b)) + \alpha)^2 \right),$$

where the sum is over all reverse tableaux T of shape λ with entries in $\{1, \dots, r\}$.

PROPOSITION 4.3. – *Fix real numbers $\alpha, \tau > 0$. For every partition $\lambda \in \mathcal{P}_r$, the polynomial*

$$Q_\lambda(x) := P_\lambda(x; \tau, \alpha) \in \mathbb{C}[x_1, \dots, x_r]$$

is the unique 2λ -monic, even-symmetric polynomial that satisfies the vanishing condition

$$(17) \quad Q_\lambda(\mu + \varrho_{\tau, \alpha}) = 0 \text{ if } |\mu| \leq |\lambda| \text{ and } \mu \neq \lambda,$$

where $\varrho_{\tau, \alpha}$ is given in (2).

Proof. – The existence statement follows from [14, Sec. 7] and the fact that the specialization of P_λ at the values of τ and α is well-defined, because when $\tau > 0$, the denominators of the coefficients $\psi_T(\tau)$ of $P_\lambda(x; \tau, \alpha)$ that appear in the combinatorial Formula (16) do not vanish. For the uniqueness statement, we use a method based on [20]. First note that from $\alpha, \tau > 0$ and [14, Eq. (7.5)] it follows that $Q_\lambda(\lambda + \varrho_{\tau, \alpha}) \neq 0$. Next fix an integer $N > 0$, set $\mathcal{J}_N := \{\mu \in \mathcal{P}_r : |\mu| \leq N\}$, and let \mathcal{S}_N denote the vector space of even-symmetric polynomials in x_1, \dots, x_r of degree at most $2N$. Note that $\dim \mathcal{S}_N = |\mathcal{J}_N|$. For every $\mu \in \mathcal{J}_N$, we consider the linear maps

$$L_\mu : \mathcal{S}_N \rightarrow \mathbb{C}, \quad f \mapsto f(\mu + \varrho_{\tau, \alpha}).$$

Next we define a total order $<$ on \mathcal{J}_N , as follows. We set $\mu < \nu$ for every $\mu, \nu \in \mathcal{J}_N$ which satisfy $|\mu| < |\nu|$, and then we extend the resulting partial order to a total order on \mathcal{J}_N . Then the matrix

$$[L_\mu(Q_{\mu'})]_{\mu, \mu' \in \mathcal{J}_N}$$

is upper triangular with nonzero entries on the diagonal. It follows that the linear map

$$\mathcal{S}_N \rightarrow \mathbb{C}^{\dim \mathcal{S}_N}, f \mapsto [f(\mu + \varrho_{\tau, \alpha})]_{\mu \in \mathcal{J}_N}$$

is invertible. Uniqueness of Q_λ follows immediately from the latter statement. □

Recall that $d := \dim(\mathbb{F})$. For every $\lambda \in \mathcal{P}_r$, we set

$$(18) \quad \gamma_\lambda := \frac{(-2d)^{|\lambda|}}{\prod_{b \in \lambda} \left(\frac{d}{2}(a_\lambda(b) + 1) + l_\lambda(b) \right)}.$$

This is the scalar that appears in the statement of Theorem 1.2.

5. The vanishing property of $c_{\lambda,s}(\mu)$

In this section, we prove a few technical statements which will be used in the proof of Theorem 1.2. The ultimate goal of this section is to prove Proposition 5.6. The proof of Theorem 1.2 will be completed in Section 6. Recall that $\varrho := (\varrho_1, \dots, \varrho_r)$ is the vector defined in (3).

LEMMA 5.1. – *Let $\lambda \in \mathcal{P}_r$. Then there exists a polynomial $d_\lambda(x, s)$ which is even-symmetric in $x := (x_1, \dots, x_r)$, has x -degree and s -degree at most $2|\lambda|$, and satisfies*

$$c_{\lambda,s}(\mu) = d_\lambda(\mu + \varrho, s),$$

for every $s \in \mathbb{C}$ and every $\mu \in \mathcal{P}_r$.

Proof. – The proof is similar to [25, Lemma 3.1]. Throughout the proof we fix λ . Recall that $K := K_n$ and $M := K_r \times K_{n-r}$. For every $s \in \mathbb{C}$, the differential operator $D_{\lambda,s}$ is K -invariant and has order at most $2|\lambda|$. By (1), for every $\mu \in \mathcal{P}_r$, the highest weight $\tilde{\mu}$ of V_μ satisfies $\tilde{\mu}|_{\mathfrak{ac}} = \sum_{i=1}^r 2\mu_i e_i$. Therefore from the Harish-Chandra homomorphism [5, Chap. II] it follows that for every $s \in \mathbb{C}$, the scalar $c_{\lambda,s}(\mu)$ is an even-symmetric polynomial of degree at most $2|\lambda|$ evaluated at $\mu + \varrho$. Next we set

$$x_\circ := \begin{bmatrix} I_{r \times r} \\ 0 \end{bmatrix} \in X,$$

and we denote the image of x_\circ in Y by y_\circ . Choose an M -fixed vector $h_\mu \in V_\mu \subseteq C^\infty(Y)$ such that $h_\mu(y_\circ) = 1$. By evaluating both sides of the relation $D_{\lambda,s}h_\mu = c_{\lambda,s}(\mu)h_\mu$ at y_\circ , we obtain

$$(19) \quad c_{\lambda,s}(\mu) = \tilde{D}_\lambda(\det(x^\dagger x)^{-s}(h_\mu \circ \varphi))(x_\circ),$$

where \tilde{D}_λ is defined in (15). Since \tilde{D}_λ is a polynomial coefficient differential operator of order at most $2|\lambda|$, from the Leibniz rule it follows that for fixed μ , the right-hand side of (19) is a polynomial in s of degree at most $2|\lambda|$. Consequently,

$$c_{\lambda,s}(\mu) = \sum_{j=0}^{2|\lambda|} a_j(\mu)s^j \text{ for every } s \in \mathbb{C}.$$

For $2|\lambda| + 1$ distinct values of s , we obtain a linear system in the coefficients $a_j(\mu)$ whose coefficients form an invertible Vandermonde matrix. Since for fixed $s \in \mathbb{C}$ we have shown that $c_{s,\lambda}(\mu)$ is a polynomial in μ of degree at most $2|\lambda|$, it follows that $a_j(\mu)$ is also a polynomial

in μ of degree at most $2|\lambda|$. Consequently, both the x -degree and the s -degree of $c_{s,\lambda}$ are at most $2|\lambda|$. The statement that d_λ is even-symmetric follows from the fact that $c_{\lambda,s}(\mu)$ is even-symmetric as a polynomial in $\mu + \varrho$. \square

DEFINITION 5.2. – For every $\chi \in \mathcal{P}_r$, let $\mathcal{P}_\chi(W)$ denote the F_χ -isotypic component of $\mathcal{P}(W)$.

We remark that $\mathcal{P}_\chi(W)$ is \mathbf{G}_n -invariant.

PROPOSITION 5.3. – Let $\lambda, \nu \in \mathcal{P}_r$, and let $m \in \mathbb{Z}$ such that $m \geq \max\{\lambda_1, \nu_1\}$. Set

$$\eta := (m - \nu_r, \dots, m - \nu_1) \text{ and } \chi := (m - \lambda_r, \dots, m - \lambda_1).$$

If $[V_\eta : \mathcal{P}_\chi(W)]_{\mathbf{K}_n} > 0$, then $\lambda \subseteq \nu$.

Proof. – Recall from Section 2 that by $(\mathrm{GL}_n, \mathrm{GL}_r)$ -duality, $\mathcal{P}_\chi(W) \simeq E_\chi^{\oplus \dim F_\chi}$ as \mathbf{G}_n -modules. Therefore we can assume that $[V_\eta : E_\chi]_{\mathbf{K}_n} > 0$. We will consider two separate cases.

Case I. – $\mathbb{F} = \mathbb{R}, \mathbb{H}$. Proposition 2.12 implies that $[E_\chi : E_\eta \otimes E_\xi]_{\mathbf{G}_n} > 0$ for some $\xi \in \mathcal{P}_r$, and Remark 2.10 implies that $\eta \subseteq \chi$, hence $\lambda \subseteq \nu$.

Case II. – $\mathbb{F} = \mathbb{C}$. Let $\tilde{\chi} \in \mathcal{P}_n$ be obtained from χ by adding $n - r$ zeros on the right of the parts of χ . Our assumption entails that the restriction of the $\mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_n(\mathbb{C})$ -module $M_{\tilde{\chi}} \otimes M_{\tilde{\chi}}^*$ to the diagonal subgroup $\mathrm{GL}_n(\mathbb{C})$ contains the $\mathrm{GL}_n(\mathbb{C})$ -module $M_{\tilde{\eta}}$, where

$$\tilde{\eta} := (m - \nu_r, \dots, m - \nu_1, 0, \dots, 0, -m + \nu_1, \dots, -m + \nu_r) \in \mathcal{P}_n.$$

Remark 2.10 implies that $-m + \nu_i \geq -m + \lambda_i$ for every $1 \leq i \leq r$, hence $\lambda \subseteq \nu$. \square

The proof of our next result, Proposition 5.4, is based on facts from the theory of symmetric functions, which we now review quickly (for a comprehensive reference, see [18]). Let

$$\mathcal{P} := \varinjlim_k \mathcal{P}_k$$

be the set of all partitions, where the maps $\mathcal{P}_k \rightarrow \mathcal{P}_{k+1}$ are given by $(\lambda_1, \dots, \lambda_k) \mapsto (\lambda_1, \dots, \lambda_k, 0)$, and let

$$\Lambda := \varprojlim_k \mathbb{C}[x_1, \dots, x_k]^{S_k}$$

denote the ring of symmetric functions, where the maps $\mathbb{C}[x_1, \dots, x_{k+1}] \rightarrow \mathbb{C}[x_1, \dots, x_k]$ are given by $f \mapsto f(x_1, \dots, x_k, 0)$. As usual, we equip Λ with a scalar product defined by $\langle h_\lambda, m_\mu \rangle_\Lambda := \delta_{\lambda,\mu}$, where h_λ and m_μ are the complete and monomial symmetric functions associated to $\lambda, \mu \in \mathcal{P}$. The Schur functions $s_\lambda, \lambda \in \mathcal{P}$, form an orthonormal basis for Λ . For every two $\lambda, \mu \in \mathcal{P}$ such that $\mu \subseteq \lambda$, the skew Schur function $s_{\lambda \setminus \mu} \in \Lambda$ satisfies the relation $\langle s_{\lambda \setminus \mu}, s_\nu \rangle_\Lambda = \langle s_\lambda, s_\mu s_\nu \rangle_\Lambda$ for every $\nu \in \mathcal{P}$. It is well known [29, Ex. 7.56(a)] that for any skew diagram $\lambda \setminus \mu$ we have

$$(20) \quad s_{\lambda \setminus \mu} = s_{(\lambda \setminus \mu)^\circ},$$

where $(\lambda \setminus \mu)^\circ$ denotes the skew diagram obtained by a 180 degree rotation of $\lambda \setminus \mu$.

PROPOSITION 5.4. – Let $m \geq 1$ be an integer, let $\mathbf{m} := (m, \dots, m) \in \mathcal{P}_r$ be the partition corresponding to the $r \times m$ rectangular Young diagram, and set $\mathcal{P}(\mathbf{m}) := \{\lambda \in \mathcal{P}_r : \lambda \subseteq \mathbf{m}\}$. Let $\mathcal{P}_{\mathbf{m}}(W)$ be as in Definition 5.2. Then as \mathbf{K}_n -modules,

$$\mathcal{P}_{\mathbf{m}}(W) \simeq \bigoplus_{\mu \in \mathcal{P}(\mathbf{m})} V_{\mu}.$$

Proof. – Recall that $\mathbf{K} := \mathbf{K}_n$ is the complexification of $K := K_n$, and that \mathbf{M} is the complexification of M . The map $\mathcal{P}_{\mathbf{m}}(W) \rightarrow C^\infty(X)^{G_r} \simeq C^\infty(Y)$, $f \mapsto \Psi^{-m} f$, is a K -equivariant embedding. It follows that $\mathcal{P}_{\mathbf{m}}(W)$ is a direct sum of irreducible \mathbf{M} -spherical \mathbf{K} -modules. Next we determine the multiplicity of every \mathbf{K} -module V_{λ} in $\mathcal{P}_{\mathbf{m}}(W)$. As in Section 2, $(\mathrm{GL}_n, \mathrm{GL}_r)$ -duality entails that $\mathcal{P}_{\mathbf{m}}(W) \simeq E_{\mathbf{m}}$ as \mathbf{G}_n -modules. We consider two separate cases.

Case I. – $\mathbb{F} = \mathbb{R}, \mathbb{H}$. For every $\eta := (\eta_1, \eta_2, \dots, \eta_r) \in \mathcal{P}_r$, we define $\eta^\bullet \in \mathcal{P}$ as in Section 2. By Proposition 2.12,

$$(21) \quad [V_{\lambda} : E_{\mathbf{m}}]_{\mathbf{K}_n} = \sum_{\xi \in \mathcal{P}_r} [E_{\mathbf{m}} : E_{\lambda} \otimes E_{\xi}]_{\mathbf{G}_n}.$$

If $[E_{\mathbf{m}} : E_{\lambda} \otimes E_{\xi}]_{\mathbf{G}_n} \neq 0$ for some $\xi \in \mathcal{P}_r$, then $[E_{\mathbf{m}}^* : E_{\lambda}^* \otimes E_{\xi}^*]_{\mathbf{G}_n} \neq 0$ and hence $\lambda, \xi \subseteq \mathbf{m}$ (see Remark 2.10). Therefore

$$[E_{\mathbf{m}} : E_{\lambda} \otimes E_{\xi}]_{\mathbf{G}_n} = \langle s_{\mathbf{m}^\bullet}, s_{\lambda^\bullet} s_{\xi^\bullet} \rangle_{\Lambda} = \langle s_{\mathbf{m}^\bullet \setminus \xi^\bullet}, s_{\lambda^\bullet} \rangle_{\Lambda} = \langle s_{(\mathbf{m}^\bullet \setminus \xi^\bullet)^\circ}, s_{\lambda^\bullet} \rangle_{\Lambda} = \delta_{(\mathbf{m} \setminus \xi)^\circ, \lambda}.$$

It follows that the value of (21) is 0 or 1, with the latter occurring exactly when $\lambda \subseteq \mathbf{m}$.

Case II. – $\mathbb{F} = \mathbb{C}$. Let $\alpha, \beta \in \mathcal{P}_n$ be defined by

$$\alpha := (\underbrace{m, \dots, m}_{r \text{ times}}, 0, \dots, 0) \text{ and } \beta := (\underbrace{m, \dots, m}_{n-r \text{ times}}, 0, \dots, 0).$$

The statement of the proposition is equivalent (after twisting $E_{\mathbf{m}}$ by $\det(\cdot)^m \otimes 1$) to showing that the restriction of the $\mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_n(\mathbb{C})$ -module $M_{\alpha} \otimes M_{\beta}$ to the diagonal $\mathrm{GL}_n(\mathbb{C})$ is a multiplicity-free direct sum of $\mathrm{GL}_n(\mathbb{C})$ -modules M_{η} for η of the form

$$(22) \quad (m + \xi_1, \dots, m + \xi_r, \underbrace{m, \dots, m}_{n-2r \text{ times}}, m - \xi_r, \dots, m - \xi_1),$$

where $\xi := (\xi_1, \dots, \xi_r)$ varies through partitions satisfying $\xi_1 \leq m$. Assume that $M_{\eta} \subseteq M_{\alpha} \otimes M_{\beta}$. From Remark 2.10 it follows that $\beta \subseteq \eta$, and in addition, every column of the skew Young diagram corresponding to $\eta \setminus \beta$ has height at most r . Since $r \leq n - r$, it follows that $\eta_i \leq m$ for every $i > n - r$. Consequently, the skew Young diagram corresponding to $\eta \setminus \beta$ is a disjoint union of two (non-skew) diagrams corresponding to the partitions

$$\eta^+ := (\eta_1 - m, \dots, \eta_r - m) \text{ and } \eta^- := (\eta_{n-r+1}, \dots, \eta_n).$$

From [18, Sec. I.5.7] it follows that $s_{\eta} = s_{\eta^+} s_{\eta^-}$, hence

$$\begin{aligned} [M_{\eta} : M_{\alpha} \otimes M_{\beta}]_{\mathrm{GL}_n(\mathbb{C})} &= \langle s_{\eta}, s_{\alpha} s_{\beta} \rangle_{\Lambda} = \langle s_{\eta \setminus \beta}, s_{\alpha} \rangle_{\Lambda} \\ &= \langle s_{\eta^+} s_{\eta^-}, s_{\alpha} \rangle_{\Lambda} = \langle s_{\eta^+}, s_{\alpha \setminus \eta^-} \rangle_{\Lambda} = \langle s_{\eta^+}, s_{(\alpha \setminus \eta^-)^\circ} \rangle_{\Lambda} = \delta_{\eta^+, (\alpha \setminus \eta^-)^\circ}. \end{aligned}$$

It is now straightforward to verify that $[M_{\eta} : M_{\alpha} \otimes M_{\beta}]_{\mathrm{GL}_n(\mathbb{C})} \leq 1$, with equality occurring if and only if η is of the form given in (22). \square

REMARK 5.5. – Recall that $\varrho := (\varrho_1, \dots, \varrho_r)$ is the vector defined in (3). Let $\varrho' \in \mathbb{C}^r$ denote the vector $\varrho_{\tau, \alpha}$ defined in (2), for $\tau := \frac{d}{2}$ and $\alpha := s - \varrho_1$. Then the vectors ϱ and ϱ' are related by the relation

$$\varrho_{r-i+1} + \varrho'_i = s \text{ for } 1 \leq i \leq r.$$

PROPOSITION 5.6. – Let $\lambda \in \mathcal{P}_r$, and let $d_\lambda(x, s)$ be as in Lemma 5.1. Let $\varrho_{\tau, \alpha}$ be as in (2) for $\tau := \frac{d}{2}$ and $\alpha := s - \varrho_1$. For every $v \in \mathcal{P}_r$, if $\lambda \not\subseteq v$ then

$$(23) \quad d_\lambda(v + \varrho_{\tau, \alpha}, s) = 0 \text{ for every } s \in \mathbb{C}.$$

Proof. – Fix $v \in \mathcal{P}_r$. Since $d_\lambda(x, s)$ is a polynomial in $x = (x_1, \dots, x_r)$ and s , it is enough to prove (23) for infinitely many values of s . Since d_λ is even-symmetric, vanishing of $d_\lambda(x, s)$ at a point (x_1, \dots, x_r) is equivalent to its vanishing at the point $(-x_r, \dots, -x_1)$. Therefore it suffices to show that $d_\lambda(y, s) = 0$ for $y := (y_1, \dots, y_r)$ given by

$$y_i := -v_{r+1-i} - \frac{d}{2}(i-1) - s + \varrho_1 \text{ for every } 1 \leq i \leq r.$$

Next set $s := -m$ for some integer $m \geq \lambda_1$. By Remark 5.5 and Lemma 5.1, it suffices to show that

$$(24) \quad c_{\lambda, -m}(\eta) = 0 \text{ for } \eta := (m - v_r, \dots, m - v_1).$$

Recall that $\mathbf{m} := (m, \dots, m) \in \mathcal{P}_r$ is the partition corresponding to an $r \times m$ Young diagram. Set

$$(25) \quad \mathcal{R}_{\mathbf{m}}(W) := \{\Psi^{-m} f : f \in \mathcal{P}_{\mathbf{m}}(W)\},$$

where Ψ is defined in Section 3. The map

$$j_m : \mathcal{R}_{\mathbf{m}}(W) \rightarrow \mathcal{P}_{\mathbf{m}}(W), f \mapsto \Psi^m f$$

is an isomorphism of \mathbf{K}_n -modules. Therefore Proposition 5.4 implies that V_η occurs as a \mathbf{K}_n -submodule of $\mathcal{R}_{\mathbf{m}}(W)$. By restriction to X , and then factoring G_r -invariant functions on X through $Y \simeq X/G_r$, we obtain an embedding $\mathcal{R}_{\mathbf{m}}(W) \subset C^\infty(Y)$. From K_n -invariance of $D_{\lambda, -m}$ it follows that $D_{\lambda, -m}\mathcal{R}_{\mathbf{m}} \subseteq \mathcal{R}_{\mathbf{m}}$. Furthermore, the diagram

$$\begin{array}{ccc} \mathcal{R}_{\mathbf{m}}(W) & \xrightarrow{D_{\lambda, -m}} & \mathcal{R}_{\mathbf{m}}(W) \\ j_m \downarrow & & \downarrow j_m \\ \mathcal{P}_{\mathbf{m}}(W) & \xrightarrow{D_\lambda} & \mathcal{P}_{\mathbf{m}}(W) \end{array}$$

commutes. From the latter commutative diagram it follows that, in order to prove (24), it suffices to show that $(D_\lambda \circ j_m)V_\eta = \{0\}$. Note that $D_\lambda \in \iota(F_\lambda \otimes F_\lambda^*)$, where ι is the map defined in (13). The elements of $\iota(F_\lambda^*)$ act on $\mathcal{P}(W)$ as K_n -invariant constant coefficient differential operators, and by considering the G_r -action it follows that they map $\mathcal{P}_{\mathbf{m}}(W)$ into $\mathcal{P}_\chi(W)$, where

$$\chi = (m - \lambda_r, \dots, m - \lambda_1).$$

Therefore the map $D_\lambda : \mathcal{P}_{\mathbf{m}}(W) \rightarrow \mathcal{P}_{\mathbf{m}}(W)$ factors through a K_n -equivariant map

$$(26) \quad \mathcal{P}_{\mathbf{m}}(W) \rightarrow \mathcal{P}_\chi(W).$$

Suppose that $(D_\lambda \circ j_m)V_\eta \neq \{0\}$. Since the map (26) is K_n -equivariant, $[V_\eta : \mathcal{P}_\chi(W)]_{\mathbf{K}_n} > 0$. From Proposition 5.3 it follows that $\lambda \subseteq \nu$, which is a contradiction. \square

6. Proof of Theorem 1.2

In this section we complete the proof of Theorem 1.2. We start by proving the following lemma.

LEMMA 6.1. – *Let $\lambda \in \mathcal{P}_r$. Then there exists a constant $\gamma'_\lambda \in \mathbb{C}$ such that for every $s \in \mathbb{C}$, and every $\mu \in \mathcal{P}_r$,*

$$(27) \quad c_{\lambda,s}(\mu) = \gamma'_\lambda P_\lambda \left(\mu + \varrho; \frac{d}{2}, s - \varrho_1 \right).$$

Proof. – First fix $s > \frac{1}{2}\varrho_1$. From Proposition 5.6 and Lemma 5.1 we obtain

$$c_{\lambda,s} \left(\nu + \left(\frac{d}{2}(r-1) + s - 2\varrho_1 \right) \mathbb{1} \right) = 0 \text{ if } |\nu| \leq |\lambda| \text{ and } \nu \neq \lambda.$$

From the vanishing part of the statement of Proposition 4.3 it follows that the polynomial $q_\lambda(x) := P_\lambda \left(x + \varrho; \frac{d}{2}, s - \varrho_1 \right)$ also vanishes for all $x := \nu + \left(\frac{d}{2}(r-1) + s - 2\varrho_1 \right) \mathbb{1}$, where ν satisfies $|\nu| \leq |\lambda|$ and $\nu \neq \lambda$. The uniqueness part of the statement of Proposition 4.3 now implies that there exists a scalar $\gamma'_\lambda \in \mathbb{C}$, possibly depending on s , such that (27) holds for every μ . Since both sides of (27) are polynomials in s and μ (see Lemma 5.1), and (27) holds for a Zariski dense subset of values $(\mu, s) \in \mathbb{C}^r \times \mathbb{C}$, it follows that (27) indeed holds for every $s \in \mathbb{C}$, and γ'_λ is a rational function of $s \in \mathbb{C}$.

Next we show that γ'_λ does not depend on s . Since the value of $c_{\lambda,s}(\mu)$ is a polynomial in s and μ , and $P_\lambda(\mu + \varrho; \frac{d}{2}, s - \varrho_1)$ is 2λ -monic in μ , it follows that γ'_λ is a polynomial in s . However, from the combinatorial formula for P_λ that is given in (16), it follows that the degree of s in $P_\lambda \left(\mu + \varrho; \frac{d}{2}, s - \varrho_1 \right)$ is exactly $2|\lambda|$, whereas the degree of s in $c_{\lambda,s}(\mu)$ is at most $2|\lambda|$. By comparing the degrees of s on both sides of (27), it follows that γ'_λ is a constant independent of s . \square

To complete the proof of Theorem 1.2, we need to prove that $\gamma'_\lambda = \gamma_\lambda$, where γ_λ is defined in (18). The rest of this section is devoted to the proof of the latter claim.

Set $t := \dim_{\mathbb{R}}(A)$ and let v_1, \dots, v_t be an orthonormal basis for A with respect to the pairing $(\cdot, \cdot)_A$. Set $\varphi_i := (\cdot, v_i)_A \in A^*$ for every $1 \leq i \leq t$, so that $\varphi_i(v_j) = \delta_{i,j}$. For every $1 \leq i \leq t$, let $\partial_{v_i} \in \mathcal{D}(A)$ denote the directional derivative corresponding to v_i , so that $\partial_{v_i}(\varphi_j) = \varphi_j(v_i) = \delta_{i,j}$. For every $1 \leq i \leq t$, set $q_i := \psi^*(\varphi_i)$, where ψ^* is as in (6), and let $\partial_{q_i} \in \mathcal{D}(W)$ denote the second-order differential operator corresponding to q_i under the isomorphism $\mathcal{P}(W) \simeq \mathcal{D}(W)$ defined in (12).

Fix an integer $m \geq 1$ and set

$$(28) \quad D^{(m)} := \sum_{|\lambda|=m} \tilde{D}_\lambda,$$

where \tilde{D}_λ is defined in (15). Then $D^{(m)}$ acts on every K -module $V_\mu \subseteq C^\infty(Y) \simeq C^\infty(X)^{G_r}$ by the scalar

$$c^{(m)}(\mu) := \sum_{|\lambda|=m} c_{\lambda,0}(\mu).$$

From the definition of the operators \widetilde{D}_λ it follows that if $\{\widetilde{v}_j\}$ is a basis for $\mathscr{P}^m(A)$ and $\{\widetilde{\partial}_j\}$ is the corresponding dual basis for $\mathscr{P}^m(A) \simeq \mathscr{P}^m(A)^*$, then $D^{(m)} = \sum_j \iota(\widetilde{v}_j)\iota(\widetilde{\partial}_j)$, where ι is as in (13). In particular, if we choose the basis

$$\{(\partial_{v_1})^{m_1} \cdots (\partial_{v_t})^{m_t} : m_1 + \cdots + m_t = m\}$$

for $\mathscr{P}^m(A)$, then the corresponding dual basis for $\mathscr{P}^m(A)$ will be

$$\left\{ \frac{1}{m_1! \cdots m_t!} \varphi_1^{m_1} \cdots \varphi_t^{m_t} : m_1 + \cdots + m_t = m \right\},$$

and thus

$$\begin{aligned} D^{(m)} &= \frac{1}{m!} \sum_{m_1 + \cdots + m_t = m} \binom{m}{m_1, \dots, m_t} q_1^{m_1} \cdots q_t^{m_t} (\partial_{q_1})^{m_1} \cdots (\partial_{q_t})^{m_t} \\ &= \frac{1}{m!} (q_1 \partial_{q_1} + \cdots + q_t \partial_{q_t})^m + D' = \frac{1}{m!} (D^{(1)})^m + D', \end{aligned}$$

where D' is a differential operator with order strictly less than the order of D .

Let $\mathfrak{g}_{\mathbb{C}}$ be the Lie algebra of G_n , and let $\Omega_{\mathfrak{k}}$ and $\Omega_{\mathfrak{g}}$ denote the Casimir operators of $\mathfrak{k}_{\mathbb{C}}$ and $\mathfrak{g}_{\mathbb{C}}$ (see the appendix for more information). Also, Let \mathbf{E} be the degree (or Euler) operator on W . The operator $D^{(1)}$ is a polynomial-coefficient differential operator on $W_{\mathbb{C}}$. The next proposition expresses $D^{(1)}$ in terms of $\Omega_{\mathfrak{k}}$, $\Omega_{\mathfrak{g}}$, and \mathbf{E} .

PROPOSITION 6.2. – *Let $D^{(1)}$ be defined as in (28). Then*

$$(29) \quad D^{(1)} = \begin{cases} -2(n-2)\Omega_{\mathfrak{k}} + \Omega_{\mathfrak{g}} - \mathbf{E} & \text{if } \mathbb{F} = \mathbb{R}, \\ -2\Omega_{\mathfrak{k}} + 2\Omega_{\mathfrak{g}} + (2n-2r)\mathbf{E} & \text{if } \mathbb{F} = \mathbb{C}, \\ -8(2n+1)\Omega_{\mathfrak{k}} + 2\Omega_{\mathfrak{g}} + 2(2n-2r+1)\mathbf{E} & \text{if } \mathbb{F} = \mathbb{H}. \end{cases}$$

Proof. – The proof is by a tedious calculation and is deferred to the appendix. □

LEMMA 6.3. – *The scalar $c^{(1)}(\mu)$ is a quadratic polynomial in μ_1, \dots, μ_r with top-degree homogeneous part equal to $-4(\mu_1^2 + \cdots + \mu_r^2)$.*

Proof. – We use the expressions for $D^{(1)}$ given in Proposition 6.2. As is well known (e.g., see [4, Lem. 3.3.8]), the action of $\Omega_{\mathfrak{k}}$ on V_μ is by a scalar which is a quadratic polynomial in μ_1, \dots, μ_n , whose top homogeneous term is given by

$$(30) \quad \begin{cases} \frac{2}{n-2}(\mu_1^2 + \cdots + \mu_r^2) & \text{if } \mathbb{F} = \mathbb{R}, \\ 2(\mu_1^2 + \cdots + \mu_r^2) & \text{if } \mathbb{F} = \mathbb{C}, \\ \frac{1}{4n+2}(\mu_1^2 + \cdots + \mu_r^2) & \text{if } \mathbb{F} = \mathbb{H}. \end{cases}$$

Fix $m \in \mathbb{Z}$ such that $m \geq \mu_1$, and let $\mathscr{R}_m(W)$ be as in (25). From Proposition 5.4 it follows that $[V_\mu : \mathscr{R}_m(W)]_{K_n} > 0$. Since elements of $\mathscr{R}_m(W)$ are homogeneous of degree zero, the degree operator \mathbf{E} vanishes on V_μ . Furthermore, we have $V_\mu \subseteq C^\infty(Y)_{K_n\text{-finite}}$, and $Y \simeq G_n/P_{r,n}$ where $P_{r,n}$ is the $(r, n-r)$ parabolic subgroup of G_n . Thus, $C^\infty(G_n/P_{r,n})_{K_n\text{-finite}}$ is the space of K_n -finite vectors of a degenerate principal series representation of G_n induced from $P_{r,n}$, and the operator $\Omega_{\mathfrak{g}}$ acts on the latter space by a scalar that is independent of μ (see [10, Prop. 8.22]). Consequently, the top-degree homogeneous part of $c^{(1)}(\mu)$ is determined by the action of $\Omega_{\mathfrak{k}}$. The lemma now follows from (29) and (30). □

Lemma 6.3 implies that the action of $D^{(m)}$ on V_μ is by a polynomial in μ_1, \dots, μ_r of degree $2m$, whose top-degree homogeneous part is $\frac{(-4)^m}{m!}(\mu_1^2 + \dots + \mu_r^2)^m$. On the other hand, from [14, Eq. (7.3)] it follows that for every λ such that $|\lambda| = m$, the top-degree homogeneous part of $c_{\lambda,0}(\mu)$ is up to a scalar equal to $P_\lambda(\mu^2, \frac{d}{2})$, where $P_\lambda(x, \tau)$ is the λ -monic Jack polynomial and $\mu^2 := (\mu_1^2, \dots, \mu_r^2)$. Let J_λ denote the normalization of the Jack polynomial introduced in [28, Thm 1.1]. The scalar relating J_λ and P_λ is given in [18, Chap. VI, Eq. (10.22)] (see also [28, Thm 5.6]). From [28, Prop. 2.3], and the relation between P_λ and J_λ , it follows that

$$(31) \quad (\mu_1^2 + \dots + \mu_r^2)^m = \sum_{|\lambda|=m} \left(\frac{d}{2}\right)^m \frac{m!}{\prod_{b \in \lambda} \left(\frac{d}{2}(a_\lambda(b) + 1) + l_\lambda(b)\right)} P_\lambda\left(\mu^2, \frac{d}{2}\right).$$

Since the polynomials $P_\lambda(\mu^2, \frac{d}{2})$ are linearly independent, by considering the top-degree homogeneous parts of both sides of (31) it follows that for every $\lambda \in \mathcal{P}_r$ such that $|\lambda| = m$,

$$c_{\lambda,0}(\mu) = \frac{(-2d)^m}{\prod_{b \in \lambda} \left(\frac{d}{2}(a_\lambda(b) + 1) + l_\lambda(b)\right)} P_\lambda\left(\mu + \varrho; \frac{d}{2}, -\varrho_1\right).$$

Lemma 6.1 completes the proof of Theorem 1.2.

Appendix: Proof of Proposition 6.2

In this appendix, we exhibit the details of the calculations that yield the Formulas (29) for the operator $D^{(1)}$.

Let $\kappa : \mathfrak{k}_\mathbb{C} \times \mathfrak{k}_\mathbb{C} \rightarrow \mathbb{C}$ denote the invariant bilinear form which is equal to the Killing form of $\mathfrak{k}_\mathbb{C}$ when $\mathbb{F} = \mathbb{R}$ or \mathbb{H} , and is given by $\kappa(x, y) := \text{tr}(xy)$ when $\mathbb{F} = \mathbb{C}$. The Casimir operator of $\mathfrak{k}_\mathbb{C}$ is $\Omega_{\mathfrak{k}} := \sum_{i=1}^{\dim \mathfrak{k}_\mathbb{C}} x_i x^i$, where $\{x_i : 1 \leq i \leq \dim \mathfrak{k}_\mathbb{C}\}$ is a basis for $\mathfrak{k}_\mathbb{C}$, and $\{x^i : 1 \leq i \leq \dim \mathfrak{k}_\mathbb{C}\}$ is the corresponding dual basis with respect to $\kappa(\cdot, \cdot)$. We define the Casimir operator $\Omega_{\mathfrak{g}}$ of $\mathfrak{g}_\mathbb{C}$ similarly. Explicit formulas for $\Omega_{\mathfrak{g}}$ are given in (33), (36), and (38). In the following, $E_{i,j}$ will always denote a matrix with a 1 in the (i, j) position and 0's elsewhere (the number of rows and columns of $E_{i,j}$ will be clear from the context).

Case I. – $\mathbb{F} = \mathbb{R}$. Recall that in this case $A = \text{Sym}_{r \times r}(\mathbb{R})$. The orthonormal basis of A with respect to $(\cdot, \cdot)_A$ is

$$E_{i,i} \text{ for } 1 \leq i \leq r \text{ and } \frac{1}{\sqrt{2}}(E_{i,j} + E_{j,i}) \text{ for } 1 \leq i < j \leq r.$$

We fix generators $x_{i,j} \in \mathcal{P}(A)$, where $1 \leq i \leq j \leq r$, and $y_{i,j} \in \mathcal{P}(W)$, where $1 \leq i \leq n$ and $1 \leq j \leq r$, such that $x_{i,j}([t_{a,b}]) := t_{i,j}$ for every matrix $[t_{a,b}] \in A$, and $y_{i,j}([t_{a,b}]) := t_{i,j}$ for every matrix $[t_{a,b}] \in W$. The isomorphism $\iota_A : A \rightarrow A^* \simeq \mathcal{P}^1(A)$ that is defined in (11) is given by

$$E_{i,i} \mapsto x_{i,i} \text{ for } 1 \leq i \leq r \text{ and } \frac{1}{\sqrt{2}}(E_{i,j} + E_{j,i}) \mapsto \sqrt{2}x_{i,j} \text{ for } 1 \leq i < j \leq r.$$

The map $\psi^* : \mathcal{P}(A) \rightarrow \mathcal{P}(W)$ of (6) is given by

$$x_{a,a} \mapsto \sum_{i=1}^n y_{i,a}^2 \text{ for } 1 \leq a \leq r \text{ and } \sqrt{2}x_{a,b} \mapsto \sqrt{2} \sum_{i=1}^r y_{i,a}y_{i,b} \text{ for } 1 \leq a < b \leq r.$$

Finally, the isomorphism $\iota_W : W \simeq W^* \simeq \mathcal{D}^1(W)$ of (11) is given by $y_{i,a} \mapsto \partial_{i,a} := \frac{\partial}{\partial y_{i,a}}$. From all of the above, it follows that

$$(32) \quad D^{(1)} = \sum_{a=1}^r \left(\sum_{i=1}^n y_{i,a}^2 \right) \left(\sum_{j=1}^n \partial_{j,a}^2 \right) + \sum_{1 \leq a \neq b \leq r} \left(\sum_{i=1}^n y_{i,a} y_{i,b} \right) \left(\sum_{j=1}^n \partial_{j,a} \partial_{j,b} \right).$$

The embedding $\mathfrak{k}_{\mathbb{C}} \hookrightarrow \mathfrak{gl}_n(\mathbb{C})$ gives the realization of $\mathfrak{k}_{\mathbb{C}}$ as

$$\mathfrak{k}_{\mathbb{C}} = \{x \in \text{Mat}_{n \times n}(\mathbb{C}) : x + x^T = 0\},$$

where x^T is the transpose of x . Recall that by the definition of $\Omega_{\mathfrak{g}}$,

$$(33) \quad \Omega_{\mathfrak{g}} = \sum_{1 \leq i, j \leq n} E_{i,j} E_{j,i}.$$

The Killing form of $\mathfrak{k}_{\mathbb{C}}$ is $\kappa(x, y) := (n - 2)\text{tr}(xy)$, and therefore

$$(34) \quad \Omega_{\mathfrak{k}} = -\frac{1}{2(n-2)} \sum_{1 \leq i < j \leq n} (E_{i,j} - E_{j,i})^2.$$

The action of every $E_{i,j} \in \mathfrak{gl}_n(\mathbb{C})$ on $\mathcal{P}(W)$, corresponding to the derived action of \mathbf{G}_n , is given by polarization operators, that is

$$(35) \quad E_{i,j} = -\sum_{a=1}^r y_{j,a} \partial_{i,a}.$$

Finally, the action of the degree operator is given by $\mathbf{E} = \sum_{i=1}^n \sum_{j=1}^r y_{i,j} \partial_{i,j}$. Thus, expanding the right-hand side of (32), and using (33), (34), and (35), we obtain

$$D^{(1)} = -2(n-2)\Omega_{\mathfrak{k}} + \Omega_{\mathfrak{g}} - \mathbf{E}.$$

Case II. – $\mathbb{F} = \mathbb{C}$. In this case A is the space of $r \times r$ complex Hermitian matrices. The orthonormal basis of A with respect to $(\cdot, \cdot)_A$ is

$$E_{i,i} \text{ for } 1 \leq i \leq r, \quad \frac{1}{\sqrt{2}}(E_{i,j} + E_{j,i}) \text{ for } 1 \leq i < j \leq r, \text{ and } \frac{\sqrt{-1}}{\sqrt{2}}(E_{i,j} - E_{j,i}) \text{ for } 1 \leq i < j \leq r.$$

Moreover, $W = \text{Mat}_{n \times r}(\mathbb{C})$, considered as a real vector space. Fix generators $x_{i,j}, y_{i,j} \in \mathcal{P}(W)$, $1 \leq i \leq n, 1 \leq j \leq r$ for the algebra $\mathcal{P}(W)$, such that

$$x_{i,j}([t_{a,b}]) := \Re(t_{i,j}) \text{ and } y_{i,j}([t_{a,b}]) := \Im(t_{i,j}).$$

The map $\psi^* : \mathcal{P}(A) \rightarrow \mathcal{P}(W)$ of (6) is given by

$$\psi^*(\iota_A(S)) := \begin{cases} \sum_{a=1}^n x_{a,i}^2 + y_{a,i}^2 & \text{if } S = E_{i,i} \text{ where } 1 \leq i \leq r, \\ \sqrt{2} \sum_{a=1}^n (x_{a,i} x_{a,j} + y_{a,i} y_{a,j}) & \text{if } S = \frac{1}{\sqrt{2}}(E_{i,j} + E_{j,i}) \text{ where } 1 \leq i < j \leq r, \\ \sqrt{2} \sum_{a=1}^n (-x_{a,j} y_{a,i} + x_{a,i} y_{a,j}) & \text{if } S = \frac{\sqrt{-1}}{\sqrt{2}}(E_{i,j} - E_{j,i}) \text{ where } 1 \leq i < j \leq r. \end{cases}$$

For the realization of the derived action of $\mathbf{G}_n \simeq \text{GL}_n(\mathbb{C}) \times \text{GL}_n(\mathbb{C})$ on $\mathcal{P}(W)$ it will be more convenient to work with the coordinates $z_{i,j}$ and $\xi_{i,j}$ on $W_{\mathbb{C}} \simeq \text{Mat}_{n \times r}(\mathbb{C}) \oplus \text{Mat}_{n \times r}(\mathbb{C})$, where $1 \leq i \leq n$ and $1 \leq j \leq r$, given by

$$z_{i,j} := x_{i,j} - \sqrt{-1}y_{i,j} \text{ and } \xi_{i,j} := x_{i,j} + \sqrt{-1}y_{i,j}.$$

From these formulas it follows that $\frac{\partial}{\partial x_{i,j}} = \frac{\partial}{\partial z_{i,j}} + \frac{\partial}{\partial \xi_{i,j}}$ and $\frac{\partial}{\partial y_{i,j}} = \sqrt{-1} \left(\frac{\partial}{\partial \xi_{i,j}} - \frac{\partial}{\partial z_{i,j}} \right)$.

Set $\partial_{\xi_{i,j}} := \frac{\partial}{\partial \xi_{i,j}}$ and $\partial_{z_{i,j}} := \frac{\partial}{\partial z_{i,j}}$. By a direct calculation, we obtain

$$\begin{aligned}
 D^{(1)} &= 4 \sum_{i=1}^r \left(\sum_{a=1}^n z_{a,i} \xi_{a,i} \right) \left(\sum_{a=1}^n \partial_{z_{a,i}} \partial_{\xi_{a,i}} \right) \\
 &+ 2 \sum_{1 \leq i < j \leq r} \left(\sum_{a=1}^n z_{a,i} \xi_{a,j} + z_{a,j} \xi_{a,i} \right) \left(\sum_{a=1}^n \partial_{z_{a,i}} \partial_{\xi_{a,j}} + \partial_{z_{a,j}} \partial_{\xi_{a,i}} \right) \\
 &+ 2 \sum_{1 \leq i < j \leq r} \left(\sum_{a=1}^n z_{a,i} \xi_{a,j} - z_{a,j} \xi_{a,i} \right) \left(\sum_{a=1}^n \partial_{z_{a,i}} \partial_{\xi_{a,j}} - \partial_{z_{a,j}} \partial_{\xi_{a,i}} \right).
 \end{aligned}$$

Next we describe the derived action of \mathbf{G}_n on $\mathcal{P}(W)$. The Lie algebra of \mathbf{G}_n is isomorphic to the direct sum $\mathfrak{gl}_n(\mathbb{C}) \oplus \mathfrak{gl}_n(\mathbb{C})$. We will denote the matrices in the standard bases of the left and right summands of this direct sum by $E_{i,j}^{(l)}$ and $E_{i,j}^{(r)}$. Recall that $\mathcal{P}(W)$ can be identified with polynomials on the complex vector space $W_{\mathbb{C}}$. The actions of $E_{i,j}^{(l)}$ and $E_{i,j}^{(r)}$ on $\mathcal{P}(W)$ are by polarization operators $\sum_{a=1}^r z_{i,a} \partial_{z_{j,a}}$ and $-\sum_{a=1}^r \xi_{j,a} \partial_{\xi_{i,a}}$. The embedding of $\mathfrak{k}_{\mathbb{C}}$ into the Lie algebra of \mathbf{G}_n is the diagonal map $\mathfrak{gl}_n(\mathbb{C}) \hookrightarrow \mathfrak{gl}_n(\mathbb{C}) \oplus \mathfrak{gl}_n(\mathbb{C})$. The Casimir operators $\Omega_{\mathfrak{g}}$ and $\Omega_{\mathfrak{k}}$ are

$$(36) \quad \Omega_{\mathfrak{g}} = \sum_{1 \leq i,j \leq n} E_{i,j}^{(l)} E_{j,i}^{(l)} + E_{i,j}^{(r)} E_{j,i}^{(r)} \quad \text{and} \quad \Omega_{\mathfrak{k}} = \sum_{1 \leq i,j \leq n} \left(E_{i,j}^{(l)} + E_{i,j}^{(r)} \right) \left(E_{j,i}^{(l)} + E_{j,i}^{(r)} \right).$$

Finally, the degree operator is $\mathbf{E} = \sum_{i=1}^n \sum_{j=1}^r (z_{i,j} \partial_{z_{i,j}} + \xi_{i,j} \partial_{\xi_{i,j}})$. With a calculation similar to Case I, we obtain $D^{(1)} = -2\Omega_{\mathfrak{k}} + 2\Omega_{\mathfrak{g}} + (2n - 2r)\mathbf{E}$.

Case III. – $\mathbb{F} = \mathbb{H}$. The calculations are similar to the previous cases, only more elaborate. In this case, A is the space of hermitian quaternionic matrices. Let $\{1, \mathbf{i}, \mathbf{j}, \mathbf{k}\}$ denote the standard \mathbb{R} -basis of \mathbb{H} . Then the orthonormal basis of A consists of $E_{a,a}$, for $1 \leq a \leq r$, and

$$\frac{1}{\sqrt{2}}(E_{a,b} + E_{b,a}), \quad \frac{\mathbf{i}}{\sqrt{2}}(E_{a,b} - E_{b,a}), \quad \frac{\mathbf{j}}{\sqrt{2}}(E_{a,b} - E_{b,a}), \quad \text{and} \quad \frac{\mathbf{k}}{\sqrt{2}}(E_{a,b} - E_{b,a}), \quad \text{for } 1 \leq a < b \leq r.$$

Similar to Case II, it will be easier to use coordinates on $W_{\mathbb{C}}$. The embedding $W \hookrightarrow W_{\mathbb{C}}$ is given in matrix form by

$$\text{Mat}_{n \times r}(\mathbb{H}) \hookrightarrow \simeq \text{Mat}_{2n \times 2r}, \quad \underline{a} + \mathbf{i}\underline{b} + \mathbf{j}\underline{c} + \mathbf{k}\underline{d} \mapsto \begin{bmatrix} \underline{a} + \sqrt{-1}\underline{b} & -\underline{c} - \sqrt{-1}\underline{d} \\ \underline{c} - \sqrt{-1}\underline{d} & \underline{a} - \sqrt{-1}\underline{b} \end{bmatrix},$$

where $\underline{a}, \underline{b}, \underline{c}, \underline{d} \in \text{Mat}_{n \times r}(\mathbb{R})$. The new coordinates on $W_{\mathbb{C}}$, and the relations between the corresponding directional derivatives are as follows.

$$(37) \quad \begin{cases} \xi_{i,j} := \underline{a}_{i,j} + \sqrt{-1}\underline{b}_{i,j}, \\ \xi_{i+n,j+r} := \underline{a}_{i,j} - \sqrt{-1}\underline{b}_{i,j}, \\ \xi_{i+n,j} := \underline{c}_{i,j} - \sqrt{-1}\underline{d}_{i,j}, \\ \xi_{i,j+r} := -\underline{c}_{i,j} - \sqrt{-1}\underline{d}_{i,j}. \end{cases} \quad \text{and} \quad \begin{cases} \partial_{\underline{a}_{i,j}} = \partial_{\xi_{i,j}} + \partial_{\xi_{i+n,j+r}}, \\ \partial_{\underline{b}_{i,j}} = \sqrt{-1}(\partial_{\xi_{i,j}} - \partial_{\xi_{i+n,j+r}}), \\ \partial_{\underline{c}_{i,j}} = \partial_{\xi_{i+n,j}} - \partial_{\xi_{i,j+r}}, \\ \partial_{\underline{d}_{i,j}} = -\sqrt{-1}(\partial_{\xi_{i+n,j}} + \partial_{\xi_{i,j+r}}). \end{cases}$$

Next we define

$$\Phi_e(a, b) := \begin{cases} \sum_{i=1}^n \xi_{i,a} \xi_{i+n,b+r} & \text{if } e = 1, \\ \sum_{i=1}^n \xi_{i,a} \xi_{i+n,b} & \text{if } e = 2, \\ \sum_{i=1}^n \xi_{i+n,a} \xi_{i,b+r} & \text{if } e = 3, \\ \sum_{i=1}^n \xi_{i,a+r} \xi_{i+n,b+r} & \text{if } e = 4. \end{cases}$$

In the coordinates defined in (37), the map $\psi^* : \mathcal{P}(A) \rightarrow \mathcal{P}(W)$ of (6) is given by

$$\psi^*(\iota_A(S)) := \begin{cases} \Phi_1(a, a) - \Phi_3(a, a) & \text{if } S = E_{a,a}, \\ \frac{1}{\sqrt{2}} (\Phi_1(a, b) + \Phi_1(b, a) - \Phi_3(a, b) - \Phi_3(b, a)) & \text{if } S = \frac{1}{\sqrt{2}}(E_{a,b} + E_{b,a}), \\ \frac{\sqrt{-1}}{\sqrt{2}} (\Phi_1(a, b) - \Phi_1(b, a) + \Phi_3(b, a) - \Phi_3(a, b)) & \text{if } S = \frac{i}{\sqrt{2}}(E_{a,b} - E_{b,a}), \\ \frac{1}{\sqrt{2}} (\Phi_2(a, b) - \Phi_4(b, a) - \Phi_2(b, a) + \Phi_4(a, b)) & \text{if } S = \frac{j}{\sqrt{2}}(E_{a,b} - E_{b,a}), \\ \frac{1}{\sqrt{2}} (\Phi_2(a, b) + \Phi_4(b, a) - \Phi_2(b, a) - \Phi_4(a, b)) & \text{if } S = \frac{k}{\sqrt{2}}(E_{a,b} - E_{b,a}). \end{cases}$$

For $1 \leq e \leq 4$, set $\partial\Phi_e(a, b) := (\iota_W)^{-1}(\Phi_e(a, b))$, where $\iota_W : \mathcal{D}(W) \rightarrow \mathcal{P}(W)$ is the isomorphism given in (12). In fact $\partial\Phi_e(a, b)$ is the constant coefficient differential operator obtained from $\Phi_e(a, b)$ by substitution of each variable $\xi_{i,j}$ by the corresponding partial derivative $\partial_{\xi_{i,j}}$. Then

$$\begin{aligned} D^{(1)} &= 4 \sum_{1 \leq a, b \leq r} (\Phi_1(a, b) - \Phi_3(a, b)) (\partial\Phi_1(a, b) - \partial\Phi_3(a, b)) \\ &\quad + 2 \sum_{1 \leq a, b \leq r} (\Phi_2(a, b) - \Phi_2(b, a)) (\partial\Phi_2(a, b) - \partial\Phi_2(b, a)) \\ &\quad + 2 \sum_{1 \leq a, b \leq r} (\Phi_4(a, b) - \Phi_4(b, a)) (\partial\Phi_4(a, b) - \partial\Phi_4(b, a)). \end{aligned}$$

The embedding of $\mathfrak{k}_{\mathbb{C}}$ into the Lie algebra of \mathbf{G}_n gives the realization of $\mathfrak{k}_{\mathbb{C}}$ as

$$\mathfrak{k}_{\mathbb{C}} := \{x \in \text{Mat}_{2n \times 2n}(\mathbb{C}) : x^T J_n = -J_n x\},$$

where J_n is as in (5), and x^T denotes the transpose of the matrix x . We will denote the matrices in the standard basis of $\mathfrak{gl}_{2n}(\mathbb{C})$ by $E_{i,j}$'s. The Killing form of $\mathfrak{k}_{\mathbb{C}}$ is $\kappa(x, y) := (4n + 2)\text{tr}(xy)$, and consequently,

$$\begin{aligned} \Omega_{\mathfrak{k}} &= \frac{1}{4(2n + 1)} \sum_{1 \leq p, q \leq n} (E_{p,q} - E_{q+n,p+n}) (E_{q,p} - E_{p+n,q+n}) \\ &\quad + \frac{1}{4(2n + 1)} \sum_{1 \leq p < q \leq n} (E_{p,q+n} + E_{q,p+n}) (E_{q+n,p} + E_{p+n,q}) + \frac{1}{2(2n + 1)} \sum_{1 \leq p \leq n} E_{p,p+n} E_{p+n,p} \\ &\quad + \frac{1}{4(2n + 1)} \sum_{1 \leq p < q \leq n} (E_{p+n,q} + E_{q+n,p}) (E_{q,p+n} + E_{p,q+n}) + \frac{1}{2(2n + 1)} \sum_{1 \leq p \leq n} E_{p+n,p} E_{p,p+n}. \end{aligned}$$

As in Cases I and II,

$$(38) \quad \Omega_{\mathfrak{g}} = \sum_{1 \leq p, q \leq 2n} E_{p,q} E_{q,p}.$$

The action of $E_{p,q}$ on $\mathcal{P}(W)$ is by the polarization operator $-\sum_{i=1}^{2r} \xi_{q,s} \partial_{\xi_{p,s}}$. Finally, the degree operator is $E = \sum_{i=1}^{2n} \sum_{j=1}^{2r} \xi_{i,j} \partial_{\xi_{i,j}}$. From all of the above, and by a straightforward calculation, we obtain $D^{(1)} = -8(2n+1)\Omega_{\mathfrak{e}} + 2\Omega_{\mathfrak{g}} + 2(2n-2r+1)E$.

BIBLIOGRAPHY

- [1] A. ALLDRIDGE, S. SAHI, H. SALMASIAN, Schur Q -functions and the Capelli eigenvalue problem for the Lie superalgebra $\mathfrak{q}(n)$, in *Representation theory and harmonic analysis on symmetric spaces*, Contemp. Math. **714**, Amer. Math. Soc., 2018, 1–21.
- [2] J. FARAUT, A. KORÁNYI, *Analysis on symmetric cones*, Oxford Mathematical Monographs, The Clarendon Press Univ. Press, 1994.
- [3] F. GAVARINI, A Brauer algebra-theoretic proof of Littlewood’s restriction rules, *J. Algebra* **212** (1999), 240–271.
- [4] R. GOODMAN, N. R. WALLACH, *Symmetry, representations, and invariants*, Graduate Texts in Math. **255**, Springer, 2009.
- [5] S. HELGASON, *Groups and geometric analysis*, Mathematical Surveys and Monographs **83**, Amer. Math. Soc., 2000.
- [6] R. HOWE, Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond, in *The Schur lectures (1992) (Tel Aviv)*, Israel Math. Conf. Proc. **8**, Bar-Ilan Univ., Ramat Gan, 1995, 1–182.
- [7] R. HOWE, S. T. LEE, Spherical harmonics on Grassmannians, *Colloq. Math.* **118** (2010), 349–364.
- [8] R. HOWE, E.-C. TAN, J. F. WILLENBRING, Stable branching rules for classical symmetric pairs, *Trans. Amer. Math. Soc.* **357** (2005), 1601–1626.
- [9] R. HOWE, T. UMEDA, The Capelli identity, the double commutant theorem, and multiplicity-free actions, *Math. Ann.* **290** (1991), 565–619.
- [10] A. W. KNAPP, *Representation theory of semisimple groups*, Princeton Mathematical Series **36**, Princeton Univ. Press, 1986.
- [11] F. KNOP, Symmetric and non-symmetric quantum Capelli polynomials, *Comment. Math. Helv.* **72** (1997), 84–100.
- [12] F. KNOP, S. SAHI, Difference equations and symmetric polynomials defined by their zeros, *Int. Math. Res. Not.* **1996** (1996), 473–486.
- [13] K. KOIKE, I. TERADA, Young diagrammatic methods for the restriction of representations of complex classical Lie groups to reductive subgroups of maximal rank, *Adv. Math.* **79** (1990), 104–135.
- [14] T. H. KOORNWINDER, Okounkov’s BC -type interpolation MacDonalld polynomials and their $q = 1$ limit, *Sém. Lothar. Combin.* **72** (2014/15), Art. B72a.
- [15] B. KOSTANT, S. SAHI, The Capelli identity, tube domains, and the generalized Laplace transform, *Adv. Math.* **87** (1991), 71–92.
- [16] B. KOSTANT, S. SAHI, Jordan algebras and Capelli identities, *Invent. math.* **112** (1993), 657–664.

- [17] D. E. LITTLEWOOD, On invariant theory under restricted groups, *Philos. Trans. Roy. Soc. London. Ser. A.* **239** (1944), 387–417.
- [18] I. G. MACDONALD, *Symmetric functions and Hall polynomials*, second ed., Oxford Classic Texts in the Physical Sciences, The Clarendon Press Univ. Press, 2015.
- [19] A. OKOUNKOV, BC-type interpolation Macdonald polynomials and binomial formula for Koornwinder polynomials, *Transform. Groups* **3** (1998), 181–207.
- [20] S. SAHI, The spectrum of certain invariant differential operators associated to a Hermitian symmetric space, in *Lie theory and geometry*, Progr. Math. **123**, Birkhäuser, 1994, 569–576.
- [21] S. SAHI, Interpolation, integrality, and a generalization of Macdonald’s polynomials, *Int. Math. Res. Not.* **1996** (1996), 457–471.
- [22] S. SAHI, The Capelli identity for Grassmann manifolds, *Represent. Theory* **17** (2013), 326–336.
- [23] S. SAHI, H. SALMASIAN, The Capelli problem for $\mathfrak{gl}(m|n)$ and the spectrum of invariant differential operators, *Adv. Math.* **303** (2016), 1–38.
- [24] S. SAHI, H. SALMASIAN, V. SERGANOVA, The Capelli problem for basic classical Lie superalgebras, *Math. Zeitschrift* (2019), doi:10.1007/s00209-019-02289-7.
- [25] S. SAHI, G. ZHANG, The Capelli identity and Radon transform for Grassmannians, *Int. Math. Res. Not.* **2017** (2017), 3774–3800.
- [26] S. SAHI, G. ZHANG, Positivity of Shimura operators, preprint arXiv:1606.05276, to appear in *Math. Res. Letters*.
- [27] A. N. SERGEEV, A. P. VESELOV, BC_∞ Calogero-Moser operator and super Jacobi polynomials, *Adv. Math.* **222** (2009), 1687–1726.
- [28] R. P. STANLEY, Some combinatorial properties of Jack symmetric functions, *Adv. Math.* **77** (1989), 76–115.
- [29] R. P. STANLEY, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Math. **62**, Cambridge Univ. Press, 1999.
- [30] N. R. WALLACH, Polynomial differential operators associated with Hermitian symmetric spaces, in *Representation theory of Lie groups and Lie algebras (Fuji-Kawaguchiko, 1990)*, World Sci. Publ., River Edge, NJ, 1992, 76–94.

(Manuscrit reçu le 30 septembre 2016 ;
 accepté, après révision, le 13 décembre 2017.)

Siddhartha SAHI
 Department of Mathematics,
 Rutgers University,
 110 Frelinghuysen Rd,
 Piscataway, NJ 08854–8019.
 E-mail: sahi@math.rutgers.edu

Hadi SALMASIAN
 Department of Mathematics and Statistics,
 University of Ottawa,
 585 King Edward Ave,
 Ottawa, Ontario Canada K1N 6N5.
 E-mail: hsalmasi@uottawa.ca