

## $\mathbb{Z}R$ and rings of Witt vectors $W_S(R)$

CHRISTOPHER DENINGER (\*) – ANTON MELLIT (\*\*)

ABSTRACT – Using  $\lambda$  operations, we give some results on the kernel of the natural map from the monoid algebra  $\mathbb{Z}R$  of a commutative ring  $R$  to the ring of  $S$ -Witt vectors of  $R$ . As a byproduct we obtain a very natural interpretation of a power series used by Dwork in his proof of the rationality of zeta functions for varieties over finite fields.

MATHEMATICS SUBJECT CLASSIFICATION (2010). Primary: 13F35; Secondary: 13J05.

KEYWORDS. Witt vectors, monoid rings, lambda rings

### 1. Introduction

For a commutative ring  $R$ , let  $\mathbb{Z}R$  be the monoid algebra of  $(R, \cdot)$ . Let  $T$  be a divisor stable subset of the natural numbers  $\mathbb{N}$  and consider the ring  $W_T(R)$  of  $T$ -Witt vectors. The Teichmüller map  $R \rightarrow W_T(R)$  is multiplicative and hence extends uniquely to a ring homomorphism  $\alpha_T: \mathbb{Z}R \rightarrow W_T(R)$ . We are interested in the kernel of this map. If  $R$  has no  $T$ -torsion, the ghost map  $\mathcal{G}_T: W_T(R) \rightarrow R^T$  is injective and hence  $\ker \alpha_T = \ker(\mathcal{G}_T \circ \alpha_T)$  consists of the elements  $x = \sum_{r \in R} n_r [r] \in \mathbb{Z}R$  which satisfy the equations

$$\sum_{r \in R} n_r r^v = 0 \quad \text{for } v \in T.$$

If  $R$  is a perfect  $\mathbb{F}_p$ -algebra and  $T = \{1, \dots, p^{n-1}\}$  then  $\ker \alpha_T = I^n$ , where  $I$  is the kernel of the map  $\mathbb{Z}R \rightarrow R$  sending  $x$  to  $\sum n_r r$ . In this case the induced map

(\*) *Indirizzo dell'A.*: Faculty of Mathematics and Computer Science, University of Münster, Einsteinstraße 62, 48149 Münster, Germany

E-mail: [c.deninger@uni-muenster.de](mailto:c.deninger@uni-muenster.de)

(\*\*) *Indirizzo dell'A.*: Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

E-mail: [anton.mellit@univie.ac.at](mailto:anton.mellit@univie.ac.at)

$\mathbb{Z}R/I^n \xrightarrow{\sim} W_T(R)$  is actually an isomorphism, see [2]. More generally,  $\ker \alpha_T$  is known for all  $\mathbb{F}_p$ -algebras  $R$  with injective Frobenius map, see [3], Theorem 7.1.

In the present note we use  $\lambda$ -ring structures to describe  $\ker \alpha_T$  for more general rings  $R$  and certain subsets  $T = S_N$  obtained as follows. Fix a divisor stable subset  $S$  which is also multiplicatively closed. Thus  $S$  consists of all natural numbers whose prime divisors lie in a given set of prime numbers. Fix some  $1 \leq N \leq \infty$  and set  $S_N := \{v \in S \mid v < N\}$ . For a  $\mathbb{Z}_S = \mathbb{Z}[p^{-1}, p \notin S]$ -algebra  $R$  all its (truncated) Witt rings are  $\mathbb{Z}_S$ -algebras as well, see [5], Lemma 1.9. Thus the Teichmüller map  $R \rightarrow W_{S_N}(R)$  induces a homomorphism of  $\mathbb{Z}_S$ -algebras:

$$(1) \quad \alpha_{S_N}: \mathbb{Z}_S R = \mathbb{Z}R \otimes_{\mathbb{Z}} \mathbb{Z}_S \longrightarrow W_{S_N}(R).$$

Let  $\pi: \mathbb{Z}_S R \rightarrow R$  be the map sending  $\sum n_r[r]$  to  $\sum n_r r$ , and for an integer  $n \geq 1$  write  $n_S = \prod_{p \in S} p^{\text{ord}_p(n)}$ . Then  $n_S \in S$  since  $S$  is multiplicatively closed. The following result holds:

**THEOREM 1.1.** *Consider the unique (special)  $\lambda$ -ring structure  $(\lambda_S^n)$  on  $\mathbb{Z}_S R$  whose associated Adams operators  $\psi_S^n: \mathbb{Z}_S R \rightarrow \mathbb{Z}_S R$  are determined by the formula  $\psi_S^n[r] = [r]^{n_S}$  for  $r \in R$ . Then we have*

$$\ker \alpha_{S_N} = \{x \in \mathbb{Z}_S R \mid \pi \lambda_S^n(x) = 0 \text{ for } 1 \leq n < N\}.$$

Existence and uniqueness of the special  $\lambda$ -ring structure are special cases of a classical result [7], Proposition 1.2.

**LEMMA 1.2 (Wilkerson).** *Let  $B$  be commutative ring without  $\mathbb{Z}$ -torsion and for  $n \geq 1$  let  $\psi^n$  be a family of ring endomorphisms of  $B$  such that  $\psi^1 = \text{id}$  and  $\psi^n \circ \psi^m = \psi^{nm}$  and such that  $\psi^p(b) \equiv b^p \pmod{pB}$  for all  $b \in B$  and all prime numbers  $p$ . Then there is a unique structure of a (special)  $\lambda$ -ring on  $B$  whose Adams operators are the given maps  $\psi^n$ .*

The main ingredient in the proof of Theorem 1.1 is a (unital) ring homomorphism  $\bar{\varphi}_S: W_S(R) \rightarrow W(R)$  for  $\mathbb{Z}_S$ -algebras  $R$  which splits the canonical projection  $W(R) \rightarrow W_S(R)$ .

Adapting a method of Dwork in the theory of  $p$ -adic formal power series, we obtain explicit albeit complicated formulas for the operations  $\lambda_S^n$  in Theorem 1.1. They are given as follows. Let  $\mu$  be the Moebius function and for  $x \in \mathbb{Z}_S R$  and  $k \in S$  set

$$(2) \quad \tau_k(x) = k^{-1} \sum_{d|k} \mu(d) \psi_S^{k/d}(x).$$

For an  $S$ -tuple  $\nu = (\nu_k)_{k \in S}$  with all  $\nu_k \geq 0$  we write

$$\binom{\tau(x)}{\nu} = \prod_{k \in S} \binom{\tau_k(x)}{\nu_k} \in \mathbb{Z}R \otimes_{\mathbb{Z}} \mathbb{Q}.$$

Set  $|\nu| = \sum_{k \in S} \nu_k$  and  $\|\nu\| = \sum_{k \in S} k \nu_k$ .

**THEOREM 1.3.** *In the situation of Theorem 1.1, for  $n \geq 1$  and  $x \in \mathbb{Z}_S R$  the following explicit formula holds in  $\mathbb{Z}_S R$ :*

$$(-1)^n \lambda_S^n(x) = \sum_{\|\nu\|=n} (-1)^{|\nu|} \binom{\tau(x)}{\nu}.$$

The methods work in the more general situation where instead of  $\mathbb{Z}_S R$  we start with a  $\mathbb{Z}$ -torsionfree  $\mathbb{Z}_S$ -algebra  $B$  which is equipped with commuting Frobenius lifts  $\psi_S^p$  for all primes  $p \in S$ . Setting  $\psi_S^p = \text{id}$  for  $p \notin S$ , we show that the corresponding  $\lambda$ -ring structure is given by the same formula as in Theorem 1.3. Moreover any homomorphism  $\pi: B \rightarrow R$  into a  $\mathbb{Z}_S$ -algebra  $R$  factors canonically over maps  $\alpha_{S_N}: B \rightarrow W_{S_N}(R)$  and the kernel of  $\alpha_{S_N}$  is obtained as in Theorem 1.1.

For background on the theory of Witt vectors we refer to [5], [6], [1], and [3]. The latter approaches avoid universal polynomials.

*Acknowledgment.* Both authors would like to thank the HIM in Bonn, where this work originated for its support.

## 2. A projector on Witt vector rings

All rings are associative, commutative and unital. All ring homomorphisms are unital. For any commutative ring  $A$  we give  $A$  the discrete topology and  $A^{\mathbb{N}}$  the product topology. Then  $W(A) \cong A^{\mathbb{N}}$  is a topological ring and the ghost map  $\mathcal{G}: W(A) \rightarrow A^{\mathbb{N}}$  is continuous.

For  $S$  as in the introduction, consider the ring homomorphism:

$$\varphi_S: A^{\mathbb{N}} \longrightarrow A^{\mathbb{N}}, \quad \varphi_S((a_n)_{n \geq 1}) = (a_{n_S})_{n \geq 1}.$$

It maps  $0 \times A^{\mathbb{N} \setminus S}$  to zero and therefore factors over  $A^S \cong A^{\mathbb{N}} / (0 \times A^{\mathbb{N} \setminus S})$ . Note that  $\varphi_S^2 = \varphi_S$ . For  $k \geq 1$ , Frobenius and Verschiebung maps from  $A^{\mathbb{N}}$  to  $A^{\mathbb{N}}$  are defined as follows:

$$F_k((a_n)_{n \geq 1}) = (a_{nk})_{n \geq 1} \quad \text{and} \quad V_k((a_n)_{n \geq 1}) = k(\delta_{k|n} a_{n/k})_{n \geq 1}.$$

Here,  $\delta_{k|n} = 1$  if  $k \mid n$  and  $= 0$  if  $k \nmid n$ . Now assume that  $A$  is a  $\mathbb{Z}_S$ -algebra and for any prime  $l \notin S$  consider the map

$$T_l = 1 + l^{-1}V_l(1 - F_l): A^{\mathbb{N}} \longrightarrow A^{\mathbb{N}}.$$

It is a ring endomorphism because of the formula:

$$T_l((a_n)_{n \geq 1}) = (b_n)_{n \geq 1}$$

where  $b_n = a_n$  if  $l \nmid n$  and  $b_n = a_{n/l}$  if  $l \mid n$ . For different primes  $l$  and  $l'$  not in  $S$ , the endomorphisms  $T_l$  and  $T_{l'}$  commute with each other. For  $m$  prime to  $S$ , set

$$T_m = \prod_l T_l^{\text{ord}_l m}.$$

Then the following limit formula holds in the pointwise topology:

$$(3) \quad \varphi_S = \lim_{v \rightarrow \infty} T_{m_v}.$$

Here  $(m_v)$  is any sequence of positive integers prime to  $S$  such that  $m_v \mid m_{v+1}$  for all  $v$  and such that any number prime to  $S$  is a divisor of some  $m_v$ . For example, if  $l_1, l_2, \dots$  are the primes not in  $S$  we could take  $m_v = (l_1 \cdots l_v)^v$ .

There are also other ways to express  $\varphi_S$ . Firstly, we have

$$\lim_{v \rightarrow \infty} T_{l^v} = \left( \sum_{k=0}^{\infty} l^{-k} V_{l^k} \right) (1 - l^{-1} V_l F_l).$$

This can be either verified directly by looking at the action on sequences or deduced from the formula for  $T_l$  and the identity  $F_k \circ V_k = k$ . This leads to the following formula, where the sums are over all positive integers prime to  $S$ :

$$\varphi_S = \left( \sum_{(n,S)=1} n^{-1} V_n \right) \left( \sum_{(n,S)=1} \mu(n) n^{-1} V_n F_n \right).$$

Frobenius and Verschiebung operators also exist on the ring  $W(A)$  of (big) Witt vectors and they correspond to Frobenius and Verschiebung on  $A^{\mathbb{N}}$  via the ghost map  $\mathfrak{G}: W(A) \rightarrow A^{\mathbb{N}}$ . If the  $\mathbb{Z}_S$ -algebra  $A$  has no  $\mathbb{Z}$ -torsion, then  $\mathfrak{G}$  is injective and it follows from formula (3) that there is a unique ring homomorphism  $\varphi_S: W(A) \rightarrow W(A)$  making the diagram

$$\begin{array}{ccc} W(A) & \xrightarrow{\mathfrak{G}} & A^{\mathbb{N}} \\ \varphi_S \downarrow & & \downarrow \varphi_S \\ W(A) & \xrightarrow{\mathfrak{G}} & A^{\mathbb{N}} \end{array}$$

commute. It also follows that  $\varphi_S$  factors uniquely over the canonical projection  $\varphi_S: W(A) \xrightarrow{\text{pr}_S} W_S(A) \xrightarrow{\bar{\varphi}_S} W(A)$ .

We have

$$(4) \quad \text{pr}_S \circ \bar{\varphi}_S = \text{id},$$

since this is true after applying the ghost map.

Now let  $R$  be any  $\mathbb{Z}_S$ -algebra and define  $\varphi_S: W(R) \rightarrow W(R)$  by the pointwise limit (3). Convergence to a well defined ring homomorphism follows by comparison with the map  $\varphi_S$  for a  $\mathbb{Z}$ -torsion free  $\mathbb{Z}_S$ -algebra  $A$  surjecting onto  $R$ . In the same way we prove a unique factorization

$$(5) \quad \varphi_S: W(R) \xrightarrow{\text{pr}_S} W_S(R) \xrightarrow{\bar{\varphi}_S} W(R)$$

and the formula

$$(6) \quad \text{pr}_S \circ \bar{\varphi}_S = \text{id} \quad \text{on } W_S(R).$$

In particular the (unital) ring homomorphism  $\bar{\varphi}_S: W_S(R) \hookrightarrow W(R)$  is injective. By construction the maps  $\varphi_S$  and  $\bar{\varphi}_S$  are functorial with respect to  $R$ . It is clear that  $\varphi_S^2 = \varphi_S$ .

We need a version of the maps  $\bar{\varphi}$  for the truncation sets  $S_N$ : for any  $\mathbb{Z}_S$ -algebra  $R$  without  $\mathbb{Z}$ -torsion, it follows by comparing with the ghost side that there is a unique ring homomorphism  $\bar{\varphi}_{S_N}$  such that the diagram

$$(7) \quad \begin{array}{ccc} W_S(R) & \longrightarrow & W_{S_N}(R) \\ \bar{\varphi}_S \downarrow & & \downarrow \bar{\varphi}_{S_N} \\ W(R) & \longrightarrow & W_N(R) \end{array}$$

commutes. Here  $W_N(R) = W_{\{1 \leq v < N\}}(R)$ . The point is that  $n < N$  implies  $n_S < N$ . For the projection  $\text{pr}_{S_N}: W_N(R) \rightarrow W_{S_N}(R)$  we have  $\text{pr}_{S_N} \circ \bar{\varphi}_{S_N} = \text{id}$ . Similarly as before, it follows that unique functorial ring homomorphisms  $\bar{\varphi}_{S_N}$  with the same properties exist for arbitrary  $\mathbb{Z}_S$ -algebras  $R$ .

### 3. Proofs of theorems 1.1 and 1.3

For any ring  $A$  we give the set  $\Lambda(A) = 1 + tA[[t]]$  the unique ring structure, for which the bijection

$$(8) \quad W(A) = A^{\mathbb{N}} \xrightarrow{\sim} \Lambda(A), \quad (a_1, a_2, \dots) \mapsto \prod_{n=1}^{\infty} (1 - a_n t^n)$$

is an isomorphism. Then  $\Lambda(A)$  is a topological ring for the  $t$ -adic topology whose multiplication is uniquely determined by the formula

$$(1 - a_1 t) \cdot (1 - a_2 t) = 1 - a_1 a_2 t \quad \text{for } a_1, a_2 \in A.$$

The addition in  $\Lambda(A)$  is given by the multiplication of power series. We will usually view the topological isomorphism (8) as an identification. There is a commutative diagram of ring homomorphisms

$$(9) \quad \begin{array}{ccc} W(A) & \xrightarrow{\mathcal{G}} & A^{\mathbb{N}} \\ \parallel & & \parallel \\ \Lambda(A) & \xrightarrow{-t\partial_t \log} & tA[[t]]. \end{array}$$

Here we view  $tA[[t]]$  as a commutative ring with the coefficientwise multiplication of power series, the Hadamard product.

As before let  $S \subset \mathbb{N}$  be divisor stable and multiplicatively closed. Let  $B$  be a  $\mathbb{Z}$ -torsion free  $\mathbb{Z}_S$ -algebra with commuting Frobenius lifts  $\psi_S^p$  for all primes  $p \in S$ . For  $n \geq 1$  we set

$$\psi_S^n = \prod_{p \in S} (\psi_S^p)^{\text{ord}_{p^n}}.$$

Let  $(\lambda_S^n)$  be the special  $\lambda$ -ring structure on  $B$  with Adams operators  $\psi_S^n$ , according to Lemma 1.2. Consider the ring homomorphism

$$\lambda_S: B \longrightarrow \Lambda(B)$$

defined by the formula

$$\lambda_S(x) = \sum_{i=0}^{\infty} (-1)^i \lambda_S^i(x) t^i.$$

Setting

$$\psi_S(x) = \sum_{n=1}^{\infty} \psi_S^n(x) t^n$$

we have

$$\psi_S(x) = -t\partial_t \log \lambda_S(x).$$

Using diagram (9) for  $A = B$  we may interpret  $\lambda_S$  as the unique ring homomorphism  $\tilde{\alpha}: B \rightarrow W(B)$  such that  $\mathcal{G} \circ \tilde{\alpha}$  maps  $x \in B$  to  $(\psi_S^n(x))_{n \geq 1} \in B^{\mathbb{N}}$ . We have  $\tilde{\alpha} = \varphi_S \circ \tilde{\alpha}$  since after applying the injective ghost map, this amounts to the equality  $\psi_S^n = \psi_S^{n_S}$  for  $n \geq 1$ . Hence we get

$$\tilde{\alpha} = \varphi_S \circ \tilde{\alpha} = \bar{\varphi}_S \circ \text{pr}_S \circ \tilde{\alpha} = \bar{\varphi}_S \circ \tilde{\alpha}_S.$$

Here  $\tilde{\alpha}_S = \text{pr}_S \circ \tilde{\alpha}: B \rightarrow W_S(B)$  is the unique ring homomorphism such that  $\mathcal{G}_S \circ \tilde{\alpha}_S$  maps  $x \in B$  to  $(\psi_S^n(x))_{n \in S} \in B^S$ . In conclusion, we have a commutative diagram:

$$(10) \quad \begin{array}{ccc} B & \xrightarrow{\tilde{\alpha}_S} & W_S(B) \\ \lambda_S \downarrow & & \downarrow \bar{\varphi}_S \\ \Lambda(B) & \equiv \equiv \equiv & W(B). \end{array}$$

REMARK. In the case  $B = \mathbb{Z}_S R$  considered in Theorem 1.1, the map  $\tilde{\alpha}_S$  is the unique  $\mathbb{Z}_S$ -algebra homomorphism extending the multiplicative map  $R \rightarrow W_S(\mathbb{Z}_S R)$  which sends  $r$  to the Teichmüller representative of  $[r]$ . This follows by comparing ghost components.

Let  $\pi: B \rightarrow R$  be a map of  $\mathbb{Z}_S$ -algebras. For  $1 \leq N \leq \infty$  consider the composition

$$\alpha_{S_N}: B \xrightarrow{\tilde{\alpha}_S} W_S(B) \xrightarrow{W_S(\pi)} W_S(R) \longrightarrow W_{S_N}(R).$$

For  $B = \mathbb{Z}_S R \xrightarrow{\pi} R$ , by the remark on  $\tilde{\alpha}_S$  above,  $\alpha_{S_N}$  agrees with the map (1) in the introduction. Hence Theorems 1.1 and 1.3 are special cases of the following result:

THEOREM 3.1. *With notations as above, we have*

$$\text{Ker } \alpha_{S_N} = \{x \in B \mid \pi \lambda_S^n(x) = 0 \text{ for } 1 \leq n < N\}.$$

Moreover, with  $\tau_k(x) \in B \otimes \mathbb{Q}$  as in (2), the following formula holds in  $\Lambda(B)$ :

$$(11) \quad \lambda_S(x) = \prod_{k \in S} (1 - t^k)^{\tau_k(x)}.$$

Equivalently, with notations as in Theorem 1.3 we have

$$(-1)^n \lambda_S^n(x) = \sum_{\|\nu\|=n} (-1)^{|\nu|} \binom{\tau(x)}{\nu} \text{ in } B, \text{ for all } n \geq 1.$$

PROOF. Using diagrams (7), (10) and the functoriality of  $\bar{\varphi}_S$  we get a commutative diagram

$$\begin{array}{ccccccc} B & \xrightarrow{\tilde{\alpha}_S} & W_S(B) & \xrightarrow{W_S(\pi)} & W_S(R) & \longrightarrow & W_{S_N}(R) \\ \lambda_S \downarrow & & \downarrow \bar{\varphi}_S & & \downarrow \bar{\varphi}_S & & \downarrow \bar{\varphi}_{S_N} \\ \Lambda(B) & \equiv \equiv \equiv & W(B) & \xrightarrow{W(\pi)} & W(R) & \longrightarrow & W_N(R). \end{array}$$

Identifying  $W_N(R)$  with  $\Lambda_N(R) = \Lambda(R)/(1 + t^N R[[t]])$ , the outer square becomes

$$\begin{array}{ccc} B & \xrightarrow{\alpha_{S_N}} & W_{S_N}(R) \\ \lambda_S \downarrow & & \downarrow \bar{\varphi}_{S_N} \\ \Lambda(B) & \xrightarrow{\Lambda(\pi)} & \Lambda_N(R). \end{array}$$

Since  $\bar{\varphi}_{S_N}$  is injective being a splitting of the projection  $\text{pr}_{S_N}: W_N(R) \rightarrow W_{S_N}(R)$ , the first assertion of Theorem 3.1 follows. In order to prove formula (11) it suffices to show the equality after applying  $-t \partial_t \log$ , i.e. the formula

$$(12) \quad \psi_S(x) = \sum_{k=1}^{\infty} \tau_k(x) \frac{k t^k}{1 - t^k}.$$

Generally, we have the identity of formal power series

$$\sum_{k=1}^{\infty} a_k \frac{t^k}{1 - t^k} = \sum_{n=1}^{\infty} A_n t^n.$$

where  $A_n = \sum_{v|n} a_v$ . In our case  $a_k = k \tau_k(x)$ , we obtain

$$A_n = \sum_{v|n} \sum_{d|v} \mu(d) \psi_S^{v/d}(x) = \psi_S^n(x),$$

by Moebius inversion. Thus formula (12) and hence the Theorem are proved.  $\square$

A priori the product  $\prod_{k \in S} (1 - t^k)^{\tau_k(x)}$  lies in  $\Lambda(B \otimes \mathbb{Q})$  but its equality with  $\lambda_S(x)$  shows that it lies in  $\Lambda(B)$ . The required integrality comes from Wilkerson's Lemma 1.2 and the congruences used in its proof. For  $S = \{1, p, p^2, \dots\}$  such products were considered by Dwork in his proof of Weil's rationality conjecture for zeta functions of varieties over finite fields. For  $p^i \in S$  we have

$$\tau_1(x) = x \quad \text{and} \quad \tau_{p^i}(x) = p^{-i} (\psi_S^{p^i}(x) - \psi_S^{p^{i-1}}(x)) \text{ for } i \geq 1.$$

Thus equation (11) asserts

$$(13) \quad \lambda_S(x) = (1 - t)^x \prod_{i=1}^{\infty} (1 - t^{p^i})^{\tau_{p^i}(x)}.$$

In [4] p. 2, using slightly different notation, Dwork considers the following product in the formal power series ring  $\mathbb{Q}[[t, X]]$ :

$$(14) \quad F(X, t) = (1 + t)^X \prod_{i=1}^{\infty} (1 + t^{p^i})^{p^{-i}(X^{p^i} - X^{p^{i-1}})}.$$



Using his well-known criterion [4] Lemma 1, he shows that the coefficients of  $F(X, t)$  are  $p$ -integral. Our sign conventions concerning  $\Lambda$ - and Witt rings are not quite compatible with Dwork's. However, for odd  $p$  we can relate  $F(X, t)$  to  $\lambda_S$  and  $\varphi_S$  as follows. For  $S$  as above, we have  $\mathbb{Z}_S = \mathbb{Z}[l^{-1} \mid l \neq p]$ . Equip the  $\mathbb{Z}$ -torsion free  $\mathbb{Z}_S$ -algebra  $B = \mathbb{Z}_S[X]$  with the Frobenius lift  $\psi_S^p$  defined by  $\psi_S^p(X) = X^p$ . Then, as in the beginning of this section the corresponding  $\lambda$ -ring structure on  $\mathbb{Z}_S[X]$  is encoded in a ring homomorphism

$$\lambda_S: \mathbb{Z}_S[X] \longrightarrow \Lambda(\mathbb{Z}_S[X]) = 1 + t\mathbb{Z}_S[X][[t]] \subset \mathbb{Z}_S[[t, X]].$$

Comparing (13) and (14), we see that for  $p \neq 2$  we have

$$(15) \quad F(X, -t) = \lambda_S(X).$$

In particular the  $p$ -integrality of (14) follows. In terms of the map

$$\bar{\varphi}_S: W_S(\mathbb{Z}_S[X]) \rightarrow W(\mathbb{Z}_S[X]) \equiv \Lambda(\mathbb{Z}_S[X])$$

we have

$$F(X, -t) = \bar{\varphi}_S(\langle X \rangle).$$

Here  $\langle X \rangle$  is the Teichmüller representative of  $X$ . This follows from diagram (10) and formula (15) noting that  $\tilde{\alpha}_S(X) = \langle X \rangle$ . The latter equality holds because  $(\mathcal{G}_S \circ \tilde{\alpha}_S)(X) = (\psi_S^{p^i}(X)) = (X^{p^i})$  by the characterization of  $\tilde{\alpha}_S$  and since  $\mathcal{G}_S(\langle X \rangle) = (X^{p^i})$ .

*Acknowledgment.* Christopher Deninger's research is supported by the CRC 878 of the DFG. Anton Mellit's research is supported by the Austrian Science Fund (FWF) through the Projects P-31705 and Y963-N35.

## REFERENCES

- [1] J. BORGER – B. WIELAND, *Plethystic algebra*, Adv. Math. 194 (2005), no. 2, pp. 246–283.
- [2] J. CUNTZ – C. DENINGER, *An alternative to Witt vectors*, Münster J. Math. 7 (2014), no. 1, pp. 105–114.
- [3] J. CUNTZ – C. DENINGER, *Witt vector rings and the relative de Rham Witt complex*, with an appendix by U. Zannier, J. Algebra 440 (2015), pp. 545–593.
- [4] B. DWORK, *On the rationality of zeta functions and L-series*, In *Proceedings of a Conference on Local Fields* (Driebergen, 1966), edited by T. A. Springer, Springer-Verlag, Berlin etc., 1967, 40–55.
- [5] L. HESSELHOLT, *The big de Rham–Witt complex*, Acta Math. 214 (2015), no. 1, pp. 135–207.

- [6] J. RABINOFF, *The theory of Witt vectors*, preprint, 2014. [arXiv:1409.7445](https://arxiv.org/abs/1409.7445) [math.NT]
- [7] C. WILKERSON, *Lambda-rings, binomial domains, and vector bundles over  $\mathbf{C}P(\infty)$* , *Comm. Algebra* 10 (1982), no. 3, pp. 311–328.

Manoscritto pervenuto in redazione il 10 maggio 2018.