

On a causal quantum stochastic double product integral related to Lévy area

Robin L. Hudson and Yuchen Pei

Abstract. We study the family of causal double product integrals

$$\prod_{a < x < y < b} \left(1 + i \frac{\lambda}{2} (dP_x dQ_y - dQ_x dP_y) + i \frac{\mu}{2} (dP_x dP_y + dQ_x dQ_y) \right),$$

where P and Q are the mutually noncommuting momentum and position Brownian motions of quantum stochastic calculus. The evaluation is motivated heuristically by approximating the continuous double product by a discrete product in which infinitesimals are replaced by finite increments. The latter is in turn approximated by the second quantisation of a discrete double product of rotation-like operators in different planes due to a result in [15]. The main problem solved in this paper is the explicit evaluation of the continuum limit W of the latter, and showing that W is a unitary operator. The kernel of $W - I$ is written in terms of Bessel functions, and the evaluation is achieved by working on a lattice path model and enumerating linear extensions of related partial orderings, where the enumeration turns out to be heavily related to Dyck paths and generalisations of Catalan numbers.

Mathematics Subject Classification (2010). 81S25, 05A15, 06A07.

Keywords. Causal double product, Lévy’s stochastic area, position and momentum Brownian motions, linear extensions, Catalan numbers, Dyck paths.

Contents

1	Introduction	468
2	The Lévy stochastic area	470
3	A discrete double product of unitary matrices	472
4	A lattice path model and linear extensions of partial orderings	476
5	Dyck paths and Catalan numbers	487
6	Proof of Theorem 7	494
7	The unitarity of W	499
	References	509

1. Introduction

Following Volterra's philosophy of product integrals as continuous limits of discrete products [28], quantum stochastic double product integrals of *rectangular* type have been constructed [15] as limits of discrete approximations obtained by replacing stochastic differentials by discrete increments of the corresponding processes. Such constructions are partially intuitive in character, involving nonrigorous manipulations of unbounded operators. Nevertheless they can be shown to yield explicit rigorously unitary operators which can then be shown in some cases [11] to satisfy the quantum stochastic differential equations (qsde's) whose solutions provide the rigorous definition of the product integral.

In this paper we initiate the much harder problem of constructing so-called *causal* (or triangular) double product integrals in the same way, first constructing discrete approximations by intuitive methods, which are then shown rigorously to enjoy the property of unitarity, which will allow rigorous verification of the qsde definitions.

The *Fock space* $\mathcal{F}(\mathcal{H})$ over a Hilbert space \mathcal{H} is conveniently defined [26] as the Hilbert space generated by the *exponential vectors* $e(f)$, $f \in \mathcal{H}$, satisfying

$$\langle e(f), e(g) \rangle = \exp \langle f, g \rangle, \quad f, g \in \mathcal{H}.$$

Rectangular product integrals live in the tensor product of two Fock spaces. This form of "double" construction was originally motivated by its use to construct explicit solutions of the quantum Yang-Baxter equation with a given classical limit ([17] and [14]), of purely algebraic character as formal power series. From the analytic point of view, the alternative causal constructs which are studied in the present paper which live naturally in a single Fock space are of wider interest.

One example which we consider in some detail is closely related to Lévy's stochastic area [23], and in particular to the Lévy area formula for its characteristic function. In effect we replace the planar Brownian motion by a quantum version in which the two components are the mutually noncommuting momentum and position Brownian motions P and Q of quantum stochastic calculus [5], which despite noncommutativity, can be shown to be independent in a certain sense [19]. Other noncommutative analogs of Lévy area are based on free probability [4]; our own less radically noncommutative form is directly related to physical applications [9]. It may also offer mathematically significant relations, for example to Riemann zeta values through the links to Euler and Bernoulli numbers [21] of the classical Lévy area formula. This is because, while the corresponding probability distribution is the atomic one concentrated at zero, it deforms naturally to the classical distribution at infinite temperature as the Fock "zero temperature"

momentum and position processes P and Q are deformed through corresponding finite temperature processes [6] to mutually commuting independent Brownian motions.

Efforts have been made to make the present article self-contained. However to gain familiarity with certain notations and ideas the reader is encouraged to read the relevant paper [15], which can be viewed as a prequel. We denote rectangular and causal product integrals by

$$\prod_{[a,b] \times [c,d]} (1 + dr), \quad \prod_{<[a,b]} (1 + dr) \tag{1}$$

respectively, where $<[a,b]$ is the set $\{(x, y) \in \mathbb{R}^2 : a \leq x < y < b\}$. They are operators in the Hilbert spaces $\mathcal{F}(L^2([a, b])) \otimes \mathcal{F}(L^2([c, d]))$ and $\mathcal{F}(L^2([a, b]))$ respectively. Both are characterised by the generator dr which is a second rank tensor over the complex vector space $\mathcal{J} = \mathbb{C}\langle dP, dQ, dT \rangle$ of differentials of the fundamental stochastic processes P, Q and the time process T of the calculus. They have rigorous definitions as solutions of either forward or backward adapted quantum stochastic differential equations [20] in which b or a in 1 is the time variable. They are related by the *coboundary relation*

$$\prod_{<[a,c]} (1 + dr) = \left(\prod_{<[a,b]} (1 + dr) \otimes I \right) \prod_{[a,b] \times [b,c]} (1 + dr) \left(I \otimes \prod_{<[b,c]} (1 + dr) \right)$$

in which the Fock space $\mathcal{F}(L^2([a, c]))$ is canonically split at time $b \in [a, c]$;

$$\begin{aligned} \mathcal{F}(L^2([a, c])) &= \mathcal{F}(L^2([a, b]) \oplus L^2([b, c])) \\ &= \mathcal{F}(L^2([a, b])) \otimes \mathcal{F}(L^2([b, c])) \end{aligned}$$

allowing it to accommodate the operator $\prod_{[a,b] \times [b,c]} (1 + dr)$.

A necessary and sufficient condition that they consist of unitary operators is [20] that

$$dr + dr^\dagger + drdr^\dagger = 0.$$

Here the space $\mathcal{J} = \mathbb{C}\langle dP, dQ, dT \rangle$ is equipped with the multiplication given by the quantum Itô product rule

	dP	dQ	dT
dP	dT	$-idT$	0
dQ	idT	dT	0
dT	0	0	0

and $\mathcal{J} \otimes \mathcal{J}$ with the corresponding tensor product multiplication, together with the natural involution \dagger derived from the self-adjointness of P, Q and T .

Two examples of such unitary generators are

$$dr_1 = i(dP \otimes dQ - dQ \otimes dP),$$

$$dr_2 = i(dP \otimes dP + dQ \otimes dQ).$$

dr_1 relates to quantum Lévy area in which the independent one-dimensional component Brownian motions of planar Brownian motion are replaced by P and Q . In the same spirit, dr_2 relates to a quantum version of the Bessel process, the radial part of planar Brownian motion. The general form of unitary generator in which the time differential dT does not appear is [15] the real linear combination

$$dr_{\lambda,\mu} = \frac{\lambda}{2}dr_1 + \frac{\mu}{2}dr_2.$$

In this paper we begin the explicit construction of the unitary causal double product integral

$$E := \prod_{<[a,b]} (1 + dr_{\lambda,\mu})$$

as the second quantisation $\Gamma(W)$ of a unitary operator W which differs from the identity operator I by an integral operator on the Hilbert space $L^2([a, b])$ whose kernel will be found explicitly.

Acknowledgements. Parts of this work were completed when the Robin Hudson visited the Mathematics Department of Chungbuk National University in Korea, whose warm hospitality is gratefully acknowledged, along with conversations with John Gough, Paul Jones and Janosch Ortmann. Other parts were completed when Yuchen Pei visited the School of Mathematics of Trinity College Dublin, Mathematical Research and Conference Center of the Institute of Mathematics of the Polish Academy of Sciences, Chungbuk National University and Gyeongbukgung. Conversations with Neil O'Connell are also acknowledged. The research of Yuchen Pei was supported by EPSRC grant number EP/H023364/1 and Center of Mathematical Sciences and Applications at Harvard University. The authors would also like to thank the anonymous referee for many suggestions and queries which have resulted in clarifications to this paper.

2. The Lévy stochastic area

Before moving on to construct E , let us take a detour and explain the motivation of this problem.

The stochastic Lévy area introduced in [23] can be defined as the signed area formed by connecting the endpoints of a 2-dimensional Brownian path. More specifically, it is defined as

$$L = \frac{1}{2} \int_{0 \leq s_1 < s_2 < t} dB_{s_1}^1 dB_{s_2}^2 - dB_{s_1}^2 dB_{s_2}^1$$

where B^1 and B^2 are two independent Brownian motions. The Lévy area formula shows the characteristic function of L :

$$\mathbb{E} \prod_{0 \leq s_1 < s_2 < t} \left(1 + \frac{i\lambda}{2} (dB_{s_1}^1 dB_{s_2}^2 - dB_{s_1}^2 dB_{s_2}^1) \right) = \mathbb{E} e^{i\lambda L} = \operatorname{sech} \frac{\lambda t}{2}. \quad (2)$$

The Lévy area formula has many interesting connotations. For example there are connections to integrable systems, Bernoulli and Euler polynomials, and hence to the values of the Riemann zeta function [31]. For some recent work and further references see [21, 22]. Also, to within normalisation and rescaling it is equal to its Fourier transform, the density of the corresponding probability distribution, which is a boundary point of the Meixner family [24].

Noncommutative analogues of Lévy area have been previously considered in free probability [4, 25, 29]. Also in this connection Deya and Schott [7] emphasise the primacy of iterated stochastic integrals which accords with our philosophy. But in this paper we are concerned with a noncommutative analogue of a more conservative kind which arises in quantum stochastic calculus [13, 26], regarded as a noncommutative extension, rather than a radically noncommutative analogue, of Itô calculus. This allows a very natural variant of the area to be constructed using the minimal one-dimensional version of the calculus. It may be regarded as a response to the call [1] for a study in this quantum context of some of the deeper properties of Brownian motion, as well as a furtherance of the theory of quantum stochastic product integrals [17, 18, 12].

By replacing B^1 and B^2 with P and Q , the iterated quantum stochastic integral

$$K(t) = \frac{1}{2} \int_{0 \leq x < y < t} (dP_x dQ_y - dQ_x dP_y)$$

has some interesting properties [19, 6]. For example it is evidently invariant under gauge transformations, which replace (P, Q) by (P^θ, Q^θ) where

$$P^\theta = P \cos \theta - Q \sin \theta, \quad Q^\theta = P \sin \theta + Q \cos \theta;$$

equivalently the corresponding creation and annihilation processes are multiplied by $e^{\pm i\theta}$. In particular, taking $\theta = -\frac{\pi}{2}$ it is invariant under the replacement (P, Q)

by $(Q, -P)$. Thus, unlike the separate processes P and Q , it can be canonically “rolled” onto a (one-dimensional) Riemannian manifold, and its multidimensional version [8] can similarly be rolled onto a multidimensional manifold, with possible applications to quantum stochastic proofs of index theorems (see for example [16, 3]), by identifying the canonical Brownian motion on the manifold generated by the Laplacian as P^θ with arbitrarily chosen θ .

It can also be verified [6] that all moments of $K(t)$ vanishes in the vacuum state, so that $K(t)$ vanishes in a probabilistic sense, even though it is not the zero operator.

But it is not K which is the main object of study. Because $\exp(a + b) \neq \exp a \exp b$ when a and b do not commute, the exponential

$$\exp(i\lambda K(t)) = \exp\left(\frac{i\lambda}{2} \int_{0 \leq x < y < t} (dP_x dQ_y - dQ_x dP_y)\right)$$

does not reflect in a coherent way the continuous tensor product structure underlying the quantum stochastic calculus. Thus, motivated by the hope of finding quantum extensions of, in particular, the Lévy area formula (2), and associated relations with Euler and Bernoulli polynomials [22] we investigate the double product integral

$$\prod_{<[a,b]} (1 + dr_1) = \prod_{a \leq x < y < b} \left(1 + \frac{i\lambda}{2} (dP_x dQ_y - dQ_x dP_y)\right).$$

However, as it turns out, the more general object

$$\begin{aligned} E &= \prod_{<[a,b]} (1 + dr_{\lambda,\mu}) \\ &= \prod_{a \leq x < y < b} \left(1 + \frac{i\lambda}{2} (dP_x dQ_y - dQ_x dP_y) + \frac{i\mu}{2} (dP_x dP_y + dQ_x dQ_y)\right). \end{aligned}$$

is more fundamental and, surprisingly, simpler to study.

3. A discrete double product of unitary matrices

The first stage of the construction of E is similar to that of the rectangular case construction outlined in [15], in that we approximate $\prod_{<[a,b]} (1 + dr_{\lambda,\mu})$ by a discrete double product $\prod_{1 \leq j < k \leq N} (I + \delta_N^{j,k} r_{\lambda,\mu})$, where $\delta_N^{j,k} r_{\lambda,\mu}$ is obtained from $dr_{\lambda,\mu}$ by replacing each basic differential $dX \in \{dP, dQ\}$ contributing to

$dr_{\lambda,\mu} \in \mathcal{J} \otimes \mathcal{J}$ in the first copy of \mathcal{J} by the j -th increment $X_{x_j} - X_{x_{j-1}}$ and in the second copy of \mathcal{J} by the k -th increment $X_{x_k} - X_{x_{k-1}}$ over the equipartition

$$[a, b) = \bigsqcup_{j=1}^N [x_{j-1}, x_j), \quad x_j = a + \frac{j}{N}(b - a) =: a + j\Delta_N.$$

Thus, for example,

$$\begin{aligned} \delta_N^{j,k} r_1 &= \frac{i}{2} ((P_{x_j} - P_{x_{j-1}})(Q_{x_k} - Q_{x_{k-1}}) \\ &\quad - (Q_{x_j} - Q_{x_{j-1}})(P_{x_k} - P_{x_{k-1}})). \end{aligned}$$

Introducing the standard canonical pairs (p_j, q_j) , $j = 1, 2, \dots, N$, given by

$$\begin{aligned} p_j &= \sqrt{\frac{N}{b-a}} (P(x_j) - P(x_{j-1})), \\ q_j &= \sqrt{\frac{N}{b-a}} (Q(x_j) - Q(x_{j-1})), \end{aligned}$$

which satisfy the canonical commutation relations

$$[p_j, q_k] = -2i\delta_{j,k}, \quad [p_j, p_k] = [q_j, q_k] = 0, \quad (3)$$

we write

$$\delta_N^{j,k} r_1 = i \frac{b-a}{2N} (p_j q_k - q_j p_k)$$

and more generally

$$\delta_N^{j,k} r_{\lambda,\mu} = i \frac{b-a}{2N} (\lambda (p_j q_k - q_j p_k) + \mu (p_j p_k + q_j q_k)).$$

Our approximation is thus

$$\begin{aligned} &\prod_{<[a,b)} (1 + dr_{\lambda,\mu}) \\ &\simeq \prod_{1 \leq j < k \leq N} \left(I + i \frac{b-a}{2N} (\lambda (p_j q_k - q_j p_k) + \mu (p_j p_k + q_j q_k)) \right) \quad (4) \\ &\simeq \prod_{1 \leq j < k \leq N} \exp \left(i \frac{b-a}{2N} (\lambda (p_j q_k - q_j p_k) + \mu (p_j p_k + q_j q_k)) \right) \end{aligned}$$

for large N .

Temporarily let us fix $j < k$ and write $(p, q) = (p_j, q_j)$, $(p', q') = (p_k, q_k)$ so that

$$[p, q] = -2i, \quad [p', q'] = -2i, \quad [p, q'] = [q, p'] = [p, p'] = [q, q'] = 0. \quad (5)$$

We recall [26] that, for an arbitrary Hilbert space \mathcal{H} and vector $f \in \mathcal{H}$ the corresponding *Weyl operator* $W(f)$ is the unique unitary operator on $\mathcal{F}(\mathcal{H})$ which acts on each exponential vector $e(g)$, $g \in \mathcal{H}$ as

$$W(f)e(g) = e^{-\frac{1}{2}\|f\|^2 - (f,g)}e(f+g).$$

The Weyl operators satisfy the Weyl relation

$$W(f)W(g) = e^{-i\text{Im}(f,g)}W(f+g). \tag{6}$$

A convenient rigorous realisation of two canonical pairs satisfying the commutation relations (5) can be constructed in terms of the one-parameter unitary groups of which they are the self-adjoint infinitesimal generators, which are Weyl operators on the Fock space $\mathcal{F}(\mathbb{C}^2)$ over \mathbb{C}^2 . Regarding \mathbb{C}^2 as a space of column vectors, we take

$$\begin{aligned} e^{ixp} &= W((x, 0)^\tau), & e^{ixq} &= W((-ix, 0)^\tau), \\ e^{ixp'} &= W((0, x)^\tau), & e^{ixq'} &= W((0, -ix)^\tau). \end{aligned}$$

for arbitrary $x \in \mathbb{R}$, noting that these four families of Weyl operators are indeed one-parameter unitary groups, and that the commutation relations (5) follow by parametric differentiation, for example from the relations

$$\begin{aligned} W((x, 0)^\tau)W((-iy, 0)^\tau) &= e^{2ixy}W((-iy, 0)^\tau)W((x, 0)^\tau), \\ W((0, x)^\tau)W((0, -iy)^\tau) &= e^{2ixy}W((0, -iy)^\tau)W((0, x)^\tau), \end{aligned}$$

all of which are consequences of (6).

Theorem 1 below, which was proved in [15, Theorem 2], gives a corresponding rigorous explicit form of the self-adjoint operator

$$L(\lambda, \mu) = \lambda(pq' - qp') + \mu(pp' + qq')$$

in this realisation. Before stating it we recall [26] that the *second quantisation* of a unitary operator U on a Hilbert space \mathcal{H} is the unique unitary operator $\Gamma(U)$ on $\mathcal{F}(\mathcal{H})$ which acts on the exponential vectors as

$$\Gamma(U)e(f) = e(Uf).$$

It is related to the Weyl operators by

$$\Gamma(U)W(f) = W(Uf)\Gamma(U) \tag{7}$$

for arbitrary $f \in \mathcal{H}$. Second quantisation is multiplicative, in the sense that

$$\Gamma(U_1U_2) = \Gamma(U_1)\Gamma(U_2) \tag{8}$$

for arbitrary unitary U_1, U_2 .

Theorem 1 (Theorem 2 in [15]). $L(\lambda, \mu)$ generates the one-parameter unitary group

$$e^{ixL(\lambda, \mu)} = \Gamma \left(\begin{bmatrix} \cos(2x|v|) & -e^{-i\phi} \sin(2x|v|) \\ e^{i\phi} \sin(2x|v|) & \cos(2x|v|) \end{bmatrix} \right), \quad x \in \mathbb{R}$$

where $v = \lambda + i\mu = e^{i\phi} |v|$ and the matrix operates on column vectors in \mathbb{C}^2 by multiplication on the left.

We now use Theorem 1 to construct an explicit form of the approximation (4).

Let us first construct a different realisation of the canonical pairs (p_j, q_j) , $j = 1, 2, \dots, n$, satisfying (3) in the Fock space $\mathcal{F}(\mathbb{C}^n)$ over \mathbb{C}^n , by defining

$$e^{ixp_j} = W(x\varepsilon_j), \quad e^{ixq_j} = W(-ix\varepsilon_j)$$

where $(\varepsilon_j)_{j=1}^n$ is the standard orthonormal basis of \mathbb{C}^n , $\varepsilon_j = (0, \dots, \overset{(j)}{1}, 0, \dots, 0)^\tau$. Correspondingly, in view of Theorem 1, each operator

$$\exp \left(i \frac{b-a}{2N} (\lambda(p_j q_k - q_j p_k) + \mu(p_j p_k + q_j q_k)) \right)$$

is realised as the second quantisation $\Gamma(R_{j,k}^N)$ where

$$R_{j,k}^N := \begin{bmatrix} & & \overset{(j)}{0} & \dots & \overset{(k)}{0} & \dots & 0 \\ & 1 & \dots & & \dots & & \dots \\ \vdots & \vdots & \ddots & & \vdots & & \vdots \\ \overset{(j)}{0} & \dots & \cos \left(\frac{(b-a)}{N} |v| \right) & \dots & -\frac{\bar{v}}{|v|} \sin \left(\frac{(b-a)}{N} |v| \right) & \dots & 0 \\ \vdots & \dots & \vdots & \ddots & \vdots & \dots & \vdots \\ \overset{(k)}{0} & \dots & \frac{\bar{v}}{|v|} \sin \left(\frac{(b-a)}{N} |v| \right) & \dots & \cos \left(\frac{(b-a)}{N} |v| \right) & \dots & 0 \\ \vdots & \dots & \vdots & \dots & \vdots & \ddots & \dots \\ 0 & \dots & 0 & \dots & 0 & \dots & 1 \end{bmatrix}.$$

In view of the multiplicativity property (8) the discrete double product (4) is correspondingly realised as the second quantisation of the product

$$\prod_{1 \leq j < k \leq N} R_{j,k}^N. \tag{9}$$

We now embed the matrix (9) as a unitary operator \mathcal{W}_N on $L^2([a, b])$ by mapping the standard basis of \mathbb{C}^N to the orthonormal family $(\chi_1, \chi_2, \dots, \chi_N)$ of normalized indicator functions

$$\chi_j(x) = \sqrt{\frac{N}{b-a}} \mathbb{1}_{[x_{j-1}, x_j]}.$$

By definition \mathcal{W}_N acts as the identity operator I on $(\chi_1, \chi_2, \dots, \chi_N)^\perp$.

Our objective in the remainder of this paper is to find an explicit form for the (weak) limit

$$W = \lim_{N \rightarrow \infty} \mathcal{W}_N$$

and to prove that W is unitarity. The corresponding problems for rectangular unitary product integrals was solved in outline in [15]. The causal case considered here is considerably more difficult, because the method of iterated limits which reduces the rectangular case to a double application of the time-orthogonal unitary dilation of [10], is not applicable. Instead a combinatorial argument based on a lattice path model is used. For a similar alternative approach, avoiding the iterated limit technique, to the rectangular product in the particular case of the generator dr_1 corresponding to the quantum Lévy area, see [11]; however the combinatorics for the rectangular case is much simpler than here and it has no direct relation to Lévy area.

4. A lattice path model and linear extensions of partial orderings

So we want to calculate the limit of the triangular double product of $N \times N$ matrices

$$\mathcal{W}_N = \prod_{1 \leq j < k \leq N} R_{j,k}^N. \tag{10}$$

Here, for elements $x_{j,k}$ of an associative algebra having the property that $x_{j,k}$ commutes with $x_{j',k'}$ whenever $\{j, k\} \cap \{j', k'\} = \emptyset$ we define the ordered double product $\prod_{1 \leq j < k \leq N} x_{j,k}$ by any of the equivalent prescriptions

$$\prod_{1 \leq j < k \leq N} x_{j,k} = \prod_{j=1}^{N-1} \left[\prod_{k=j+1}^N x_{j,k} \right] = \prod_{k=2}^N \left[\prod_{j=1}^{k-1} x_{j,k} \right] = \prod_{r=1}^{\frac{1}{2}N(N-1)} x_{j_r, k_r}$$

where

$$((j_1, k_1), (j_2, k_2), \dots, (j_{\frac{1}{2}N(N-1)}, k_{\frac{1}{2}N(N-1)}))$$

is any ordering of the $\frac{1}{2}N(N - 1)$ pairs (j, k) , $1 \leq j < k \leq N$ which is *allowed*, meaning that it has the property that

$$(j_r, k_r) \text{ precedes } (j_s, k_s) \text{ if both } j_r \leq j_s \text{ and } k_r \leq k_s. \tag{11}$$

In constructing the limit as $N \rightarrow \infty$ we use the small angle approximations for sine and cosine, so that

$$\begin{pmatrix} \cos \frac{b-a}{N} |\nu| & -\frac{\bar{\nu}}{|\nu|} \sin \frac{b-a}{N} |\nu| \\ \frac{\nu}{|\nu|} \sin \frac{b-a}{N} |\nu| & \cos \frac{b-a}{N} |\nu| \end{pmatrix} = I + \frac{b-a}{N} \begin{pmatrix} 0 & -\bar{\nu} \\ \nu & 0 \end{pmatrix} + O(N^{-2})$$

hence

$$R_{j,k}^N = I + \frac{(b-a)}{N} (-\bar{\nu} |\chi_j\rangle \langle \chi_k| + \nu |\chi_k\rangle \langle \chi_j|) + O(N^{-2}).$$

When there is no ambiguity, for any integers j and k , we use abbreviations

$$|j\rangle := |\chi_j\rangle \quad \text{and} \quad \langle k| := \langle \chi_k|.$$

Then the product (10) becomes

$$W_N \simeq \prod_{1 \leq j < k \leq N} \left(I + \frac{(b-a)}{N} Z^N(j, k) \right) =: W_N$$

where

$$Z^N(j, k) = -\bar{\nu} |j\rangle \langle k| + \nu |k\rangle \langle j|.$$

To compute this, we introduce and work on a lattice path model. Consider a lattice $L_s := \{(m, n) : 1 \leq m \leq s, 0 \leq n \leq 1\}$. We call $(m, 1)_{1 \leq m \leq s}$ the upper vertices, and $(m, 0)_{1 \leq m \leq s}$ the lower vertices. Denote by Π_s the set of lattice path $\pi = (m_i, b_i)_{i=1}^s$ satisfying the following two conditions:

- (1) $m_i = i$ for $i = 1, \dots, s$;
- (2) there does not exist an i such that $b_i = b_{i+1} = 0$.

For convenience, we write $\pi(i) = b_i$ and let $\pi = (\pi(i))_i$. We call any $\pi \in \Pi_s$ a path of length $s - 1$. It is straightforward to verify by induction that

$$|\Pi_s| = \text{Fib}_{s+2} = \frac{\Phi^{s+2} - (-\Phi)^{-s-2}}{\sqrt{5}},$$

where Fib_n is the n th Fibonacci number and Φ is the golden ratio $\frac{\sqrt{5}+1}{2}$.

If we assign weight $\theta(v)$ to each vertex v in L_s , then we can define the weight $\theta(\pi)$ of a path $\pi \in \Pi_s$ by the product of the weights of its vertices:

$$\theta(\pi) := \prod_{i=1}^s \theta(i, \pi(i)).$$

For any array of s pairs $\{p_{ij}: 1 \leq i \leq s, 1 \leq j \leq 2\}$, define its associated weight $\theta_p(v)$ for any $v = (m, b) \in L_s$ to be

$$\theta_p(v) = \begin{cases} v |p_{i2}\rangle \langle p_{i1}| & \text{if } b = 0; \\ -\bar{v} |p_{i1}\rangle \langle p_{i2}| & \text{if } b = 1. \end{cases}$$

Finally, define the weight $\theta_p(\pi)$ of a path in the same way as before.

For example, if we label the vertices by their weights associated to p , the path in Figure 1 is a path in Π_5 , but not the path in Figure 2 because the third edge connects two bottom vertices.

Any array of s pairs $p = (p_{ij})_{1 \leq i \leq s, 1 \leq j \leq 2}$ satisfying the condition

$$\mathbb{I}_{p_{i,1}=p_{i+1,1}} + \mathbb{I}_{p_{i,2}=p_{i+1,2}} + \mathbb{I}_{p_{i,2}=p_{i+1,1}} = 1, \quad p_{i,1} \neq p_{i+1,2} \quad (12)$$

can be associated with a path $\pi_p \in \Pi_s$ in the following way:

$$(\pi(i), \pi(i + 1)) = \begin{cases} (0, 1) & \text{if } p_{i,1} = p_{i+1,1}, \\ (1, 1) & \text{if } p_{i,2} = p_{i+1,1}, \\ (1, 0) & \text{if } p_{i,2} = p_{i+1,2}. \end{cases}$$

Note that this is equivalent to

$$\theta_p(\pi_p) = \prod_{i=1}^s Z^N(p_{i,1}, p_{i,2}).$$

Lemma 2. *We have*

$$W_N = I + \sum_{s=1}^{N(N-1)/2} \tilde{w}_{s,N},$$

where

$$\tilde{w}_{s,N} = \left(\frac{b-a}{N}\right)^s \sum_{(*)} \theta_p(\pi_p)$$

where the domain $(*)$ of the summation is

$$\begin{aligned} & (1 \leq p_{i,j} \leq N, \text{ for all } 1 \leq i \leq s, 1 \leq j \leq 2) \text{ and} \\ & (p_{i,1} < p_{i+1,1} < p_{i,2} = p_{i+1,2} \text{ or} \\ & p_{i,1} < p_{i,2} = p_{i+1,1} < p_{i+1,2} \text{ or} \\ & p_{i,1} = p_{i+1,1} < p_{i,2} < p_{i+1,2}, \text{ for all } 1 \leq i \leq s - 1). \end{aligned}$$

$$\begin{array}{l}
 -\bar{v} |p_{11}\rangle \langle p_{12}| - -\bar{v} |p_{21}\rangle \langle p_{22}| -\bar{v} |p_{31}\rangle \langle p_{32}| -\bar{v} |p_{4,1}\rangle \langle p_{4,2}| - -\bar{v} |p_{5,1}\rangle \langle p_{5,2}| \\
 +v |p_{12}\rangle \langle p_{11}| +v |p_{22}\rangle \langle p_{21}| +v |p_{32}\rangle \langle p_{31}| +v |p_{4,2}\rangle \langle p_{4,1}| +v |p_{5,2}\rangle \langle p_{5,1}|
 \end{array}$$

Figure 1

$$\begin{array}{l}
 -\bar{v} |p_{11}\rangle \langle p_{12}| - -\bar{v} |p_{21}\rangle \langle p_{22}| -\bar{v} |p_{31}\rangle \langle p_{32}| -\bar{v} |p_{4,1}\rangle \langle p_{4,2}| -\bar{v} |p_{5,1}\rangle \langle p_{5,2}| \\
 +v |p_{12}\rangle \langle p_{11}| +v |p_{22}\rangle \langle p_{21}| +v |p_{32}\rangle \langle p_{31}| - +v |p_{4,2}\rangle \langle p_{4,1}| +v |p_{5,2}\rangle \langle p_{5,1}|
 \end{array}$$

Figure 2

Proof. For any rearrangement $(j_i, k_i)_{1 \leq i \leq \frac{N(N-1)}{2}}$ of $\{1 \leq j < k \leq N\}$ satisfying (11),

$$\begin{aligned} \prod_{1 \leq j < k \leq N} (I + Z(j, k)) &= \prod_{i=1}^{N(N-1)/2} (I + Z(j_i, k_i)) \\ &= I + \sum_{s=1}^{N(N-1)/2} \sum_{(**)} \prod_{r=1}^s Z(p_{r1}, p_{r2}), \end{aligned}$$

where the domain (**) is

$$(p_{11}, p_{12}), (p_{21}, p_{22}), \dots, (p_{s1}, p_{s2})$$

is a subsequence of $(j_1, k_1), (j_2, k_2), \dots, (j_{N(N-1)/2}, k_{N(N-1)/2})$.

Now for the product $\prod_{r=1}^s Z(p_{r1}, p_{r2})$ to be nonzero, the product of

$$Z(p_{i,1}, p_{i,2}) \times Z(p_{i+1,1}, p_{i+1,2})$$

for each i has to be nonzero, that is

$$(-|p_{i,1}\rangle \langle p_{i,2}| + |p_{i,2}\rangle \langle p_{i,1}|)(-|p_{i+1,1}\rangle \langle p_{i+1,2}| + |p_{i+1,2}\rangle \langle p_{i+1,1}|) \neq 0.$$




This in turn is equivalent to

$$(p_{i,2} = p_{i+1,2}) \text{ or } (p_{i,2} = p_{i+1,1}) \text{ or } (p_{i,1} = p_{i+1,1}) \text{ or } (p_{i,1} = p_{i+1,2}).$$

We analyse these four possibilities one by one.

- (1) If $p_{i,2} = p_{i+1,2}$, then by (11), and since $(p_{i,1}, p_{i,2}) \neq (p_{i+1,1}, p_{i+1,2})$, only when $p_{i,1} < p_{i+1,1}$ can the product be nonzero. In this case the coordinates are ordered as $p_{i,1} < p_{i+1,1} < p_{i,2} = p_{i+1,2}$.
- (2) If $p_{i,2} = p_{i+1,1}$, then since $p_{i,1} < p_{i,2} = p_{i+1,1}$ and $p_{i,2} = p_{i+1,1} < p_{i+1,2}$, we have that (11) is satisfied. Therefore this case is also included / permitted in the product. The ordering of the coordinates is $p_{i,1} < p_{i,2} = p_{i+1,1} < p_{i+1,2}$.
- (3) If $p_{i,1} = p_{i+1,1}$, then similar to Case 1, the coordinates have to satisfy $p_{i,1} = p_{i+1,1} < p_{i,2} < p_{i+1,2}$ for the product to be nonzero.
- (4) If $p_{i,1} = p_{i+1,2}$, then $p_{i,1} = p_{i+1,2} > p_{i,1}$ and $p_{i,2} > p_{i,1} = p_{i+1,2}$ violates (11), hence this case never happens.

The three feasible cases are illustrated as below.

$i \quad i + 1$	$i \quad i + 1$	$i \quad i + 1$	(13)
			
$p_{i1} < p_{i+1,1}$	$p_{i1} < p_{i+1,1}$	$p_{i1} = p_{i+1,1}$	
$\wedge \quad \wedge$	$\wedge \not\parallel \wedge$	$\wedge \quad \wedge$	
$p_{i2} = p_{i+1,2}$	$p_{i2} < p_{i+1,2}$	$p_{i2} < p_{i+1,2}$	

The concatenation of these edges gives a path in Π_s . Case 4 corresponds to a horizontal bottom edge in the path which is not allowed in the definition of Π_s . Therefore we have established a correspondence between the possibilities of orderings in the product and Π_s . \square

Denote by A_s^* the set of array of s pairs p satisfying condition (*), and $\Omega_\pi := \{p \in A_s^* : \pi_p = \pi\}$. Then

$$\tilde{w}_{s,N} = \left(\frac{b-a}{N}\right)^s \sum_{\pi \in \Pi_s} \sum_{p \in \Omega_\pi} \theta_p(\pi).$$

Given a path $\pi \in \Pi_s$, by the correspondence in (13) there exist $m_1, m_2, \dots, m_{s-1} \in \{1, 2\}$ such that for any $p \in \Omega_\pi$, $p_{x,m_x} = p_{x+1,m'_{x+1}}$, where $m'_x := 3 - m_x$. Therefore, Ω_π is characterised by a partial ordering on the $s + 1$ coordinates $p_{1,m'_1}, p_{1,m_1}, p_{2,m_2}, \dots, p_{s-1,m_{s-1}}, p_{s,m'_{s-1}}$. We call them the *essential coordinates* of p . This also shows we can associate π with $(m_1, m_2, \dots, m_{s-1})$. In the following we do not differentiate between π and the corresponding partial ordering.

Any ordering $\pi \in \Pi_s$ can be decomposed into (strict) total orderings without any repetition of the essential coordinates and those with repeated essential coordinates. We call any of the former orderings B a *linear extension* of π which is denoted by $B \vdash \pi$, and the latter degenerate orderings, which, for reasons that will emerge in the proof of Lemma 3, are ignored. Thus we have

$$\Omega_\pi = \bigsqcup_{B \vdash \pi} B \cup \text{set of degenerate orderings.}$$

For any $p \in \Omega_\pi$, there exists a $B \vdash \pi$ such that $p \in B$. Denote $(j_p, k_p) = (p_{1,m'_1}, p_{s,m'_{s-1}})$. In the total ordering imposed by B , let r_B be the number of essential coordinates less than j_p and r'_B the number of those greater than k_p . That is, the essential coordinates are ordered as follows.

- If $r_B + r'_B < s$,

$$\begin{aligned}
 &1 < l_1 < \dots \\
 &< l_{r_B} < j_p < m_1 < \dots \\
 &< m_{s-1-r_B-r'_B} < k_p < n_1 < \dots \\
 &< n_{r'_B} \leq N.
 \end{aligned}$$

- If $r_B + r'_B > s$,

$$\begin{aligned}
 &1 < l_1 < \dots \\
 &< l_{s-r'_B} < k_p < m_1 < \dots \\
 &< m_{r_B+r'_B-s-1} \\
 &< j_p < n_1 < \dots \\
 &< n_{s-r_B} \leq N.
 \end{aligned}$$

We call (r_B, r'_B) the rank of B .

Let $\epsilon(\pi)$ be the number of upper vertices of the path π . Since horizontal edges between lower vertices are not allowed, there is at least one upper vertex between two consecutive lower vertices, hence

$$2\epsilon(\pi) \geq s - 1.$$

The location (upper or lower) of the first vertex of π , the number of upper vertices $\epsilon(\pi)$ and the parity of the length of π together determine the number of horizontal edge in π . An extreme case is when there is no horizontal edges in π , for which $\epsilon(\pi) \approx \frac{s-1}{2}$. This will be later specified and exploited in the proof of Lemma 12.

The weight of π is

$$\theta_p(\pi) = (-\bar{\nu})^{\epsilon(\pi)} \nu^{s-\epsilon(\pi)} |j_p\rangle \langle k_p|.$$


So

$$\begin{aligned}
 \tilde{w}_{s,N} &= \left(\frac{b-a}{N}\right)^s \sum_{\pi \in \Pi_s} (-\bar{\nu})^{\epsilon(\pi)} \nu^{s-\epsilon(\pi)} \sum_{p \in \Omega_\pi} |j_p\rangle \langle k_p| \\
 &\simeq \left(\frac{b-a}{N}\right)^s \sum_{\pi \in \Pi_s} (-\bar{\nu})^{\epsilon(\pi)} \nu^{s-\epsilon(\pi)} \sum_{B \vdash \Omega_\pi} \sum_{p \in B} |j_p\rangle \langle k_p| \\
 &= \sum_{\pi \in \Pi_s} (-\bar{\nu})^{\epsilon(\pi)} \nu^{s-\epsilon(\pi)} \sum_{B \vdash \pi} H_s^N(r_B, r'_B) + v_{s,N} =: w_{s,N} + v_{s,N},
 \end{aligned}$$

where $v_{s,N}$ is the contribution from the degenerate orderings, on which one can carry out the same calculation for $w_{s,N}$ below, and

$$H_s^N(r, r') = \begin{cases} \left(\frac{b-a}{N}\right)^s \sum_{1 < l_1 < \dots < l_r < j < m_1 < \dots < m_{s-1-r-r'} < k < n_1 < \dots < n_{r'} \leq N} |j\rangle \langle k| & \text{if } r + r' < s, \\ \left(\frac{b-a}{N}\right)^s \sum_{1 < l_1 < \dots < l_{s-r'} < k < m_1 < \dots < m_{r+r'-s-1} < j < n_1 < \dots < n_{s-r} \leq N} |j\rangle \langle k| & \text{if } r + r' > s. \end{cases}$$

For example, for the following path π of length 2,



$$\begin{array}{ccc} p_{1,1} = p_{2,1} < p_{3,1} \\ \wedge \quad \wedge \quad \wedge \\ p_{1,2} < p_{2,2} = p_{3,2} \end{array}$$

we have $m_1 = 1, m_2 = 2, m_3 = 1$. The ordering of the essential coordinates imposed by π is:

$$(p_{1,1} < p_{1,2} < p_{2,2}) \wedge (p_{2,1} < p_{3,1} < p_{2,2}),$$

and the non-repeated starting and ending coordinates are $j_p = p_{12}$ and $k_p = p_{31}$. The total ordering decomposition of Ω_π is

$$\begin{aligned} \Omega_\pi &= \{p_{11} < p_{12} < p_{31} < p_{22}\} \sqcup \{p_{11} < p_{31} < p_{12} < p_{22}\} \\ &\quad \sqcup \{p_{11} < p_{12} = p_{31} < p_{22}\}. \end{aligned}$$

The last term is a degenerate case as p_{12} is repeated. There is only one upper vertex, hence this path contributes $-v|v|^2(H_3^N(1, 1) + H_3^N(2, 2))$ to $w_{s,N}$.

Define the Volterra-type kernels

$$\mathop{>}_a^b(x, y) := 1_{a \leq y < x < b} \quad \text{and} \quad \mathop{<}_a^b(x, y) := \mathop{>}_a^b(y, x),$$

and

$$[m, n, p](x, y) := \frac{(x-a)^m (y-x)^n (b-y)^p}{m! n! p!},$$

and

$$[m, n, p]^\dagger(x, y) := [m, n, p](y, x).$$

The asymptotics of H_s^n can be written down explicitly.

Lemma 3. $H_s^N(r, r')$ converges weakly to an integral operator $H_s(r, r')$, with the integral kernel

$$h_s(r, r') = \begin{cases} [r, s - 1 - r - r', r'] <_a^b & \text{if } r + r' < s, \\ [s - r', r + r' - s - 1, s - r]^\dagger >_a^b & \text{if } r + r' > s. \end{cases}$$

Proof. Suppose $r + r' < s$ (the case $r + r' > s$ can be done in the same way). Then

$$\begin{aligned} H_s^N(r, r') &= \left(\frac{b-a}{N}\right)^s \sum_{1 \leq j < k \leq N} |j\rangle \langle k| \sum_{1 \leq l_1 < \dots < l_r < j < m_1 < \dots < m_{s-1-r-r'} < k < n_1 < \dots < n_{r'} \leq N} 1 \\ &= \left(\frac{b-a}{N}\right)^s \sum_{1 \leq j < k \leq N} |j\rangle \langle k| \binom{j-1}{r} \binom{k-j-1}{s-1-r-r'} \binom{N-k}{r'} \end{aligned}$$

We denote

$$\Delta_N := \frac{b-a}{N} \quad \text{and} \quad A_j := [x_{j-1}, x_j),$$

then the kernel of $H_s^N(r, r')$ is

$$\begin{aligned} h_s^N(x, y) &= \sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y) \frac{1}{r!(s-1-r-r')!r'!} \\ &\quad \prod_{\alpha=0}^{r-1} (x_{j-1} - a - \alpha\Delta_N) \prod_{\beta=0}^{s-2-r-r'} (x_{k-1} - x_j - \beta\Delta_N) \\ &\quad \prod_{\gamma=0}^{r'-1} (b - x_k - \gamma\Delta_N). \end{aligned}$$

This, as $N \rightarrow \infty$, converges weakly (as an integral kernel) to $[r, s - 1 - r - r', r'] <_a^b(x, y)$. □

It can also be seen from the proof of this lemma that the degenerate orderings contribute 0 to the total sum. More specifically, the degenerate version of $h_s^N(x, y)$ where there are d repeated essential coordinates is

$$\begin{aligned} q_s^N(x, y) &= \sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y) \frac{1}{r!(s-1-r-r')!r'!} \frac{(b-a)^d}{N^d} \\ &\quad \prod_{\alpha=0}^{r-1} (x_{j-1} - a - \alpha\Delta_N) \prod_{\beta=0}^{s-d-2-r-r'} (x_{k-1} - x_j - \beta\Delta_N) \\ &\quad \prod_{\gamma=0}^{r'-1} (b - x_k - \gamma\Delta_N) \longrightarrow 0 \end{aligned}$$

as $N \rightarrow \infty$. We will examine carefully the rate of convergence of this lemma and the (in)significance of the degenerate orderings later in the proof of Theorem 7. This lemma immediately gives the following corollary.

Corollary 4. *There exist two integer arrays $(D_{m,n,p;q})_{m,n,p \geq 0, 0 \leq q \leq m+n+p+1}$ and $(E_{m,n,p;q})_{m,n,p \geq 0, 0 \leq q \leq m+n+p+1}$ such that $w_{s,N}$ converges weakly as $N \rightarrow \infty$ to w_s with kernel*

$$f_s <_a^b + g_s >_a^b$$

where f_s and g_s are defined by

$$f_s = \sum_{m,n,p \geq 0, m+n+p=s-1} \sum_{q=0}^s D_{m,n,p;q} (-\bar{v})^q v^{s-q} [m, n, p],$$

$$g_s = \sum_{m,n,p \geq 0, m+n+p=s-1} \sum_{q=0}^s E_{m,n,p;q} (-\bar{v})^q v^{s-q} [m, n, p]^\dagger.$$

Indeed, $D_{m,n,p;q}$ (resp. $E_{m,n,p;q}$) enumerates the linear extensions of all possible paths of length $m + n + p$ with q upper vertices and rank (m, p) (resp. $(m + n + 1, n + p + 1)$).

Corollary 5. *The functions f_s and g_s both are symmetric in the following sense:*

$$f_s(x, y) = f_s(a + b - y, a + b - x), \quad g_s(x, y) = g_s(a + b - y, a + b - x)$$

Proof. This follows from the fact that the path inversion $(i, b_i) \mapsto (i, b_{s+1-i})$ is a weight-preserving bijection between Π_s and itself. \square

For example, some calculation yields

$$f_1 = -\bar{v}[0, 0, 0], \tag{14a}$$

$$f_2 = -|v|^2[0, 0, 1] - |v|^2[1, 0, 0] + \bar{v}^2[0, 1, 0], \tag{14b}$$

$$f_3 = (\bar{v}|v|^2 - \bar{v}^3)[0, 2, 0] + \bar{v}|v|^2[0, 1, 1] + \bar{v}|v|^2[1, 1, 0] - v|v|^2[1, 0, 1], \tag{14c}$$

$$g_1 = v[0, 0, 0]^\dagger, \tag{14d}$$

$$g_2 = 0, \tag{14e}$$

$$g_3 = -v|v|^2[1, 0, 1]^\dagger. \tag{14f}$$

The following three theorems are the main results of this paper.

Theorem 6. *The closed form expression of D and E are*

$$D_{m,n,p;q} = \begin{cases} \binom{n}{q-1} - \binom{n}{q} & \text{if } 2q > m + n + p, \\ \binom{n}{q-m} - \binom{n}{q} & \text{if } 2q = m + n + p, \\ 0 & \text{if } 2q < m + n + p, \end{cases} \tag{15}$$

$$E_{m,n,p;q} = \mathbb{I}_{m=p=q,n=0}.$$

Proof. See Section 5. □

Theorem 7. *Recall the definition of w_s in Corollary 4. The operator W_N converges weakly to*

$$W = I + \sum_{s \geq 1} w_s.$$

Proof. See Section 6. □

For $j \geq 0$, let B_j be power series in two variables related to the Bessel functions of the first kind J_j .

$$B_j(x, y) := \sum_{n \geq 0} \frac{(-1)^{n+j} x^{n+j} y^n}{(n+j)!n!} = (-1)^j (x/y)^{j/2} J_j(2\sqrt{xy}).$$

Let I be the identity, then the kernel of the operator $W - I$ can be written in terms of B_j .

Theorem 8. *The integral operator $W - I$ has kernel*

$$\begin{aligned} \ker(W - I)(x, y) &= (vB_0((y-a)|v|, (b-x)|v|) + |v|B_1((b-a)|v|, (y-x)|v|) \\ &\quad - (v + \bar{v}) \sum_{q \geq 0} B_q((y-x)|v|, (b-a)|v|) \left(\frac{\bar{v}}{|v|}\right)^q) <_a^b(x, y) \\ &\quad + vB_0((y-a)|v|, (b-x)|v|) >_a^b(x, y). \end{aligned}$$

Moreover, W is unitary.

Proof. See section 7. □

For example, when $\mu = 0$ and $\lambda > 0$, the kernel of the operator corresponding to the Lévy stochastic area is

$$\begin{aligned} \ker(W - I)(x, y) &= \left(\lambda B_0((y - a)\lambda, (b - x)\lambda) + \lambda B_1((b - a)\lambda, (y - x)\lambda) \right. \\ &\quad \left. - 2\lambda \sum_{q \geq 0} B_q((y - x)\lambda, (b - a)\lambda) \right) <_a^b(x, y) \\ &\quad + \lambda B_0((y - a)\lambda, (b - x)\lambda) >_a^b(x, y). \end{aligned}$$

Moreover by plugging $D_{m,n,p;q}$ and $E_{m,n,p;q}$ into the integral identity (19) below, the unitarity of W implies the following combinatorial identity:

$$\begin{aligned} D_{\alpha,\beta,\gamma;\xi} - \mathbb{I}_{\alpha=\gamma=\xi-1,\beta=0} - \binom{\alpha + \gamma - 1}{\gamma} \mathbb{I}_{\beta+\gamma+1=\alpha=\xi} \\ - \sum_{m=0}^{\alpha} \sum_{p=0}^{\gamma-\alpha+m} \sum_{n=0}^{\alpha+\beta-\gamma-m+p-1} D_{m,n,\alpha+\beta-\gamma-m-n+2p-1;\xi-\gamma+p-1} \\ \binom{\alpha}{m} \binom{\gamma - \alpha + m + n - p}{n} \binom{\gamma}{p} \\ + \sum_{m_1=0}^{\alpha} \sum_{m_2=0}^{\beta} \sum_{n_1=0}^{\gamma-1} \sum_{n_2=0}^{\gamma-1-n_1} \sum_{p_1=0}^{\gamma-1-n_1-n_2} \sum_{t_1=0}^{\xi} (-1)^{\alpha+\gamma-m_1-n_1-p_1+m_2} \\ D_{m_1,n_1+\beta-m_2,p_1;t_1} \\ D_{m_2+\alpha-m_1,n_2,\gamma-1-n_1-n_2-p_1;\alpha+\gamma-\xi-m_1-n_1-p_1+m_2+t_1} \\ \binom{\alpha}{m_1} \binom{\beta}{m_2} \binom{n_1 + n_2}{n_1} \binom{\gamma - 1 - n_1 - n_2}{p_1} \\ = 0. \end{aligned}$$

5. Dyck paths and Catalan numbers

For $m \in \mathbb{Z}_{\geq 0}$ and $n \in \mathbb{Z}$, define the binomial coefficient the usual way

$$\binom{m}{n} := \frac{m!}{n!(m-n)!} \mathbb{I}_{0 \leq n \leq m}.$$

For integers m, n, p define a double generalisation of the Catalan numbers and the Catalan’s triangle

$$C_{m,n,p} := \binom{m+n}{m} - \binom{m+n}{m+p+1}.$$

For $\alpha, m, n, p \in \mathbb{Z}_{\geq 0}$, denote by $T_{\alpha,m,n,p}$ the set of lattice paths $(\rho_i)_{i=0}^{m+n}$ such that $\rho_0 = \alpha, |\rho_i - \rho_{i-1}| = 1, \rho_i \geq -p, \rho_{m+n} = \alpha + m - n$. That is, $T_{\alpha,m,n,p}$ is the set of Dyck paths starting from α , having m up-steps, n down-steps that never cross the line $y = -p$. By the reflection principle we obtain the following lemma, which shows these numbers have a similar combinatorial interpretation to the Catalan numbers.

Lemma 9. *When $m, n, p \geq 0$ and $m - n \geq -p - 1$, $C_{m,n,p} = |T_{0,m,n,p}|$.*

More specifically, the reflection principle we use here identifies the forbidden paths that cross $y = -p$ with a path starting at $-2p - 2$, having $m + p + 1$ up-steps and $n - p - 1$ down-steps, and hence ending at $m - n$. This is done by reflecting the part of the paths preceding the point when it first touches the horizontal line $y = -p - 1$ about the same line. The doubly generalised Catalan numbers have been discussed in e.g. [27].

When $m, n \geq 0$ and $p = 0$, $C_{m,n,0}$ is reduced to the (m, n) th entry in the Catalan triangle (OEIS:A009766) which we denote by $C_{m,n}$; furthermore when $m = n$, $C_{n,n}$ is the n th Catalan number which we denote by C_n .

The following recurrence relation will be useful:

Lemma 10. *If $n \geq 0, m \geq p$ and $m + n + p + 1 \geq 0$, then*

$$\sum_{k=0}^{\lfloor \frac{m+p}{2} \rfloor} C_{k+n,k} C_{m-k,p-k} = C_{m+n+1,p}.$$

Proof. We first show a basic version of this formula is true: for $n \geq 0, m \geq p \geq 0$,

$$\sum_{k=0}^p C_{k+n,k} C_{m-k,p-k} = C_{m+n+1,p}.$$

This can be proved using a combinatorial argument similar to one used to prove the recurrence relation of the Catalan numbers which is a special case of the identity above (note that $C_{p+1,p} = C_{p+1,p+1}$):

$$\sum_{k=0}^p C_k C_{p-k} = C_{p+1}.$$

Define a “stopping time” σ on $T_{0,m+n+1,p,0}$ by

$$\sigma(\rho) = \max\{i \geq 0 : \rho_i = n\}.$$

Then

$$\begin{aligned} C_{m+n+1,p} &= \sum_{k=0}^p |\{\rho \in T_{0,m+n+1,p,0} : \sigma(\rho) = 2k + n\}| \\ &= \sum_{k=0}^p |T_{0,k+n,k,0}| |T_{n+1,m-k,p-k,n+1}| \\ &= \sum_{k=0}^p |T_{0,k+n,k,0}| |T_{0,m-k,p-k,0}| \\ &= \sum_{k=0}^p C_{n+k,k} C_{m-k,p-k}. \end{aligned}$$

If the condition $n \geq 0, m \geq p$ are retained, but $p < 0$ and $m + n + 1 + p \geq 0$, then the LHS is zero because the domain of the summation is empty. The RHS is also zero because

$$\binom{m+n+1+p}{m+n+1} = \binom{m+n+1+p}{m+n+2} = 0.$$

Since $m \geq p$, we have $p \leq \lfloor \frac{m+p}{2} \rfloor$. Moreover, for any $k \in (p, \lfloor \frac{m+p}{2} \rfloor]$,

$$C_{m-k,p-k} = \binom{m+p-2k}{m-k} - \binom{m+p-2k}{m-k+1} = 0.$$

Therefore we can extend the domain of the summation from $0 \leq k \leq p$ to $0 \leq k \leq \lfloor \frac{m+n}{2} \rfloor$. □

Lemma 11. For any $B \vdash \pi \in \Pi_s$, if $\pi(0) = 1$ then $r_B = 0$, and if $\pi(0) = 0$ then $r_B > 0$. If $\pi(s) = 1$ then $r'_B = 0$, and if $\pi(s) = 0$ then $r'_B > 0$.

Proof. We show the claim for $\pi(0)$, as the one for $\pi(s)$ can be deduced from the symmetry property. If $\pi(0) = 1$, then $\pi(1) = 0$ or 1 . If $\pi(1) = 0$ then by the correspondence (13), for any $p \in \Omega_\pi$, the first four coordinates have the ordering $j_p = p_{1,1} < p_{2,1} < p_{1,2} = p_{2,2}$. Since $p_{1,1} \leq p_{k,1}, k \geq 2$, and $p_{1,1} < p_{1,2} \leq p_{k,2}, k \geq 2$. Thus j_p is the smallest (essential) coordinate and $r_B = 0$. If $\pi(1) = 1$ then $j_p = p_{1,1} < p_{1,2} = p_{2,1} < p_{2,2}$ hence it's also the smallest coordinates and $r_B = 0$.

If $\pi(1) = 0$, then $\pi(1) = 1$ and by (13), for any $p \in \Omega_\pi$, the first four coordinates are ordered as $p_{1,1} = p_{2,1} < p_{1,2} = j_p < p_{2,2}$. Hence j_p is greater than at least one other essential coordinate and $r_B > 0$. \square

In some extreme cases the coefficient $D_{m,n,p;q}$ can be calculated directly. We denote by $D_{m,n,p;q}^\pi$ the contribution to $D_{m,n,p;q}$ from path π . In the following we denote by $\setminus, /$ and \wedge the edges in paths as shown in (13). We also denote paths by concatenating them with superscripts representing multiplicities. For example $-\vee^2 \setminus$ represents the paths $-\setminus \setminus \setminus \setminus$.

Lemma 12. *The following case hold.*

Case A. $D_{0,2k,0;k+1} = D_{0,2k,0;k+1}^{\vee^k} = C_k$. Conversely, if $m = 0, 2q = n + p + 2$, then $D_{m,n,p;q} > 0$ only if $p = 0, 2q - 2 = n$.

Case B. $D_{0,n,2k-n+1;k+1} = D_{0,n,2k-n+1,k+1}^{\setminus^k} = C_{k,n-k}$ for $0 \leq n \leq 2k + 1$.

Case C. $D_{2k-n+1,n,0;k+1} = D_{2k-n+1,n,k+1}^{\wedge^k /} = C_{k,n-k}$ for $0 \leq n \leq 2k + 1$.

Case D. $D_{r,2k-r-r',r';k} = D_{r,2k-r-r',r';k}^{\wedge^k} = C_{k-r,k-r',r-1} = C_{k-r',k-r,r'-1}$ for $0 \leq r + r' \leq 2k$.

Moreover, $E_{m,n,p;q} = \mathbb{I}_{m=p=q,n=0}$.

Proof. First we show the first identity in each case. In Case D, for there to be k upper vertices and $k + 1$ lower vertices, the path can only be \wedge^k .

For Case A, since the rank is $(0, 0)$, by Lemma 11, any path π contributing to $D_{0,2k,0;k+1}$ has to begin and end with upper vertices. Removing these two vertices resulting a path of length $2k - 2, k - 1$ upper vertices and k lower vertices, which is the same as Case D. Therefore $\pi = \vee^k$.

With the same arguments the paths for Case B and C are also determined to be \setminus^k and $\wedge^k /$ respectively.

Now we show the second identity in Case D, as Cases A, B and C are simpler variations of D and can be verified similarly. We achieve this by associating partial orderings with sets of Dyck paths. The path $\pi = \wedge^k$ imposes the following ordering of the essential coordinates:

$$\begin{array}{ccccccccccc}
 p_{1,1} & < & p_{3,1} & < & p_{5,1} & < & \cdots & < & p_{2k-1,1} & < & k_p = p_{2k+1,1} \\
 \wedge & & \wedge & & \wedge & & \cdots & & \wedge & & \wedge \\
 j_p = p_{1,2} & < & p_{2,2} & < & p_{4,2} & < & \cdots & < & p_{2k-2,2} & < & p_{2k,2}
 \end{array}$$

We relabel these coordinates by $t_{1,1} := p_{1,1}, t_{2,1} = p_{3,1}$ and so on, to obtain

$$\begin{array}{ccccccccccc} t_{1,1} & < & t_{2,1} & < & t_{3,1} & < & \cdots & < & t_{k,1} & < & t_{k+1,1} \\ \wedge & & \wedge & & \wedge & & \cdots & & \wedge & & \wedge \\ t_{1,2} & < & t_{2,2} & < & t_{3,2} & < & \cdots & < & t_{k,2} & < & t_{k+1,2} \end{array} \tag{16}$$

There is a one-one correspondence between the set of all linear extensions of this partial ordering (namely $\{B : B \vdash \pi\}$) and $T_{0,k+1,k+1,0}$. The Dyck path ρ corresponding to the linear extension $t_{m_1,b_1} < t_{m_2,b_2} < \cdots < t_{m_{2k+2},b_{2k+2}}$ is defined by

$$\rho(i) = \begin{cases} \rho(i-1) + 1 & \text{if } b_i = 1, \\ \rho(i-1) - 1 & \text{if } b_i = 2. \end{cases}$$

Clearly, the rank of a linear extension B being (r, r') is equivalent to the corresponding Dyck path starting with r up-steps followed by a down-step and concluding with one down-step with r' up-steps. These cut off the first $r + 1$ and the last $r' + 1$ steps corresponds to $T_{r-1,k-r,k-r',0}$. Therefore

$$D_{r,2k-r-r',r';k}^{\wedge} = |T_{r-1,k-r,k-r',0}| = |T_{0,k-r,k-r',r-1}| = C_{k-r,k-r',r-1}.$$

If $r = 0, r' = 0$, then by Lemma 11 $D_{r,2k-r-r',r';q}^{\wedge} = 0$, which agrees with $C_{k-r,k-r',r-1}$ as well. On the other hand, since the paths of $T_{0,k,k,0}$ only have k up- and down-steps, the LHS is 0 if $r > k$ or $r' > k$, which agrees with the right hand side.

Finally, the \wedge^k are the only possible paths to contribute to the coefficients E , which record the instances when $k_p < j_p$. The corresponding linear extension $B \vdash \wedge^k$ is $\{t_{1,1} < t_{2,1} < \cdots < t_{k+1,1} < t_{1,2} < t_{2,2} < \cdots < t_{k+1,2}\}$. For any other paths, by Lemma 11, any path starting with \backslash or ending with $/$ has j_p as the smallest or k_p as the greatest essential coordinate; on the other hand, any horizontal edge will result in $j_p < k_p$. □

From the above proof we deduce a stronger version of Lemma 11: $D_{0,n,p;q}^{\pi} \neq 0$ only if $\pi(0) = 1$, and for $m \geq 1, D_{m,n,p;q}^{\pi} \neq 0$ only if $\pi(0) = 0$ and begins with \wedge^{m-1} . We also refer to this stronger version as Lemma 11.

Proof of Theorem 6. Case D covers the $2q = m + n + p$ case; moreover, by the same argument as in the proof of the first identities in each case of Lemma 12, for $\pi \in \Pi_{m+n+p+s}$,

$$\epsilon(\pi) + 1 \geq m + n + p + 1 - \epsilon(\pi)$$

therefore $D_{m,n,p;q} \neq 0$ only if $2q \geq m + n + p$. Thus it suffices to show

$$D_{m,n,p;q} = C_{q-1,n-q+1}, \quad 2q > m + n + p.$$

We group the paths into ones starting with $\wedge^k/-$ (call the set of such paths $\Pi^{\wedge k}$) and ones starting with \vee^k- (call the set of such paths $\Pi^{\vee k}$). Then by Lemma II, $D_{m,n,p;q}$ are contributed from $\Pi^{\wedge k}$ if $k - 1 \geq m > 0$, and from $\Pi^{\vee k}$ if $m = 0$:

$$D_{m,n,p;q} = \begin{cases} \sum_{k \geq m-1} \sum_{\pi \in \Pi^{\wedge k}} D_{m,n,p;q}^\pi & \text{if } m > 0, \\ \sum_{k \geq 0} \sum_{\pi \in \Pi^{\vee k}} D_{m,n,p;q}^\pi & \text{if } m = 0, \end{cases}$$

where $D_{m,n,p;q}^\pi$ is the contribution from path π to the coefficient $D_{m,n,p;q}$. Therefore we divide the proof into two cases, $m = 0$ and $m > 0$. The formula for $D_{m,n,p;q}$ with $m + n + p \leq 2$ can be verified by hand (the reader can check their calculation against (14)), so we assume the formula is true for $s \leq m + n + p$, and we want to use induction to verify the formula of $D_{m,n,p;q}$ in general.

5.1. $m > 0$. When $m > 0$, for any $\pi \in \Pi^{\wedge k}$,

$$D_{m,n,p;q}^\pi = D_{m,2k+1-m,0;k+1} D_{0,m+n-2k-2,p;q-k-1}^{\theta_{2k+2}\pi}$$

where $\theta_r: \Pi_s \rightarrow \Pi_{s-r}$, for all $s \geq r$, is the shifting operator such that $(\theta_r \pi)(j) = \pi(j + r)$.

Summing over k and $\pi \in \Pi^{\wedge k}$ we have

$$\begin{aligned} D_{m,n,p;q} &= \sum_{k \geq m-1} \sum_{\pi \in \Pi^{\wedge k}} D_{m,n,p;q}^\pi \\ &= \sum_{k=m-1}^{\lfloor \frac{m+n-2}{2} \rfloor} \sum_{\pi \in \Pi^{\wedge k}} D_{m,2k+1-m,0;k+1} D_{0,m+n-2k-2,p;q-k-1}^{\theta_{2k+2}\pi} \\ &= \sum_{k=m-1}^{\lfloor \frac{m+n-2}{2} \rfloor} D_{m,2k+1-m,0;k+1} D_{0,m+n-2k-2,p;q-k-1} \\ &= \sum_{k=m-1}^{\lfloor \frac{m+n-2}{2} \rfloor} C_{k,k-m+1} C_{q-k-2,m+n-q-k} \\ &= \sum_{k=0}^{\lfloor \frac{n-m}{2} \rfloor} C_{k+m-1,k} C_{q-m-1-k,n-q+1-k} \end{aligned}$$

where the last two equalities comes from the Case C and the induction assumption.

To apply Lemma 10, we check the three conditions hold: (1) The condition “ $n \geq 0$ ” becomes $m - 1 \geq 0$: this is correct as $m > 0$. (2) “ $m \geq p$ ” is $2q \geq m + n + 2$: we know that $2q > m + n + p$, so either $2q \geq m + n + 2$ or $2q = m + n + 1$, in the latter since $m + n + 1 \leq m + n + p + 1 \leq 2q$, we have $p = 0$ and this is covered by Case C. (3) “ $m + n + p + 1 \geq 0$ ” becomes $n \geq 0$, which is evidently true by the definition of $D_{m,n,p;q}$. The upper bound of the summation domain “ $\lfloor \frac{m+p}{2} \rfloor$ ” becomes $\lfloor \frac{n-m}{2} \rfloor$. Therefore we can apply Lemma 10 to the sum above and obtain

$$D_{m,n,p;q} = C_{q-1,n-q+1}.$$

5.2. $m = 0$. When $m = 0$, similarly, for any $\pi \in \Pi^{\vee k}$,

$$D_{0,n,p;q}^\pi = D_{0,2k,0;k+1} D_{0,n-2k-1,p;q-k-1}^{\theta_{2k+1}\pi}.$$

Again, summing over k and $\pi \in \Pi^{\vee k}$ we have

$$\begin{aligned} D_{0,n,p;q} &= \sum_{k \geq 0} \sum_{\pi \in \Pi^{\vee k}} D_{m,n,p;q}^\pi \\ &= \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} \sum_{\pi \in \Pi^{\vee k}} D_{0,2k,0;k+1} D_{0,n-2k-1,p;q-k-1}^{\theta_{2k+1}\pi} \\ &= \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} D_{0,2k,0;k+1} D_{0,n-2k-1,p;q-k-1} \\ &= \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} C_{k,k} C_{q-k-2,n-q-k+1} \end{aligned}$$

where the last equality comes from Case A, the induction assumption and the fact that $C_k = C_{k,k}$. Once again, we want to check the conditions in order to apply Lemma 10. The condition “ $n \geq 0$ ” is obvious. “ $m \geq p$ ” is equivalent to $2q \geq n + 3$. Since $2q > n + p$, there are two possibilities apart from “ $m \geq p$ ”:

- (1) $2q = n + p + 1$. This is covered by Case B;
- (2) $2q = n + p + 2$ and $p = 0$. This is covered by Case A.

“ $m + n + p + 1 \geq 0$ ” is again equivalent to $n \geq 0$, which is evidently true. The upper bound of the summation domain “ $\lfloor \frac{m+p}{2} \rfloor$ ” is $\lfloor \frac{n-1}{2} \rfloor$. Therefore we can apply Lemma 10 to the sum above and obtain

$$D_{0,n,p;q} = C_{q-1,n-q+1}. \quad \square$$

In the above proof of Theorem 6, an induction argument is used to show that the coefficients are indeed the generalised Catalan numbers. Therefore it would be interesting to seek a more direct proof with an explicit correspondence between the linear extensions and Catalan objects (e.g. the Dyck paths).

6. Proof of Theorem 7

In this section we often abuse notations and do not differentiate between operators and their kernels. Without loss of generality assume $|v| = 1$ (otherwise one can scale (a, x, y, b)). We only consider the generating function of coefficient D , as the case for E can be dealt with similarly. We write

$$\phi^N(m, n, p) = \sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y) \prod_{\alpha=0}^{m-1} (x_{j-1} - a - \alpha \Delta_N) \prod_{\beta=0}^{n-1} (x_{k-1} - x_j - \beta \Delta_N) \prod_{\gamma=0}^{p-1} (b - x_k - \gamma \Delta_N),$$

$$\phi(m, n, p) = (x - a)^m (y - x)^n (b - y)^p \mathbb{I}_{a \leq x < y < b},$$

where we recall $\Delta_N = \frac{b-a}{N}$. Then

$$0 \leq \phi^N(m, n, p) \leq \phi(m, n, p) \leq (b - a)^{m+n+p}.$$

We also write

$$\mathcal{D}_{m,n,p} = \sum_{q=0}^s \frac{D_{m,n,p;q}}{m!n!p!} (-1)^q v^{m+n+p+1-2q}.$$

Then

$$W_N = \sum_{s=1}^{N-1} w_{s,N} + \sum_{s=1}^{2N-3} v_{s,N}$$

where $v_{s,N}$ are the degenerate terms, and

$$w_{s,N} = \sum_{m+n+p=s-1} \mathcal{D}_{m,n,p} \phi^N(m, n, p).$$

The reason for the range of the sum for s to be $1 \leq s \leq N - 1$ is because $w_{s,N} = 0$ for $s \geq N$, as

$$\phi^N(m, n, p) = 0, \quad m + n + p \geq N - 1 \tag{17}$$

Proof of Theorem 7. We divide the proof into three parts.

- (1) $\sum_{m+n+p \geq N-1} \mathcal{D}_{m,n,p} \phi^N(m, n, p) \xrightarrow{N \rightarrow \infty} 0$ uniformly on $[a, b]^2$. This shows the limit exists.
- (2) $\sum_{s=1}^{N-1} w_{s,N}$ converges weakly to W .
- (3) The degenerate terms vanish uniformly: $\sum_{s=1}^{2N-3} v_{s,N}$ are arbitrarily small as N grows bigger.

6.1. Part 1. This follows from (17).

6.2. Part 2. We want to show that for any $\epsilon > 0$ and sufficiently large N , we have $\langle f, \sum_{s \leq N-1} (w_{s,N} - w_s)g \rangle < \epsilon \|f\|_2 \|g\|_2$ for testing functions $f, g \in L^2([a, b])$. Equivalently, we must show that

$$\left| \int_a^b \int_a^b \tilde{f}(x, y) \sum_{m+n+p \leq N-2} \mathcal{D}_{m,n,p} (\phi^N(m, n, p) - \phi(m, n, p)) g(x, y) dx dy \right| \leq \epsilon \|f\|_2 \|g\|_2.$$

We divide it into two further parts.

(1) We have

$$\begin{aligned} & (x - a)^m (y - x)^n (b - y)^p \sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j \times A_k}(x, y) \\ & \approx (x - a)^m (y - x)^n (b - y)^p \mathbb{I}_{a \leq x < y < b} \end{aligned}$$

(2) and

$$\begin{aligned} & (x - a)^m (y - x)^n (b - y)^p \sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j \times A_k}(x, y) \\ & \approx \sum_{1 \leq j < k \leq N} \tau^N(j, k; m, n, p) \cdot \mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y), \end{aligned}$$

where

$$\begin{aligned} \tau^N(j, k; m, n, p) := & \prod_{\alpha=0}^{m-1} (x_{j-1} - a - \alpha \Delta_N) \prod_{\beta=0}^{n-1} (x_{k-1} - x_j - \beta \Delta_N) \\ & \prod_{\gamma=0}^{p-1} (b - x_k - \gamma \Delta_N). \end{aligned}$$

6.2.1. Part 2.1. We want to show that

$$\left| \int_a^b \int_a^b \bar{f}(x, y) \sum_{m+n+p \leq N-2} \mathcal{D}_{m,n,p} \sum_{1 \leq j < k \leq N} (x-a)^m (y-x)^n (b-y)^p \right. \\ \left. (\mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y) - \mathbb{I}_{a \leq x < y < b}) g(x, y) dx dy \right| \\ \leq \epsilon.$$

Denote the left hand side by B_N , then by (1) and that

$$(x-a)^m (y-x)^n (b-y)^p \leq (b-a)^{m+n+p}$$

for $a \leq x \leq y \leq b$, we have that

$$B_N \leq \sum_{m+n+p \leq N-2} (m+n+p+1) \frac{(6(b-a))^{m+n+p}}{(m+n+p)!} \\ \left| \sum_{j=1}^N \int \int_{x_{j-1} \leq x < y < x_j} \bar{f}(x) g(y) dx dy \right| \\ \leq \sum_{r \leq N-1} (r+1) \binom{r+2}{2} \frac{(6(b-a))^r}{r!} \\ \left| \sum_{j=1}^N \int \int_{x_{j-1} \leq x < y < x_j} \bar{f}(x) g(y) dx dy \right|.$$

The term in the modulus can be bounded by repeated use of Cauchy–Schwartz inequality:

$$\left| \sum_{j=1}^N \int \int_{x_{j-1} \leq x < y < x_j} \bar{f}(x) g(y) dx dy \right| \leq \sum_{j=1}^N \int_{x_{j-1}}^{x_j} |f(x)| dx \int_x^{x_j} |g(y)| dy \\ \leq \sum_{j=1}^N \int_{x_{j-1}}^{x_j} |f(x)| \sqrt{x_j - x} \|g\|_2 dx \\ \leq \sum_{j=1}^N \frac{(b-a)^2}{2N^2} \|f\|_2 \|g\|_2 \\ = \frac{(b-a)^2}{2N} \|f\|_2 \|g\|_2.$$

Therefore

$$\begin{aligned} B_N &\leq \sum_{r=0}^{N-1} (r+1) \binom{r+2}{2} \frac{(6(b-a))^r (b-a)^2}{r!} \frac{1}{2N} \|f\|_2 \|g\|_2 \\ &\leq CN^{-1} \|f\|_2 \|g\|_2 \end{aligned}$$

for some constant C , where the second bound comes from the fact that

$$\sum_{r \geq 0} (r+1) \binom{r+2}{2} \frac{(6(b-a))^r}{r!} < \infty.$$

6.2.2. Part 2.2. We establish the following uniform convergence, from which weak convergence will follow:

$$\begin{aligned} &\sum_{m+n+p \leq N-2} \mathcal{D}_{m,n,p} \\ &\sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y) \\ &\quad (\tau^N(j, k; m, n, p) - (x-a)^m (y-x)^n (b-y)^p) \longrightarrow 0 \end{aligned}$$

When $a \leq x \leq y \leq b$, τ^N is non-negative, and for $m+n+p \leq N$ we use a telescoping series:

$$\begin{aligned} &(x-a)^m (y-x)^n (b-y)^p - \tau^N(j, k; m, n, p) \\ &= (x-x_{j-1}) \cdots \\ &\quad + (x_{j-1}-a)(x-x_{j-2}) \cdots \\ &\quad + (x_{j-1}-a)(x_{j-2}-a)(x-x_{j-3}) \cdots \\ &\quad + \cdots \\ &\quad + (x_{j-1}-a)(x_{j-2}-a)(x_{j-3}-a) \cdots (x_{k+p-2}-y)(b-y) \\ &\quad + (x_{j-1}-a)(x_{j-2}-a)(x_{j-3}-a) \cdots (b-x_{k+p-2})(x_{k+p-1}-y) \\ &\quad + (x_{j-1}-a)(x_{j-2}-a)(x_{j-3}-a) \cdots (b-x_{k+p-2})(b-x_{k+p-1}) \\ &\quad - (x_{j-1}-a)(x_{j-2}-a)(x_{j-3}-a) \cdots (b-x_{k+p-2})(b-x_{k+p-1}) \\ &\leq N^{-1} (b-a)^{m+n+p} \left(\sum_{\alpha=1}^m \alpha + \sum_{\beta=1}^n \beta + \sum_{\gamma=1}^p \gamma \right) \\ &\leq N^{-1} (b-a)^{m+n+p} \frac{3}{2} (m+n+p)(m+n+p+1). \end{aligned}$$

Therefore

$$\begin{aligned} & \left| \sum_{m+n+p \leq N} \mathcal{D}_{m,n,p} \right. \\ & \quad \left. \sum_{1 \leq j < k \leq N} \mathbb{I}_{A_j}(x) \mathbb{I}_{A_k}(y) (\tau^N(j, k; m, n, p) - (x-a)^m (y-x)^n (b-y)^p) \right| \\ & \leq N^{-1} \sum_{r \leq N} \binom{k+2}{2} \frac{3}{2} r(r+1)^2 \frac{(6(b-a))^r}{r!} \\ & \leq CN^{-1}. \end{aligned}$$

6.3. Part 3. The degenerate terms are the total orderings of path of length s with some repeated essential coordinates. If such a total ordering J has $s + 1 - d$ non-repeated coordinates, then we call d the *degree* of degeneration, or we say that there are d degenerations in J . Each degeneration happens on a wedge part of a path, that is, any two essential coordinates $p_{i_1, j_1} = p_{i_2, j_2}$ if and only if they correspond to parts of the same \wedge^k part of a path for some k . On the other hand, degenerations happen in pairs. That is, for an array p , there do not exist three essential coordinates equal to each other, which would violate the partial ordering. Therefore, given a path of length $s - 1$, the number of degenerate total orderings with d degenerations is bounded by $\left(\frac{s-1}{d}\right)^2$. Moreover, the number of paths of length $s - 1$ is the Fibonacci number $\frac{\Phi^{s+2} - (-\Phi)^{-s-2}}{\sqrt{5}}$ where $\Phi = \frac{\sqrt{5}+1}{2}$. Since a path of length $s - 1$ can have at most $\frac{s-1}{2}$ wedges, there are at most $\frac{s-1}{2}$ degenerations.

Therefore for each N , we have

$$V_N = \sum_{s=1}^{2N-3} v_{s,N}$$

where $V_{s,N}$ is the counterpart of $W_{s,N}$ that collects all degenerate cases of paths of lengths $s - 1$. By applying the calculation in the proof of Lemma 3, we can see the degenerate (of degree d) version of h_s^N is

$$q_s^N(x, y) = \frac{(b-a)^d}{m!n!p!N^d} \phi^N(m, n, p)$$

where $m + n + p + d = s - 1$. Thus the sum is bounded uniformly by

$$\frac{(b-a)^{s-1}}{m!n!p!N^d}.$$

Therefore the total sum of degenerate terms is bounded as follows:

$$\begin{aligned}
 |V_N| &\leq \left| \sum_{s=1}^{N-1} \sum_{d=1}^{\frac{s-1}{2}} \sum_{m+n+p=s-1-d} C \left(\frac{s-1}{d} \right)^2 \Phi^{s+2} \frac{(b-a)^{s-1}}{m!n!p!N^d} \sum_{q=0}^s (-1)^q v^{s-2q} \right| \\
 &\leq \sum_{s=1}^{N-1} \sum_{d=1}^{\frac{s-1}{2}} \sum_{m+n+p=s-1-d} C \left(\frac{s-1}{d} \right)^2 \Phi^{s+2} \frac{(3(b-a))^{s-1}}{(s-1)!} s N^{-d} \\
 &\leq \sum_{d=1}^{\frac{N-2}{2}} \sum_{s=2d+2}^{N-1} C \frac{(6\Phi(b-a))^{s-1} s}{(s-1)!} \binom{s-d+2}{2} \\
 &\leq \sum_{d=1}^{\frac{N-2}{2}} \sum_{s=1}^{N-1} C \frac{(12\Phi(b-a))^{s-1} s}{(s-1)!} \\
 &\leq C(N^{-1} + \sum_{d=2}^{\frac{N-2}{2}} N^{-2}) \\
 &\leq CN^{-1}.
 \end{aligned}$$

□

7. The unitarity of W

Let f and g be the generating functions of D and E :

$$\begin{aligned}
 f &:= \sum_{m,n,p \geq 0} \sum_{q=0}^{m+n+p+1} D_{m,n,p;q} [m, n, p] (-\bar{v})^q v^{m+n+p+1-q} = \sum_{s \geq 1} f_s, \\
 g &:= \sum_{m,n,p \geq 0} \sum_{q=0}^{m+n+p+1} E_{m,n,p;q} [m, n, p]^\dagger (-\bar{v})^q v^{m+n+p+1-q} = \sum_{s \geq 1} g_s.
 \end{aligned}$$

And the kernel of $W - I$ is

$$f(x, y) <_a^b (x, y) + g(x, y) >_a^b (x, y). \tag{18}$$

One can write down the equation that f and g have to satisfy for W to be unitary.

Proposition 13. For W to be unitary, it suffices to show that for any $a < x < y < b$,

$$\begin{aligned} f(x, y) + \overline{g(y, x)} + \int_a^x g(x, z)\overline{g(y, z)}dz \\ + \int_x^y f(x, z)\overline{g(y, z)}dz + \int_y^b f(x, z)\overline{f(y, z)} \\ = 0. \end{aligned} \quad (19)$$

Proof. For W to be unitary it is necessary and sufficient to show it is both a coisometry and an isometry

$$W^*W = WW^* = I$$

Plugging in (18) and using the formulas for kernels of products and adjoints of integral operators we obtain equation (19) and three “other” equations:

$$\begin{aligned} \overline{f(y, x)} + g(x, y) + \int_a^y g(x, z)\overline{g(y, z)}dz + \int_y^x g(x, z)\overline{f(y, z)}dz \\ + \int_x^b f(x, z)\overline{f(y, z)}dz \\ = 0, \quad a < y < x < b, \end{aligned} \quad (20)$$

$$\begin{aligned} \overline{g(y, x)} + f(x, y) + \int_a^x f(z, y)\overline{f(z, x)}dz + \int_x^y f(z, y)\overline{g(z, x)}dz \\ + \int_y^b g(z, y)\overline{g(z, x)}dz \\ = 0, \quad a < x < y < b, \end{aligned} \quad (21)$$

$$\begin{aligned} \overline{f(y, x)} + g(x, y) + \int_a^y f(z, y)\overline{f(z, x)}dz + \int_y^x g(z, y)\overline{f(z, x)}dz \\ + \int_x^b g(z, y)\overline{g(z, x)}dz \\ = 0, \quad a < y < x < b. \end{aligned} \quad (22)$$

The equation (19) and (20) are equivalent: one can interchange x with y and take a conjugate in the former to obtain the latter. So are (21) and (22). By the symmetry of f and g from Corollary 5, (19) and (21) are equivalent, hence it suffices to verify (19) to show the unitarity of W . \square

Proof of Theorem 8. We divide the proof into two parts: first we write down f and g in a more amenable form, then we proceed to proving the integral identity.

7.1. The formulas for f and g . As previous, without loss of generality suppose $|\nu| = 1$. Recall that

$$B_j(x, y) = \sum_{n \geq 0} \frac{(-1)^{n+j} x^{n+j} y^n}{(n+j)!n!} = (-1)^j (x/y)^{j/2} J_j(2\sqrt{xy}).$$

Since $E_{m,n,p;q} = \mathbb{I}_{m=p=q,n=0}$,

$$g(x, y) = \nu \sum_{m \geq 0} (-1)^m \frac{(y-a)^m (b-x)^m}{m!m!} = \nu B_0(y-a, b-x). \quad (23)$$

Let f_e and f_o be the parts of the sum of f where $m+n+p$ are even and odd respectively, i.e.,

$$f_e = \sum_{2|m+n+p} \sum_{q=0}^{m+n+p+1} (x-a)^m (y-x)^n (b-y)^p (-\bar{\nu})^q \nu^{m+n+p+1-q},$$

$$f_o = \sum_{2 \nmid m+n+p} \sum_{q=0}^{m+n+p+1} (x-a)^m (y-x)^n (b-y)^p (-\bar{\nu})^q \nu^{m+n+p+1-q}.$$

The function f_e can be further divided into the $\binom{n}{q-m}$ part and the rest. Let $u := (x-a)\nu$, $v := (y-x)\nu$, $w := (b-y)\nu$ and $z = -\frac{\bar{\nu}}{\nu} = -\bar{\nu}^2$, then

$$\begin{aligned} & \nu \sum_{q \geq 0} \sum_{m+n+p=2q} \frac{u^m v^n w^p z^q}{m!n!p!} \binom{n}{q-m} \\ &= \nu \sum_{q \geq 0} \sum_{m+n+p=2q} \frac{u^m v^{2q-m-p} w^p z^q}{m!(2q-m-p)!p!} \binom{2q-m-p}{q-m} \\ &= \nu \sum_{q \geq 0} \sum_{0 \leq m, p \leq q} \frac{u^m v^{2q-m-p} w^p z^q}{m!(2q-m-p)!p!} \binom{2q-m-p}{q-m} \quad (24) \\ &= \nu \sum_{q \geq 0} v^{2q} z^q \sum_{m=0}^q \frac{(u/v)^m}{m!(q-m)!} \sum_{p=0}^q \frac{(w/v)^p}{p!(q-p)!} \\ &= \nu \sum_{q \geq 0} \frac{(u+v)^q (v+w)^q z^q}{q!q!} = \nu B_0(y-a, b-x). \end{aligned}$$

The rest of f_e is slightly more complicated. Let $k = \frac{m+n+p}{2}$. We observe that

$$\begin{aligned} & \sum_q D_{m,n,p;q} z^q - \binom{n}{k-m} z^k \\ &= -\binom{n}{k} z^k + \binom{n}{k} z^{k+1} - \binom{n}{k+1} z^{k+1} + \dots \\ & \quad + \binom{n}{n-1} z^n - \binom{n}{n} z^n + \binom{n}{n} z^{n+1} \\ &= (z-1) \sum_{q=k}^n \binom{n}{q} z^q. \end{aligned}$$

Therefore the rest of f_e , i.e. the sum excluding the terms corresponding to $\binom{n}{k-m}$ is

$$\begin{aligned} & v(z-1) \sum_{k \geq 0} \sum_{m+n+p=2k} \frac{u^m v^n w^p}{m!n!p!} \sum_{q=k}^n \binom{n}{q} z^q \\ &= v(z-1) \sum_{k \geq 0} \sum_{n=k}^{2k} \sum_{q=k}^n \binom{n}{q} z^q \frac{(u+w)^{2k-n}}{(2k-n)!} \frac{v^n}{n!} \\ &= v(z-1) \sum_{k \geq 0} \sum_{n=k}^{2k} \sum_{q=k}^n \frac{z^q}{q!} \frac{(u+w)^{2k-n}}{(2k-n)!} \frac{v^n}{(n-q)!} \\ &= v(z-1) \sum_{k \geq 0} \sum_{n=0}^k \sum_{q=k}^{k+n} \frac{z^q}{q!} \frac{(u+w)^{k-n}}{(k-n)!} \frac{v^{k+n}}{(k+n-q)!} \\ &= v(z-1) \sum_{k \geq 0} \sum_{n=0}^k \sum_{q=0}^n \frac{z^{k+q}}{(k+q)!} \frac{(u+w)^{k-n}}{(k-n)!} \frac{v^{k+n}}{(n-q)!} \\ &= v(z-1) \sum_{k \geq 0} \sum_{q=0}^k \sum_{n=0}^{k-q} \frac{z^{k+q}}{(k+q)!} \frac{(u+w)^{k-q-n}}{(k-q-n)!} \frac{v^{k+q+n}}{n!} \\ &= v(z-1) \sum_{k \geq 0} \sum_{q=0}^k \frac{(zv)^{k+q}}{(k+q)!} \frac{(u+v+w)^{k-q}}{(k-q)!} \\ &= v(z-1) \sum_{q \geq 0} \sum_{k \geq 0} \frac{(zv)^{k+2q}}{(k+2q)!} \frac{(u+v+w)^k}{k!}. \end{aligned} \tag{25}$$

We keep (25) to later merge it with a similar term in f_o .

For f_o , let $k = \frac{m+n+p-1}{2}$. We observe that

$$\begin{aligned} & \sum_q D_{m,n,p;q} z^q - \binom{n}{k-m} z^k \\ &= \binom{n}{k} z^{k+1} - \binom{n}{k+1} z^{k+1} + \dots + \binom{n}{n-1} z^n - \binom{n}{n} z^n + \binom{n}{n} z^{n+1} \\ &= (z-1) \sum_{q=k+1}^n \binom{n}{q} z^q + \binom{n}{k} z^{k+1}. \end{aligned}$$

Following the same procedure which leads (25), the sum corresponding to the first term in the RHS is

$$v(z-1) \sum_{q \geq 0} \sum_{k \geq 0} \frac{(zv)^{k+2q+1}}{(k+2q+1)!} \frac{(u+v+w)^k}{k!} \tag{26}$$

whereas the contribution from the second term is computed as follows:

$$\begin{aligned} & v \sum_{k \geq 0} \sum_{m+n+p=2k+1} \frac{u^m v^n w^p}{m!n!p!} \binom{n}{k} z^{k+1} \\ &= v \sum_{k \geq 0} \sum_{n=k}^{2k+1} \frac{v^n (u+w)^{2k+1-n}}{n! (2k+1-n)!} \binom{n}{k} z^{k+1} \\ &= v \sum_{k \geq 0} \sum_{n=0}^{k+1} \frac{v^{n+k}}{n!k!} \frac{(u+w)^{k+1-n}}{(k+1-n)!} z^{k+1} \\ &= v \sum_{k \geq 0} \frac{v^k z^{k+1} (v+u+w)^{k+1}}{k! (k+1)!} = B_1(b-a, y-x). \end{aligned} \tag{27}$$

By summing up (24), (25), (26), and (27) and plugging in u, v, w, z we obtain

$$\begin{aligned} f(x, y) &= vB_0(y-a, b-x) + B_1(b-a, y-x) \\ &\quad - (v + \bar{v}) \sum_{q \geq 0} B_q(y-x, b-a) \bar{v}^q. \end{aligned} \tag{28}$$

Note that

$$\sum_{q \geq 0} B_q(y-x, b-a) \bar{v}^q = \sum_{q \geq 0} J_q(2\sqrt{(b-a)(y-x)}) \left(-\sqrt{\frac{y-x}{b-a}} \bar{v} \right)^q$$

is a generating function of the Bessel functions.

7.2. Verifying the identity (19). We list a few useful properties of B_j (where we let $B_{-1}(x, y) := -B_1(y, x)$):

- (1) $\partial_x B_j(x, y) = -B_{j-1}(x, y), j \geq 0,$
- (2) $B_0(x, y) = B_0(y, x),$
- (3) $B_j(0, y) = \delta_{j0}, j \geq 0,$
- (4) $\partial_y B_j(x, y) = B_{j+1}(x, y).$

And an integral:

$$\begin{aligned}
 & \int_y^b B_0(b-x, z-a) B_j(z-y, b-a) dz \\
 &= - \sum_{k \geq 0} B_k(b-x, z-a) B_{k+j+1}(z-y, b-a) \Big|_y^b \\
 &= \begin{cases} - \sum_{k \geq 0} B_k(b-x, b-a) B_{k+j+1}(b-y, b-a) & \text{if } j \geq 0, \\ \sum_{k \geq 0} B_k(b-x, b-a) B_k(b-y, b-a) - B_0(b-x, y-a) & \text{if } j = -1. \end{cases}
 \end{aligned} \tag{29}$$

Substituting for f from (28) and g from (23) into (19) gives

$$\begin{aligned}
 & B_1(b-a, y-x) \\
 &+ \int_a^b B_0(z-a, b-x) B_0(z-a, b-y) dz \\
 &+ \int_y^b B_1(b-a, z-x) B_1(b-a, z-y) dz \\
 &+ v B_0(y-a, b-x) + \bar{v} B_0(x-a, b-y) \\
 &+ \bar{v} \int_x^b B_1(b-a, z-x) B_0(z-a, b-y) dz \\
 &+ v \int_y^b B_1(b-a, z-y) B_0(z-a, b-x) dz \\
 &- (v + \bar{v}) \int_y^b \sum_{q \geq 0} B_1(b-a, z-x) B_q(z-y, b-a) v^q dz \\
 &- (v + \bar{v}) \int_y^b \sum_{q \geq 0} B_1(b-a, z-y) B_q(z-x, b-a) \bar{v}^q dz \\
 &- \bar{v}(v + \bar{v}) \int_x^b \sum_{q \geq 0} B_0(z-a, b-y) B_q(z-x, b-a) \bar{v}^q dz
 \end{aligned}$$

$$\begin{aligned}
 & -v(v + \bar{v}) \int_y^b \sum_{q \geq 0} B_0(z - a, b - x) B_q(z - y, b - a) v^q dz \\
 & + (v + \bar{v})^2 \int_y^b \sum_{q \geq 0} B_q(z - x, b - a) \bar{v}^q \sum_{q \geq 0} B_q(z - y, b - a) v^q dz \\
 & - (v + \bar{v}) \sum_{q \geq 0} B_q(y - x, b - a) \bar{v}^q \\
 & = 0.
 \end{aligned} \tag{30}$$

Let

$$G_j(x) = \sum_{k \geq 0} \frac{(-1)^k x^k}{k!(k + 1)!},$$

then for $\alpha > 0$ the following two integrals hold:

$$\begin{aligned}
 & \int_0^x G_0(\alpha z) G_0(\beta z) dz \\
 & = (\alpha - \beta)^{-1} (\alpha x G_1(\alpha x) G_0(\beta x) - \beta x G_1(\beta x) G_0(\alpha x))
 \end{aligned}$$

and

$$\begin{aligned}
 & \int_0^z G_1(w) G_1(w + \alpha) dw \\
 & = \alpha^{-1} (z G_1(z) G_0(z + \alpha) - (z + \alpha) G_1(z + \alpha) G_0(z)) + G_1(\alpha).
 \end{aligned}$$

The first of these two integrals is the well-known Lommel’s integral, see e.g. Section 11 and 94 of [2]. The second integral written in terms of an indefinite integral of the Bessel functions is

$$\begin{aligned}
 & \int \frac{1}{\sqrt{w^2 + \beta^2}} J_1(w) J_1(\sqrt{w^2 + \beta^2}) dw \\
 & = \beta^{-2} (w J_1(w) J_0(\sqrt{w^2 + \beta^2}) - \sqrt{w^2 + \beta^2} J_1(\sqrt{w^2 + \beta^2}) J_0(w)), \quad \beta > 0.
 \end{aligned} \tag{31}$$

This is a special case of the so-called Sonine-Gegenbauer type integral (see e.g. page 415 of [30]). However, the authors have not found an explicit formula like the one on the right hand side of (31) in the literature.

By these two integrals we have

$$\begin{aligned}
 & \int_a^b B_0(z - a, b - x) B_0(z - a, b - y) dz \\
 & = (y - x)^{-1} ((b - y) B_1(b - a, b - y) B_0(b - a, b - x) \\
 & \quad - (b - x) B_1(b - a, b - x) B_0(b - a, b - y))
 \end{aligned}$$

and

$$\begin{aligned} & \int_y^b B_1(b-a, z-x)B_1(b-a, z-y)dz \\ &= B_1(b-a, y-x) \\ & \quad - (y-x)^{-1}((b-y)B_1(b-a, b-y)B_0(b-a, b-x) \\ & \quad - (b-x)B_1(b-a, b-x)B_0(b-a, b-y)). \end{aligned}$$

Therefore the first and the second lines of (30) vanishes.

By the integral (29) (and interchanging x and y when necessary), the third and the fourth lines are reduced to their real part.

Now if the real part of ν is 0 then we are done. Otherwise by subtracting the first and the second lines and the imaginary part of the third and the fourth lines from (30), and dividing the remainder by the real part $\frac{\nu+\bar{\nu}}{2}$, we simplify the integral identity into

$$\begin{aligned} & B_0(y-a, b-x) \\ & + B_0(x-a, b-y) \\ & + \int_x^b B_1(b-a, z-x)B_0(z-a, b-y)dz \\ & + \int_y^b B_1(b-a, z-y)B_0(z-a, b-x)dz \\ & - 2 \sum_{q \geq 0} B_q(y-x, b-a)\bar{\nu}^q \\ & - 2 \int_y^b \sum_{q \geq 0} B_1(b-a, z-x)B_q(z-y, b-a)\nu^q dz \\ & - 2 \int_y^b \sum_{q \geq 0} B_1(b-a, z-y)B_q(z-x, b-a)\bar{\nu}^q dz \\ & - 2\bar{\nu} \int_x^b \sum_{q \geq 0} B_0(z-a, b-y)B_q(z-x, b-a)\bar{\nu}^q dz \\ & - 2\nu \int_y^b \sum_{q \geq 0} B_0(z-a, b-x)B_q(z-y, b-a)\nu^q dz \\ & + 2(\nu + \bar{\nu}) \int_y^b \sum_{q \geq 0} B_q(z-x, b-a)\bar{\nu}^q \sum_{q \geq 0} B_q(z-y, b-a)\nu^q dz \\ & = 0 \end{aligned}$$

By the integral formulas (29), the above identity can be further simplified to

$$\begin{aligned}
 & \sum_{k \geq 0} B_k(b-x, b-a) B_k(b-y, b-a) \\
 & + \sum_{q > 0} \sum_{k \geq 0} B_k(b-x, b-a) B_{k+q}(b-y, b-a) v^q \\
 & + \sum_{q > 0} \sum_{k \geq 0} B_k(b-y, b-a) B_{k+q}(b-x, b-a) \bar{v}^q \\
 & - \sum_{q \geq 0} B_q(y-x, b-a) \bar{v}^q \\
 & - \int_y^b \sum_{q \geq 0} B_1(b-a, z-x) B_q(z-y, b-a) v^q dz \\
 & - \int_y^b \sum_{q \geq 0} B_1(b-a, z-y) B_q(z-x, b-a) \bar{v}^q dz \\
 & + (v + \bar{v}) \int_y^b \sum_{q \geq 0} B_q(z-x, b-a) \bar{v}^q \sum_{q \geq 0} B_q(z-y, b-a) v^q dz \\
 & = 0
 \end{aligned}$$

It remains to verify the coefficient of v^q for each $q \in \mathbb{Z}$ in the LHS is 0, which can be done by repeated use of integration by parts and the properties of the functions B_j .

When $q = 0$, the coefficient is

$$\begin{aligned}
 & \sum_{j \geq 0} B_j(b-x, b-a) B_j(b-y, b-a) \\
 & - B_0(y-x, b-a) \\
 & + \int_y^b -B_1(b-a, z-y) B_0(z-x, b-a) \\
 & \quad - B_1(b-a, z-x) B_0(z-y, b-a) \\
 & + \sum_{j \geq 0} B_{j+1}(z-x, b-a) B_j(z-y, b-a) \\
 & + \sum_{j \geq 0} B_{j+1}(z-y, b-a) B_j(z-x, b-a) dz
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{j \geq 0} B_j(b-x, b-a) B_j(b-y, b-a) - B_0(y-x, b-a) \\
&\quad + \sum_{j \geq -1} \int_y^b B_j(z-x, b-a) B_{j+1}(z-y, b-a) \\
&\quad\quad + B_j(z-y, b-a) B_{j+1}(z-z, b-a) dz \\
&= \sum_{j \geq 0} B_j(b-x, b-a) B_j(b-y, b-a) - B_0(y-x, b-a) \\
&\quad - \sum_{j \geq 0} B_j(z-x, b-a) B_j(z-y, b-a) \Big|_y^b \\
&= 0,
\end{aligned}$$

where in the fourth line we use integration by parts and the properties of B_j .

When $q > 0$, the coefficient is

$$\begin{aligned}
&\sum_{k \geq 0} B_k(b-x, b-a) B_{k+q}(b-y, b-a) \\
&\quad + \int_y^b -B_1(b-a, z-x) B_q(z-y, b-a) \\
&\quad\quad + \sum_{j \geq 0} B_j(z-x, b-a) B_{j+q-1}(z-y, b-a) \\
&\quad\quad + \sum_{j \geq 0} B_j(z-x, b-a) B_{j+q+1}(z-y, b-a) dz \\
&= \sum_{k \geq 0} B_k(b-x, b-a) B_{k+q}(b-y, b-a) \\
&\quad + \int_y^b \sum_{j \geq 0} B_j(z-x, b-a) B_{j+q-1}(z-y, b-a) \\
&\quad\quad + \sum_{j \geq -1} B_j(z-x, b-a) B_{j+q+1}(z-y, b-a) dz \\
&= \sum_{k \geq 0} B_k(b-x, b-a) B_{k+q}(b-y, b-a) \\
&\quad + \int_y^b \sum_{j \geq 0} B_j(z-x, b-a) B_{j+q-1}(z-y, b-a) \\
&\quad\quad + \sum_{j \geq 0} B_{j-1}(z-x, b-a) B_{j+q}(z-y, b-a) dz
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{k \geq 0} B_k(b-x, b-a) B_{k+q}(b-y, b-a) \\
 &\quad - \sum_{k \geq 0} B_k(z-x, b-a) B_{k+q}(z-y, b-a) \Big|_y^b \\
 &= 0.
 \end{aligned}$$

When $q < 0$, the coefficient is

$$\begin{aligned}
 &\sum_{k \geq 0} B_k(b-y, b-a) B_{k+q}(B-x, b-a) \\
 &\quad - B_q(y-x, b-a) \\
 &\quad + \int_y^b -B_1(b-a, z-y) B_q(z-x, b-a) \\
 &\quad \quad + \sum_{k \geq 0} B_{k+q-1}(z-x, b-a) B_k(z-y, b-a) \\
 &\quad \quad + \sum_{k \geq 0} B_{k+q+1}(z-x, b-a) B_k(z-y, b-a) dz \\
 &= \sum_{k \geq 0} B_k(b-y, b-a) B_{k+q}(B-x, b-a) \\
 &\quad - B_q(y-x, b-a) \\
 &\quad + \sum_{k \geq 0} \int_y^b B_{k+q-1}(z-x, b-a) B_k(z-y, b-a) \\
 &\quad \quad + B_{k+q}(z-x, b-a) B_{k-1}(z-y, b-a) dz \\
 &= \sum_{k \geq 0} B_k(b-y, b-a) B_{k+q}(B-x, b-a) - B_q(y-x, b-a) \\
 &\quad - \sum_{k \geq 0} B_{k+q}(z-x, b-a) B_k(z-y, b-a) \Big|_y^b = 0. \quad \square
 \end{aligned}$$

References

- [1] D. B. Applebaum, Robin Hudson's pathless path to quantum stochastic calculus. *Commun. Stoch. Anal.* **4** (2010), no. 4, 481–491. [MR 2743665](#) [Zbl 1331.81165](#)
- [2] F. Bowman, *Introduction to Bessel functions*. Longmans, Green & Co., London etc., 1938. Reprint, Dover Publications, New York, 1958. [MR 0097539](#) [Zbl 0083.05602](#) (reprint) [JFM 64.1087.01](#) [MR 0020.11205](#)

- [3] F. Baudoin and J. Wang, Stochastic areas, winding numbers and Hopf fibrations. *Probab. Theory Related Fields* **169** (2017), no. 3–4, 977–1005. [MR 3719061](#) [Zbl 06814045](#)
- [4] M. Capitaine and C. Donati-Martin, The Lévy area process for the free Brownian motion. *J. Funct. Anal.* **179** (2001), no. 1, 153–169. [MR 1807256](#) [Zbl 0979.60044](#)
- [5] A. M. Cockroft and R. L. Hudson, Quantum mechanical Wiener processes. *J. Multivariate Anal.* **7** (1977), no. 1, 107–124. [MR 0450495](#) [Zbl 0450495](#)
- [6] S. Chen and R. Hudson, Some properties of quantum Lévy area in Fock and non-Fock quantum stochastic calculus. *Probab. Math. Statist.* **33** (2013), no. 2, 425–434. [MR 3158567](#) [Zbl 1283.81099](#)
- [7] A. Deya and R. Schott, On the rough-paths approach to non-commutative stochastic calculus. *J. Funct. Anal.* **265** (2013), no. 4, 594–628. [MR 3062538](#) [Zbl 1291.46056](#)
- [8] P. K. Friz and N. B. Victoir, *Multidimensional stochastic processes as rough paths*. Theory and applications. Cambridge Studies in Advanced Mathematics, 120. Cambridge University Press, Cambridge, 2010. [MR 2604669](#) [Zbl 1193.60053](#)
- [9] M. R. Hush, A. R. R. Carvalho, M. Hedges, and M. R. James, Analysis of the operation of gradient echo memories using a quantum input–output model. *New J. Phys.* **15** (2013), no. 8, 085020.
- [10] R. L. Hudson, P. D. F. Ion, and K. R. Parthasarathy, Time-orthogonal unitary dilations and noncommutative Feynman–Kac formulae. *Comm. Math. Phys.* **83** (1982), no. 2, 261–280. [MR 0649162](#) [Zbl 0485.46038](#)
- [11] R. L. Hudson and P. Jones, Explicit construction of a unitary double product integral. In M. Bożejko, A. Krystek, and Ł. Wojakowski (eds.), *Noncommutative harmonic analysis with applications to probability III*. Papers from the 13th Workshop held in Będlewo, July 11–17, 2010. Banach Center Publications, 96. Polish Academy of Sciences, Institute of Mathematics, Warsaw, 2012, 215–236. [MR 2986829](#) [Zbl 1271.81091](#)
- [12] R. L. Hudson and K. R. Parthasarathy, The Feynman–Kac formula for boson wiener processes. In K. E. Gustafson and W. P. Reinhardt (eds.) *Quantum mechanics in mathematics, chemistry, and physics*. Plenum Press, New York and London, 1981, 479–493.
- [13] R. L. Hudson and K. R. Parthasarathy, Quantum Ito’s formula and stochastic evolutions. *Comm. Math. Phys.* **93** (1984), no. 3, 301–323. [MR 0745686](#) [Zbl 0546.60058](#)
- [14] R. L. Hudson and S. Pulmannová, Double product integrals and Enriquez quantisation of Lie bialgebras. II. The quantum Yang–Baxter equation. *Lett. Math. Phys.* **72** (2005), no. 3, 211–224. [MR 2164611](#) [Zbl 1079.53139](#)
- [15] R. Hudson and Y. Pei, Unitary causal quantum stochastic double products as universal interactions I. *Acta Phys. Polon. B* **46** (2015), no. 9, 1851–1869. [MR 3403844](#) [Zbl 1371.81194](#)
- [16] R. L. Hudson, Boson-fermion correspondence on the circle via quantum stochastic calculus. *Internat. J. Theoret. Phys.* **39** (2000), no. 3, 737–744. [MR 1792192](#) [Zbl 0987.81046](#)

- [17] R. L. Hudson, A double dilation constructed from a double product of rotations. *Markov Process. Related Fields* **13** (2007), no. 1, 169–190. J. T. Lewis memorial volume. [MR 2321756](#) [Zbl 1134.81030](#)
- [18] R. L. Hudson, An analytic double product integral. In L. Accardi, W. Freudenberg, and M. Schürmann (eds.), *Quantum probability and infinite dimensional analysis*. Proceedings of the 26th Conference held in Levico, February 20–26, 2005. Quantum Probability and White Noise Analysis, 20. World Scientific, Hackensack, N.J., 2007, 241–250. [MR 2359397](#) [Zbl 1135.28300](#)
- [19] R. L. Hudson, Quantum Lévy area as a quantum martingale limit. In L. Accardi and F. Fagnola (eds.), *Quantum probability and related topics*. Proceedings of the 32nd Conference Levico Terme, Italy, 29 May–4 June 2011. World Scientific, Hackensack, N.J., 2013, 169–188.
- [20] R. L. Hudson, Forward and backward adapted quantum stochastic calculus and double product integrals. *Russ. J. Math. Phys.* **21** (2014), no. 3, 348–361. [MR 3255131](#) [Zbl 1311.81164](#)
- [21] N. Ikeda and S. Taniguchi, The Itô–Nisio theorem, quadratic Wiener functionals, and 1-solitons. *Stochastic Process. Appl.* **120** (2010), no. 5, 605–621. [MR 2603056](#) [Zbl 1203.60059](#)
- [22] N. Ikeda and S. Taniguchi, Euler polynomials, Bernoulli polynomials, and Lévy’s stochastic area formula. *Bull. Sci. Math.* **135** (2011), no. 6-7, 684–694. [MR 2838097](#) [Zbl 1236.60049](#)
- [23] P. Lévy, Wiener’s random function, and other Laplacian random functions. In J. Neyman (ed.), *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*. 1950. University of California Press, Berkeley and Los Angeles, 1951, 171–187. [MR 0044774](#) [Zbl 0044.13802](#)
- [24] J. Meixner, Orthogonale polynomsysteme mit einer besonderen gestalt der erzeugenden funktion. *J. London Math. Soc.* **9** (1934), no. 1, 6–13. [MR 1574715](#) [Zbl 0008.16205](#)
- [25] J. Ortman, Functionals of the Brownian bridge. In C. Donati-Martin, A. Lejay, and A. Rouault (eds.), *Séminaire de Probabilités XLV*. Including papers from the Conference on Stochastic Filtrations held in Strasbourg, September 2011. Lecture Notes in Mathematics, 2078. Springer, Cham, 2013, 433–458. [MR 3185925](#) [Zbl 1291.46063](#)
- [26] K. R. Parthasarathy, An Introduction to quantum stochastic calculus. Monographs in Mathematics, 85. Birkhäuser Verlag, Basel, 1992. [MR 1164866](#) [Zbl 0751.60046](#)
- [27] S. Reuveni, Catalan’s trapezoids. *Probab. Engrg. Inform. Sci.* **28** (2014), no. 3, 353–361. [MR 3214794](#) [Zbl 1351.05014](#)
- [28] A. Slavík, Product integration, its history and applications. Corrected translation of the Czech original. Nečas Center for Mathematical Modeling, 1. Dějiny Matematiky/History of Mathematics, 29. Matfyzpress, Prague, 2007. [MR 2917851](#) [Zbl 1216.28001](#)
- [29] N. Victoir, Lévy area for the free Brownian motion: existence and non-existence. *J. Funct. Anal.* **208** (2004), no. 1, 107–121. [MR 2034293](#) [Zbl 1062.46055](#)

- [30] G. N. Watson, A treatise on the theory of Bessel functions. Reprint of the second (1944) edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1995. [MR 1349110](#) [Zbl 0849.33001](#)
- [31] M. Yor, Remarques sur une formule de Paul Lévy. In J. Azéma and M. Yor (eds.), *Séminaire de Probabilités. XIV*. Held in Paris, 1978/1979. Lecture Notes in Mathematics, 784. Springer, Berlin, 1980, 343–346. [MR 0580140](#) [Zbl 0429.60045](#)

© European Mathematical Society

Communicated by Adrian Tanasă

Received April 12, 2016; accepted November 11, 2016

Robin L. Hudson, Mathematics Department, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK

e-mail: r.hudson@lboro.ac.uk

Yuchen Pei, KTH Mathematics, Lindstedtsvägen 25, 100 44 Stockholm, Sweden

e-mail: me@ypei.me