

# ANNALES SCIENTIFIQUES DE L'É.N.S.

HEINZ SIEDENTOP

RUDI WEIKARD

**A new phase space localization technique with application to the sum of negative eigenvalues of Schrödinger operators**

*Annales scientifiques de l'É.N.S. 4<sup>e</sup> série*, tome 24, n° 2 (1991), p. 215-225

[http://www.numdam.org/item?id=ASENS\\_1991\\_4\\_24\\_2\\_215\\_0](http://www.numdam.org/item?id=ASENS_1991_4_24_2_215_0)

© Gauthier-Villars (Éditions scientifiques et médicales Elsevier), 1991, tous droits réservés.

L'accès aux archives de la revue « Annales scientifiques de l'É.N.S. » (<http://www.elsevier.com/locate/ansens>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

# A NEW PHASE SPACE LOCALIZATION TECHNIQUE WITH APPLICATION TO THE SUM OF NEGATIVE EIGENVALUES OF SCHRÖDINGER OPERATORS

BY HEINZ SIEDENTOP AND RUDI WEIKARD

## 1. Introduction

Let  $H$  be the hamiltonian of  $N$  electrons in the field of a nucleus of charge  $Z$ , *i. e.*,

$$(1) \quad H = \sum_{i=1}^N \left( -\Delta_i - \frac{Z}{|\mathbf{r}_i|} \right) + \sum_{\substack{i, j=1 \\ i < j}}^N \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|}$$

selfadjointly realized in  $\bigwedge_{i=1}^N (L^2(\mathbb{R}^3) \otimes \mathbb{C}^q)$ ,  $q$  being the number of spin states of a single electron. It has been shown (upper bound: Siedentop and Weikard [10], Bach [1]; lower bound: Siedentop and Weikard [9], Bach [1] based on a work of Hughes [3], [4]) that for fixed or negative degree of ionization  $\lambda = 1 - N/Z$  — an assumption that we make throughout the whole paper without further mentioning — the following holds:

THEOREM 1:

$$E_Q(Z, N) = E_{TF}(1, N/Z) Z^{7/3} + \frac{q}{8} Z^2 + O(Z^{47/24})$$

where  $E_Q(Z, N) = \inf \sigma(H)$  and  $E_{TF}(Z, N)$  is the corresponding Thomas-Fermi energy.

The upper bound is obtained by a variational calculation using eigenfunctions of the operator

$$(2) \quad H_l^0 = -\frac{d^2}{dr^2} + \frac{l(l+1)}{r^2} - \frac{Z}{r}$$

for small  $l$ , *i. e.*,  $l < L = [Z^{1/12}]$ , the integer part of  $Z^{1/12}$ , and Macke orbitals, *i. e.*,

$$(3) \quad \varphi_{l, n}(r) = \left( \frac{\rho_l(r)}{N_l} \right)^{1/2} \exp \left( i\pi k_{l, n} \int_0^r \frac{\rho_l}{N_l} dt \right)$$

for  $l \geq L$  where  $N_l = \int \rho_l$  and the  $k_{l, n}$  differ for fixed  $l$  by even integers from each other.  $\rho_l$  is some suitable nonnegative function which may be interpreted as radial electron density in the angular momentum channel  $l$  (Siedentop and Weikard [8]).

For the lower bound one observes that

$$(4) \quad E_Q(Z, N) \geq \sum_{l=0}^{\infty} q(2l+1) \operatorname{tr}(\tilde{H}_l)_- + \frac{1}{3} E_{\text{TF}}(Z, N) - \text{const } Z^{5/3}$$

(Lieb [5], Siedentop and Weikard [9], Bach [1]). Here

$$(5) \quad \tilde{H}_l = -\frac{d^2}{dr^2} + \frac{l(l+1)}{r^2} - (\phi(r) - \mu)_+ = -\frac{d^2}{dr^2} - \phi_l,$$

$\phi - \mu$  being the Thomas-Fermi potential of the atom, *i. e.*, the solution of

$$\left(\frac{6\pi^2}{q}\right)^{2/3} \rho^{2/3}(r) = (\phi(r) - \mu)_+$$

$$\phi(r) = \frac{Z}{r} - \rho * \frac{1}{|\cdot|}(r)$$

under the condition  $\int \rho = \min\{Z, N\}$ , and  $(\cdot)$  denotes the restriction of the operator under consideration onto its negative spectral subspace. Again one distinguishes large and small angular momenta. For small  $l$  one may drop  $\rho_{\text{TF}} * 1/|\cdot|$  in (5) to obtain a bound which can be evaluated explicitly. However, the channels for large  $l$  were treated by a cumbersome WKB analysis (Hughes [3], [4], Siedentop and Weikard [9]).

The purpose of the present paper is to give a simpler proof of the lower bound using Macke orbitals requiring almost only known properties from the upper bound. We remark that  $\{\varphi_{l, n} | n \in \mathbb{Z}\}$  is an orthonormal basis on  $L^2(\mathbb{R}^+)$ , if  $\rho_l$  is positive almost everywhere. Moreover the Macke orbitals yield a phase space localization through their « densities »  $\rho_l$  and their « momenta »  $\pi k_{l, n}$ .

The strategy of our proof is somewhat reminiscent to those of Berezin [2] and Lieb [5] who got upper and lower bounds using coherent states. Our idea is as follows. We break the one-particle operators  $\tilde{H}_l$  into a sum of operators  $H_{l, n}$  operating on almost orthogonal two-dimensional subspaces each of which has at most one bound state. The sum of the corresponding eigenvalues turns out to be the energy in the angular momentum channel  $l$  up to tolerable errors. This is the content of Section 2. The proof of the Scott type lower bound is basically the collection of the various pieces and an optimization of the various error bounds which is done in Section 3. The appendix is a collection of useful estimates most of which are transcriptions from [10] and [1].

## 2. Phase space localization by Macke orbitals

Now we introduce the Macke orbitals  $\varphi_{l, n}$  of (3) explicitly by defining the radial densities  $\rho_l$ . Let

$$(6) \quad L_Z(r) = r(\phi(r) - \mu)_+^{1/2}$$

and define  $k \in \mathbb{R}^+$  by  $\beta_k = \max \{ L_Z(r)^2 \mid r \in \mathbb{R}^+ \}$  where  $\beta_x = x(x+1)$  in our case. For  $l \leq [k] - 1$  we set

$$(7) \quad \rho_l(r) = q(2l+1) \begin{cases} \alpha^2 r^{2l+2}, & \text{if } 0 \leq r \leq x_1, \\ \frac{1}{\pi r} (L_Z(r)^2 - \beta_l)_+^{1/2}, & \text{if } x_1 \leq r \leq x_2, \\ \beta^2 \exp(-2Z^{2/3}r), & \text{if } x_2 \leq r. \end{cases}$$

The  $l$  and  $Z$  dependent numbers  $\alpha^2$  and  $\beta^2$  ensure continuity of  $\rho_l$  at  $x_1 = r_1 + T(\beta_l^{1/2}/Z)$  and  $x_2 = r_2 - SZ^{-2/3}$  where  $T$  and  $S$  are suitable positive constants and  $r_1$  and  $r_2$  are the boundary points of the support of  $(L_Z(r)^2 - \beta_l)_+$ . Remark that  $x_1 < x_2$  for  $Z$  large enough by (27). Furthermore let  $n_l = N_l/(q(2l+1))$  and  $k_{l, n} = 2n$ .

Now let  $V_l$  be the potential generated by the Fermi-Hellmann equations and the given radial densities  $\rho_l$ , *i. e.*,

$$(8) \quad -V_l = \alpha_l \rho_l^2 \quad \text{for } l \leq [k] - 1$$

where  $\alpha_l = \pi^2/q^2(2l+1)^2$  and let

$$(9) \quad H_l = -\frac{d^2}{dr^2} + V_l$$

in  $L^2(0, \infty)$  with Dirichlet boundary conditions. The next lemma shows that the quantum mechanical problem involving  $V_l$  does not lower the sum of the negative eigenvalues – up to our required accuracy – for high enough angular momenta. Moreover angular momentum channels near to  $k$  may be omitted completely.

LEMMA 1. — Let  $L = [Z^\delta]$  and  $L' = [k - Z^\varepsilon]$  with  $0 < \varepsilon < 2/9, 1/9 < \delta < 1/3$ . Then

$$\sum_{l=L}^{\infty} q(2l+1) \operatorname{tr}(\tilde{H}_l)_- \geq \sum_{l=L}^{L'-1} q(2l+1) \operatorname{tr}(H_l)_- - \operatorname{Const.} Z^{\max\{4/3+3\varepsilon, 19/9-\delta\}}.$$

*Proof.* — 1.  $L' \leq l$ :

$\tilde{H}_l$  has the same negative eigenvalues as

$$-\frac{d^2}{dr^2} + \begin{cases} -\phi_l(r), & \text{if } r > 0, \\ 0, & \text{if } r \leq 0, \end{cases}$$

in  $L^2((-\infty, 0) \cup (0, \infty))$  with Dirichlet boundary conditions. Thus, using the Lieb-Thirring inequality ([7], Theorem 1) and by dropping the Dirichlet condition at zero we

obtain

$$\begin{aligned} -\operatorname{tr}(\tilde{\mathbf{H}}_l)_- &\leq \operatorname{Const.} \int (\phi_l)_+^{3/2} = \operatorname{Const.} \int_{r_1}^{r_2} \left( \frac{L_Z(r)^2 - \beta_l}{r^2} \right)^{3/2} dr \\ &\leq \operatorname{Const.} Z^{3/2} \int_{r_1}^{r_2} (L_Z(r) - \beta_l^{1/2})^{3/2} dr \end{aligned}$$

where we used (24) and  $L_Z(r) \leq \beta_k^{1/2}$ .

We cut the last integral into two pieces, namely  $r_1$  to the  $RZ^{-1/3}$ , where  $L_Z$  has its maximum, and  $RZ^{-1/3}$  to  $r_2$ . Since  $k - L' = O(Z^\varepsilon)$  the point  $r_2$  is to the left of the smallest point of inflexion of  $L_Z$  with a distance of order  $Z^{-1/3}$ . Thus  $L_Z''$  is negative on  $(r_1, r_2)$  and may be estimated from above and below by  $-\operatorname{Const.} Z$ . By expansion we have

$$\begin{aligned} L_Z'(r) &= (r - RZ^{-1/3}) L_Z''(r') \\ L_Z(r) &= \beta_k^{1/2} + \frac{1}{2} (r - RZ^{-1/3})^2 L_Z''(r'') \end{aligned}$$

with suitable points  $r', r'' \in (r_1, r_2)$ . Thus we have

$$(10) \quad \operatorname{Const.} Z^{1/2} (\beta_k^{1/2} - L_Z(r))^{1/2} \leq |L_Z'(r)| \leq \operatorname{Const.} Z^{1/2} (\beta_k^{1/2} - L_Z(r))^{1/2}$$

and therefore by a change of variables

$$-\operatorname{tr}(\tilde{\mathbf{H}}_l)_- \leq \operatorname{Const.} Z \int_{\beta_l^{1/2}}^{\beta_k^{1/2}} \frac{(x - \beta_l^{1/2})^{3/2}}{(\beta_k^{1/2} - x)^{1/2}} dx \leq \operatorname{Const.} Z (\beta_k^{1/2} - \beta_l^{1/2})^2.$$

Since  $\tilde{\mathbf{H}}_l \geq 0$  for  $l \geq k$  summation yields

$$\sum_{l=L'}^{\infty} q(2l+1) \operatorname{tr}(\tilde{\mathbf{H}}_l)_- = O(Z^{4/3+3\varepsilon}).$$

## 2. $L \leq l < L'$ :

The difference of the infima of the spectra of

$$\sum_{i=1}^{n_l^0} \left( -\frac{d^2}{dr_i^2} + V_l(r_i) \right)$$

and

$$H_{\psi_l} := \sum_{i=1}^{n_l^0} \left( -\frac{d^2}{dr_i^2} - \phi_l(r_i) \right)$$

may be estimated from above by

$$e_l = \int (\phi_l + V_l)_+ \rho_{\psi_l}$$

where  $\rho_{\psi_l}$  is the ground state density (or the densities of a minimizing sequence) of  $H_{\psi_l}$ . Thus

$$\sum_{l=L}^{L'-1} q(2l+1) \operatorname{tr}(\tilde{\mathbf{H}}_l)_- \geq \sum_{l=L}^{L'-1} q(2l+1) \operatorname{tr}(H_l)_- - \sum_{l=L}^{L'-1} q(2l+1) e_l.$$

By the Hölder and the Lieb-Thirring inequality we have

$$\begin{aligned}
 (11) \quad & \sum_{l=L}^{L'-1} q(2l+1) e_l \\
 & \leq \left( \sum_{l=0}^{\infty} q(2l+1) \int_0^{\infty} \rho_{\psi_l}^3 dr \right)^{1/3} \left( \sum_{l=L}^{L'-1} q(2l+1) \int_{(r_1, x_1) \cup (x_2, r_2)} \phi_l^{3/2} dr \right)^{2/3} \\
 & \leq \text{Const.} \left( \sum_{l=0}^{\infty} q(2l+1) T_{\psi_l} \right)^{1/3} \left( \sum_{l=L}^{L'-1} q(2l+1) \int_{(r_1, x_1) \cup (x_2, r_2)} \left( \frac{L_Z(r)^2 - \beta_l}{r^2} \right)^{3/2} \right)^{2/3}
 \end{aligned}$$

where  $T_{\psi_l}$  is the kinetic energy of  $\psi_l$ . The first factor of the right hand side can be estimated by the cubic root of the ground state kinetic energy of  $\sum_{i=1}^{N^Q} (-\Delta_i - (\phi(r_i) - \mu)_+)$

where  $N^Q = \sum_{l=0}^{\infty} q(2l+1) n_l^Q$  using the fact that we treat an (effective) one-particle operator. This total kinetic energy, however, can be bounded by a Lieb-Simon type argument by  $\text{Const.} Z^{7/3}$  (Lieb and Simon [6], Theorem III.2). The first factor on the right hand side of (11) is therefore  $O(Z^{7/9})$ . The integrals over the two intervals of the second factor of the right hand side of (11) can be estimated by

$$\begin{aligned}
 \int_{r_1}^{x_1} \frac{(L_Z(r)^2 - \beta_l)^{3/2}}{r^3} dr & \leq (L_Z(x_1)^2 - \beta_l)^{3/2} \frac{x_1 - r_1}{r_1^3} = O(Z^2 l^{-7/2}) \\
 \int_{x_2}^{r_2} \frac{(L_Z(r)^2 - \beta_l)^{3/2}}{r^3} dr & \leq (L_Z(x_2)^2 - \beta_l)^{3/2} \frac{r_2 - x_2}{x_2^3} = O(Z^{5/6})
 \end{aligned}$$

using (24)-(26), (28) and (29). Thus the left hand side of (11) is of the order of  $Z^{19/9-\delta}$ . ■

Now we introduce some notations. For  $l \leq [k] - 1$  let

$$\begin{aligned}
 A_l &= \frac{1}{n_l^2} \int_0^{\infty} \alpha_l \rho_l^3 dr \\
 B_l &= \int_0^{\infty} \sqrt{\rho_l}{}^{\prime 2} + V_l \rho_l dr
 \end{aligned}$$

and

$$C_l = \int_0^{\infty} \sqrt{\rho_l}{}^{\prime 2} dr \int_0^{\infty} -V_l \rho_l dr - \left( \int_0^{\infty} \sqrt{\rho_l} \sqrt{\rho_l}{}^{\prime} (-V_l)^{1/2} dr \right)^2.$$

Observe that  $A_l$  and  $C_l$  are positive.

Since the Macke orbitals constitute an orthonormal basis we have by Bessels equality and by partial integration in the weak sense

$$(12) \quad H_l = \sum_{n=-\infty}^{\infty} H_{l, n}$$

where

$$(13) \quad H_{l, n} = |\varphi'_{l, n}\rangle \langle \varphi'_{l, n}| - |(-V_l)^{1/2} \varphi_{l, n}\rangle \langle (-V_l)^{1/2} \varphi_{l, n}|.$$

Remark that

$$(14) \quad \lambda_{l, n} = \frac{1}{2N_l} \left\{ 4n^2 A_l + B_l - \sqrt{(4n^2 A_l + B_l)^2 + 4C_l} \right\}$$

is the only negative eigenvalue of  $H_{l, n}$ .

LEMMA 2. —  $\sum_{n=-\infty}^{\infty} \lambda_{l, n}$  is absolutely convergent and

$$\begin{aligned} & \sum_{l=L}^{L'-1} q(2l+1) \sum_{n=-\infty}^{\infty} \lambda_{l, n} \\ & \geq -\frac{2}{3} \sum_{l=L}^{L'-1} \alpha_l^{-1/2} \int_0^{\infty} \left( (\phi - \mu)_+ - \frac{l(l+1)}{r^2} \right)_+^{3/2} - \text{Const. } Z^{2-\delta/2} \log Z \end{aligned}$$

for  $L, L'$ , and  $\delta$  as in Lemma 1 but with  $\varepsilon > 1/8$ .

*Proof.* — We have for  $L \leq l < L'$  by inequalities (30) and (32)

$$\begin{aligned} 0 & \leq \frac{\int \sqrt{\rho_l}^2}{\int \alpha_l \rho_l^3} = O \left( \left( \frac{Z^2}{l^{3/2}} + Z^{17/12} l^{1/2} \right) \frac{1}{Z^{4/3} (k-l)^2} \right) \\ & = \begin{cases} O(L^{-3/2} + Z^{-5/12}) \leq 1/2, & \text{if } l \leq k/2, \\ O(Z^{3/2-4/3-2\varepsilon} + Z^{1/4-2\varepsilon}) \leq 1/2, & \text{if } l > k/2, \end{cases} \end{aligned}$$

if  $\varepsilon > 1/8$  and  $Z$  sufficiently large. Thus, by definition of  $V_l$ ,

$$(15) \quad \frac{1}{2} n_l^2 \leq \frac{-B_l}{A_l} = n_l^2 \left( 1 - \frac{\int \sqrt{\rho_l}^2}{\int \alpha_l \rho_l^3} \right) \leq n_l^2.$$

Now for  $n$  such that  $\gamma_{l, n} := ||2n| - \sqrt{-B_l/A_l}| \geq 4$  the estimate

$$(16) \quad q(2l+1) \lambda_{l, n} \geq \frac{1}{2n_l} \left\{ B_l + 4n^2 A_l - |B_l + 4n^2 A_l| - \frac{2C_l}{A_l} \frac{1}{|4n^2 + B_l/A_l|} \right\}$$

is appropriate, otherwise we use

$$(17) \quad q(2l+1)\lambda_{l, n} \geq \frac{1}{2n_l} \{B_l + 4n^2 A_l - |B_l + 4n^2 A_l| - 2\sqrt{C_l}\}.$$

By monotonicity of the function  $1/|x^2 - c^2|$  in the intervals  $(0, |c|)$  and  $(|c|, \infty)$  we estimate as follows

$$(18) \quad \frac{1}{n_l} \sum_{\substack{n=-\infty \\ \gamma_l, n \geq 4}}^{\infty} \frac{C_l}{A_l} \frac{1}{|4n^2 + B_l/A_l|} \leq \text{Const.} \frac{C_l}{n_l^2 A_l} \log n_l = O\left(\left(\frac{Z^2}{l^{3/2}} + Z^{17/12} l^{1/2}\right) \log Z\right)$$

by (15), (30), and (34).

Moreover

$$(19) \quad \frac{1}{n_l} \sum_{\substack{n=-\infty \\ \gamma_l, n < 4}}^{\infty} \sqrt{C_l} = O\left(\frac{Z^{5/3}}{l^{3/4}} + Z^{33/24} l^{1/4}\right),$$

since we have only finitely many summands and by (30), (32) and (34).

The main terms of (16) and (17) yield

$$(20) \quad -\frac{1}{n_l} \sum_{\substack{n=-\infty \\ 4n^2 < -B_l/A_l}}^{\infty} (B_l + 4n^2 A_l) = -\frac{1}{n_l} A_l \left(-4v_l^2 + 8 \sum_{n=1}^{[v_l]} (n^2 - v_l^2)\right)$$

with  $v_l = \sqrt{-B_l/4A_l}$ . Thus (20) becomes

$$(21) \quad \int \alpha_l \rho_l^3 \left(\frac{16}{3} \left(\frac{v_l}{n_l}\right)^3 + O\left(\frac{v_l}{n_l^3}\right)\right) \leq \frac{2}{3} \int \alpha_l \rho_l^3 (1 + O(n_l^{-2}))$$

$$\leq \frac{2}{3} \int \alpha_l^{-1/2} (\phi_l)_+^{3/2} + \frac{2}{3} \int_0^{x_1} \alpha_l \rho_l^3 + \frac{2}{3} \int_{x_2}^{\infty} \alpha_l \rho_l^3 + \text{Const.} n_l^{-2} \int \alpha_l \rho_l^3$$

$$\leq \frac{2}{3} \int \alpha_l^{-1/2} (\phi_l)_+^{3/2} + \text{Const.} (Z^2 l^{-5/2} + n_l^{-2} \int \alpha_l \rho_l^3)$$

using (15), (22), (23) and the Fermi-Hellmann equation. Thus, collecting terms, we obtain from (16), (17), (18), (19), and (21)

$$\sum_{n=-\infty}^{\infty} q(2l+1)\lambda_{l, n} \geq -\frac{2}{3} \int \alpha_l^{-1/2} (\phi_l)_+^{3/2} - \text{Const.} \left(\frac{Z^2 \log Z}{l^{3/2}} + Z^{17/12} l^{1/2} \log Z + n_l^{-2} \int \alpha_l \rho_l^3\right).$$

Summation over  $l$  using (31) yields the desired result. ■

The following theorem is a basic fact of our microlocalization.

THEOREM 2. — Let  $H_l$  be given as in (9) and  $\lambda_{l, n}$  as in (14). Then

$$\operatorname{tr}(H_l)_- \geq \sum_{n=-\infty}^{\infty} \lambda_{l, n}.$$

*Proof.* — Let  $\mathbf{M}$  be the set of all trace class operators  $d$  with finite kinetic energy and  $0 \leq d \leq 1$ . Then

$$\operatorname{tr}(H_l)_- = \inf \{ \operatorname{tr}(H_l d) \mid d \in \mathbf{M} \} \geq \sum_{n=-\infty}^{\infty} \inf \{ \operatorname{tr}(H_{l, n} d) \mid d \in \mathbf{M} \} = \sum_{n=-\infty}^{\infty} \operatorname{tr}(H_{l, n})_-$$

where  $H_{l, n}$  is given by (13). The claim follows now from the fact that  $\lambda_{l, n}$  is the only negative eigenvalue of  $H_{l, n}$ . ■

### 3. Application to the Scott problem

By the above arguments we are now in a position to bound the hamiltonian (1) from below.

THEOREM 3:

$$H \geq E_{\text{TF}}(1, N/Z) Z^{7/3} + \frac{q}{8} Z^2 - \text{Const. } Z^{53/27}.$$

*Proof.* — Estimating the eigenvalues of  $\tilde{H}_l$  for small  $l$ , i. e.,  $l < L$ , by the eigenvalues of  $H_l^0$  we obtain

$$\begin{aligned} \sum_{l=0}^{L-1} q(2l+1) \operatorname{tr}(\tilde{H}_l)_- &\geq -\frac{qZ^2}{2} L + \frac{q}{8} Z^2 - \text{Const. } Z^2 L^{-1} \\ &\geq -\frac{2}{3} \sum_{l=0}^{L-1} \int \alpha_l^{-1/2} \left( (\phi - \mu)_+ - \frac{(l+(1/2))^2}{r^2} \right)_+^{3/2} + \frac{q}{8} Z^2 - \text{Const. } (Z^{2-\delta} + Z^{5/3+2\delta}) \end{aligned}$$

using page 190, second but last inequality, of [9] and replacing  $Z/r$  by  $(\phi - \mu)_+$  analogously to the proof of Lemma 12, second part, of [9]. By Lemmata 1 and 2 and Theorem 2 we have for  $\varepsilon = 1/6$ .

$$\begin{aligned} \sum_{l=L}^{\infty} q(2l+1) \operatorname{tr}(\tilde{H}_l)_- &\geq -\frac{2}{3} \sum_{l=L}^{\infty} \int \alpha_l^{-1/2} \left( (\phi - \mu)_+ - \frac{(l+(1/2))^2}{r^2} \right)_+^{3/2} \\ &\quad - \text{Const. } (Z^{19/9-\delta} + (Z^{2-\delta/2} + Z^{23/12}) \log Z) \end{aligned}$$

where we used again an argument analogously to Lemma 12, first part, of [9] to substitute  $l(l+1)$  by  $(l+(1/2))^2$ .

Optimizing  $\delta$  yields  $\delta=4/27$  and the error terms of order  $Z^{53/27}$ . Then Poisson summation – generating errors of order  $Z^{5/3}$  only (see [10]) – and formula (4) yield the desired result. ■

**A. Some useful formulae**

Throughout this appendix assume  $L \leq l \leq [k] - 1$  and  $Z \gg 1$ .

By explicit integration

$$(22) \quad \alpha_l \int_0^{x_1} \rho_l^3 = O\left(\frac{Z^2}{l^{5/2}}\right),$$

$$(23) \quad \alpha_l \int_{x_2}^{\infty} \rho_l^3 = O(Z^{7/6} l).$$

The following results are easily transcribed from [10], Lemma C.1, formulae (3.9), (3.12), and (3.13), and [1], Lemma 18, 19, and 20.

$$(24) \quad \frac{\beta_l}{Z} \leq r_1 \leq \text{Const.} \frac{\beta_l}{Z},$$

$$(25) \quad \frac{\text{Const.}}{l} \leq r_2 \leq \frac{\text{Const.}}{l} \quad \text{for } \lambda \leq 0,$$

$$(26) \quad \frac{\text{Const.}}{Z^{1/3}} \leq r_2 \leq \frac{\text{Const.}}{Z^{1/3}} \quad \text{for } \lambda > 0.$$

Furthermore

$$(27) \quad x_1 \leq RZ^{-1/3} - \text{Const.} Z^{-1/2} < RZ^{-1/3} + \text{Const.} Z^{-1/2} \leq x_2,$$

$$(28) \quad L_Z(x_1)^2 - \beta_l \leq T \beta_l^{1/2},$$

$$(29) \quad L_Z(x_2)^2 - \beta_l \leq \text{Const.} Z^{1/3},$$

$$(30) \quad \int \sqrt{\rho_l}^2 = O\left(\frac{Z^2}{l^{3/2}} + Z^{17/12} l^{1/2}\right),$$

$$(31) \quad \sum_{l=L}^{[k]-1} \frac{\alpha_l \int \rho_l^3}{n_l^2} \leq \text{Const.} \sum_{l=L}^{[k]-1} \frac{Z^{4/3} (k-l)^2}{(k-l)^2} = O(Z^{5/3}).$$

Next we note

$$(32) \quad \text{Const.} Z^{4/3} (k-l)_+^2 \leq \int \alpha_l \rho_l^3 \leq \text{Const.} Z^{4/3} (k-l)_+^2.$$

*Proof.* – The upper bound is a transcription as above. The lower bound requires a separate treatment: Note the scaling  $L_{\lambda, Z}(r) = Z^{1/3} L_{\lambda, 1}(Z^{1/3} r)$ . Denote the unique maximum of  $L_{\lambda, 1}$  (see [10], Lemma A.1, and [1], Lemma 5) by R. (At this point we

make explicit the  $\lambda$  dependence of  $L_Z$ .) This yields that there are positive constants  $c_1, c_2$  such that  $L_{\lambda, z}(r)^2 \geq c_1 Z r$  for  $r \leq c_2 Z^{-1/3}$ . Moreover by [1], Lemma 3.(ii) and [10], formula (B. 4)

$$L'_{\lambda, z}(r) < L'_{0, z}(r) \leq \text{Const.} \left(\frac{Z}{r}\right)^{1/2} \quad \text{for } r \leq c_2 Z^{-1/3}$$

possibly by decreasing  $c_2$ .

For  $\beta_l \leq c_3^2 \beta_k$  with  $c_3$  positive but sufficiently small

$$(33) \quad \int \alpha_l \rho_l^3 \geq \text{Const.} l \int_{x_1}^{c_2 Z^{-1/3}} \frac{\sqrt{L_Z(r)^2 - \beta_l^3}}{r^3 L'_Z(r)} L'_Z(r) dr.$$

which may be estimated from below by

$$\text{Const.} l \int_{L_Z(x_1)}^{L_Z(c_2 Z^{-1/3})} Z^2 \frac{(y^2 - \beta_l)^{3/2}}{y^5} dy \geq \text{Const.} Z^2 \int_a^b \frac{(x^2 - 1)^{3/2}}{x^5} dx \geq Z^{4/3} (k - l)^2,$$

since by (28) for suitable  $c_3$

$$a := \frac{L_Z(x_1)}{\beta_l^{1/2}} \leq 1 + T < \frac{\text{Const.}}{c_3} \leq \frac{L_Z(c_2 Z^{-1/3})}{c_3 \beta_k^{1/2}} \leq \frac{L_Z(c_2 Z^{-1/3})}{\beta_l^{1/2}} =: b.$$

For  $\beta_l > c_3^2 \beta_k$  the integral  $\int \alpha_l \rho_l^3$  – noting  $x_1 \geq \text{Const.} Z^{-1/3}$  – may be estimated from below by

$$\text{Const.} l Z^{1/2} \int_{L_Z(x_1)}^{L_Z(RZ^{-1/3})} \frac{(L^2 - \beta_l)^{3/2}}{(\beta_k^{1/2} - L)^{1/2}} dL$$

because of (10). Moreover, since

$$c := \left(\frac{\beta_k^{1/2} - L_Z(x_1)}{\beta_k - \beta_l}\right)^{1/2} \geq (1 - T/2)^{1/2},$$

we obtain

$$\int \alpha_l \rho_l^3 \geq \text{Const.} l^{5/2} Z^{1/2} (k - l)^2 \int_0^c (1 - x^2)^{3/2} dx \geq \text{Const.} Z^{4/3} (k - l)^2,$$

which proves (32).

Finally

$$(34) \quad \text{Const.} (k - l) \leq n_l \leq \text{Const.} (k - l).$$

*Proof.* – In this case the lower bound is a transcription as above, and the upper bound requires a separate treatment: By direct integration

$$\int_0^{x_1} \rho_l + \int_{x_2}^\infty \rho_l = q(2l + 1) O(1).$$

Now we estimate the middle term: First let  $l \leq k/2$ . We split the integral at  $RZ^{-1/3}$  and the use for  $\phi(r)$  the upper bound  $Z/r$  on the left part and  $\text{Const.}/r^4$  (Sommerfeld solution) on the right part. By dropping the  $\beta_l$  one directly obtains the desired inequality. For  $k/2 \leq l$  we split the interval of integration at  $RZ^{-1/3}$  once more, but by using (10) we obtain

$$\int_{x_1}^{x_2} \rho_l \leq \text{Const.} \frac{Z^{1/3} k^{1/2}}{Z^{1/2}} \int_{\sqrt{\beta_l}}^{\sqrt{\beta_k}} \left( \frac{L - \sqrt{\beta_l}}{\sqrt{\beta_k} - L} \right)^{1/2} dL \leq \text{Const.} (k-l).$$

### Acknowledgement

We thank V. Bach, E. A. Carlen, and E. H. Lieb for various helpful discussions, V. Bach for a critical reading of the manuscript, E. H. Lieb, and B. Simon for their hospitality at Princeton University and Caltech where the present investigation started, and R. Lewis and L. Oversteegen for a pleasant stay at the University of Alabama at Birmingham where the main part of the work was done.

### REFERENCES

- [1] V. BACH, *A Proof of Scott's Conjecture for Ions*. [*Rep. Math. Phys.*, (to appear).].
- [2] F. A. BEREZIN, *Wick and Anti-Wick Operator Symbols*. (*Math. U.S.S.R. Sbornik*, Vol. 15, 1971, pp. 578-610).
- [3] W. HUGHES, *An Atomic Energy Lower Bound that Gives Scott's Correction*. (*Ph. D. thesis*, Princeton, Department of Mathematics, 1986).
- [4] W. HUGHES, *An Atomic Lower Bound that Agrees with Scott's Correction*. (*Advances in Mathematics*, 1990, pp. 213-270).
- [5] E. H. LIEB, Thomas-Fermi and Related Theories of Atoms and Molecules (*Rev. Mod. Phys.*, 53, 1981, pp. 603-604).
- [6] E. H. LIEB and B. SIMON, *The Thomas-Fermi Theory of Atoms, Molecules and Solids*. (*Adv. Math.*, Vol. 23, 1977, pp. 22-116).
- [7] E. H. LIEB and W. E. THIRRING, *Inequalities for the Moments of the Eigenvalues of the Schrödinger Hamiltonian and their Relation to Sobolev Inequalities*, in E. H. LIEB, B. SIMON and A. S. WIGHTMAN Ed., *Studies in Mathematical Physics: Essays in Honor of Valentine Bargmann*, Princeton University Press, Princeton, 1976.
- [8] H. SIEDENTOP and R. WEIKARD, *On Some Basic Properties of Density Functionals for Angular Momentum Channels*. (*Rep. Math. Phys.*, Vol. 28, 1986, pp. 193-218).
- [9] H. SIEDENTOP and R. WEIKARD, *On the Leading Correction of the Thomas-Fermi Model: Lower Bound—with an Appendix by A. M. K. Müller*. (*Invent. Math.*, 97, 1989, pp. 159-193).
- [10] H. SIEDENTOP and R. WEIKARD, *On the Leading Energy Correction for the Statistical Model of the Atom: Interacting Case*. (*Commun. Math. Phys.*, Vol. 112, 1987, pp. 471-490).

(Manuscript received April 23, 1990).

H. SIEDENTOP\*,  
Department of Mathematics,  
Princeton University,  
Princeton, NJ 08544-1000,  
U.S.A.

R. WEIKARD\*,  
Institut für Mathematische Physik,  
Technische Universität Carolo-Wilhelmina,  
Mendelssohnstraße 3,  
D-3300 Braunschweig.

\*Address as of September 1, 1990:  
Department of Mathematics  
University of Alabama at Birmingham  
Birmingham, AL 35294  
U.S.A.