BULLETIN DE LA S. M. F.

AYDIN AYTUNA

Extension operators for analytic functions defined on certain closed subvarieties of a Stein space

Bulletin de la S. M. F., tome 123, n° 2 (1995), p. 243-255

<http://www.numdam.org/item?id=BSMF_1995__123_2_243_0>

© Bulletin de la S. M. F., 1995, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (http://smf.emath.fr/Publications/Bulletin/Presentation.html) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Bull. Soc. math. France,
123, 1995, p. 243–255.

EXTENSION OPERATORS FOR ANALYTIC FUNCTIONS DEFINED ON CERTAIN CLOSED SUBVARIETIES OF A STEIN SPACE

BY

AYDIN AYTUNA (*)

RÉSUMÉ. — Soient M un espace de Stein irréductible et V une sous-variété fermée de M telle que $\mathcal{O}(V)$ soit un espace de séries de puissance. Dans cet article, nous donnons des conditions nécessaires et suffisantes pour l'existence d'un opérateur d'extension linéaire et continu de $\mathcal{O}(V)$ dans $\mathcal{O}(M)$ en termes de functions plurisous-harmoniques définies sur ces variétés. En fait, nous obtenons ces résultats en résolvant un problème d'extension plus général. Nous considérons aussi quelques conséquences de ces résultats.

ABSTRACT. — Let M be an irreducible Stein space and let V a closed subvariety of M with the property that $\mathcal{O}(V)$ is a power series space. In this paper we give a necessary and sufficient condition for the existence of a continuous linear extension operator from $\mathcal{O}(V)$ into $\mathcal{O}(M)$ in terms of plurisubharmonic functions defined on these varieties. Actually we obtain these results by solving a general lifting problem. We also consider some consequences of these results.

0. — Let M be an irreducible Stein space and V a closed subvariety of M. One of the consequences of the Oka-Cartan theory is that every analytic function on V can be extended to an analytic function on M. The question as to whether this extension process can be achieved by a continuous linear extension operator was studied by various authors.

Such a continuous operator if it exists, will imbed the Fréchet space of all analytic functions on $V, \mathcal{O}(V)$, into $\mathcal{O}(M)$ as a closed complemented subspace. In some cases this simple observation exhibits an obstruction, for the existence of a continuous linear extension operator. This situation

(*) Texte reçu le 20 août 1993.
A. AYTUNA, Department of Mathematics, 06531 Ankara, Turkey.
This work was partially supported by the Scientific and Technical Research Council of Turkey (TBAG-Ç2).

Keywords : Extension operators from subvarieties, plurisubharmonic functions.

Classification AMS : 46E10.

BULLETIN DE LA SOCIÉTÉ MATHÉMATIQUE DE FRANCE 0037–9484/1995/243/$ 5.00
© Société mathématique de France

occurs for example, when $\mathcal{O}(V)$ has no continuous norm (i.e. when V has infinite number of irreducible components) or when every continuous linear mapping from $\mathcal{O}(V)$ into $\mathcal{O}(M)$ is compact (see [9]). On the other hand positive answers in the cases :

(i) when M is a strictly pseudoconvex domain in a Stein manifold and V is of the form $V = M \cap \widetilde{V}$ where $\widetilde{V}$ is a closed submanifold near $\overline{M}$ intersecting ∂M transversally, and

(ii) when $M = \mathbb{C}^n$ and V a closed submanifold for which $\mathcal{O}(V)$, is isomorphic to $\mathcal{O}(\mathbb{C}^d)$ for some d as Fréchet spaces, e.g. when V is a smooth algebraic variety (see [17]),

were obtained in [10] by using $\overline{\partial}$-methods. In both of the cases considered above, the spaces $\mathcal{O}(V)$ turns out to belong to a well studied and well understood class of Fréchet spaces. A *power series space* is a sequence space of the form

$$\Lambda_R(\alpha) = \Big\{ x = \{x_n\}_{n=1}^{\infty} \,;\ \|x\|_r \doteq \sum_{k=1}^{\infty} |x_k|\, r^{\alpha_k} < +\infty \quad \text{for all } \ 0 < r < R \Big\}$$

where $0 < R \leq +\infty$ and $\alpha = \{\alpha_n\}$ is an increasing unbounded sequence of positive numbers. The space $\Lambda_R(\alpha)$ equipped with the norms $\| \ \|_r$, for $0 < r < R$ is a Fréchet space. It is easy to see that for a fixed α, the spaces $\Lambda_R(\alpha)$, for $0 < R < +\infty$, are all isomorphic to each other and so we have two types of power series spaces; the ones that are isomorphic to $\Lambda_1(\alpha)$, (finite type), and the ones that are isomorphic to $\Lambda_\infty(\alpha)$ (infinite type). A large number of Fréchet function spaces occuring in analysis are actually power series spaces [14]. In the case (i) considered above, $\mathcal{O}(V)$ is (isomorphic to) $\Lambda_1(n^{1/d})$ and in the case (ii) is $\Lambda_\infty(n^{1/d})$ where in both cases d is the dimension of V.

In this article we shall investigate the above mentioned question in the case when $\mathcal{O}(V)$ is a power series space. More generally we consider for a given data (M, V, W, T) consisting of a irreducible Stein space M, a subvariety V of M, a Stein space W for which $\mathcal{O}(W)$ is a power series space and a continuous linear operator T from $\mathcal{O}(W)$ into $\mathcal{O}(V)$, the problem of finding a continuous linear operator $\widetilde{T}$ such that the following diagram commutes

$$\begin{array}{ccc} & & \mathcal{O}(M) \\ & \overset{\widetilde{T}}{\nearrow} & \big\downarrow T \\ \mathcal{O}(W) & \xrightarrow{\quad R \quad} & \mathcal{O}(V) \end{array}$$

where R is the restriction operator. Observe that in the special case

$W = V$ and $T = I$ the identity of $\mathcal{O}(V), \tilde{I}$ if it exists, is a continuous linear extension operator. The obstruction to finding $\widetilde{T}$ for an arbitrary T in the above set up is due to the non vanishing of the first derived functor $\mathrm{Ext}^1(\cdot,\cdot)$ of the functor Pro in the terminology of the locally convex homological algebra developed by PALAMADOV [11] (cf. [15]). Indeed in the above set up, denoting by $I(V)$ the ideal of the variety V, the short exact sequence

$$O \to I(V) \longrightarrow \mathcal{O}(M) \xrightarrow{R} \mathcal{O}(V) \to O$$

gives rise to the exact sequence

$$\begin{aligned} 0 \to L\big(\mathcal{O}(W), I(V)\big) &\longrightarrow L\big(\mathcal{O}(W), \mathcal{O}(M)\big) \\ &\longrightarrow L\big(\mathcal{O}(W), \mathcal{O}(V)\big) \xrightarrow{\delta} \mathrm{Ext}^1\big(\mathcal{O}(W), I(V)\big) \\ &\longrightarrow \mathrm{Ext}^1\big(\mathcal{O}(W), \mathcal{O}(M)\big) \longrightarrow \mathrm{Ext}^1\big(\mathcal{O}(W), \mathcal{O}(V)\big) \to 0 \end{aligned}$$

where $L(E,F)$ denotes the space of all continuous linear operators from E into F (see [15]). For a nuclear Fréchet space E, $\mathrm{Ext}^1(E, I(V))$ can be identified with the first Čech cohomology group of the sheaf $I^{E^*}(V)$, of germs of E^* valued analytic functions on M that vanish on V (see for example [1]). Hence the possible non vanishing of Ext^1 in this case reflects the failure of the Cartan theorem (B) for E^* valued coherent analytic sheaves on M. Various conditions on the pair of Fréchet spaces which assure the vanishing of this derived functor are given in [15] (see also [1]). In particular the vanishing of $\mathrm{Ext}^1(\mathcal{O}(W), I(V))$ when $\mathcal{O}(W)$ is a power series space of infinite type follows from these general considerations (see also Remark 1). Hence in the above mentioned set up we will restrict our attention to Stein spaces W for which $\mathcal{O}(W)$ is isomorphic to a finite type power series space.

We shall use the standard terminology and notation of complex analysis as in [6], [7] except perhaps in our usage of the term Stein space. In this note by a Stein space we mean a reduced, irreducible Stein space in the sense of [6] which has a Hausdorff, separable topology.

Some results of this work was announced in [3].

1. — Returning to our problem, let us fix a Stein space M, a closed subvariety V of M and a Stein space W for which $\mathcal{O}(W)$ is a power series space. Since we will be investigating the extendibility of continuous linear operators from $\mathcal{O}(W)$ into $\mathcal{O}(V)$, we can, without loss of generality take W to be either Δ^d, the unit polydisc in $\mathbb{C}^d$, or $\mathbb{C}^d$ itself depending

upon the type of the power series space $\mathcal{O}(W)$, where $d = \dim W$. In both case a continuous linear operator T from $\mathcal{O}(W)$ into $\mathcal{O}(V)$ induces a plurisubharmonic function ρ_T on V via the formula

$$\rho_T(z) = \overline{\lim_{\xi \to z}} \; \overline{\lim_{|n| \to \infty}} \frac{\ln |T(z^n)(\xi)|}{|n|}$$

where we have used the multi index notation $z^n = z_1^{n_1} \cdots z_d^{n_d}$ for $n = (n_1, \ldots, n_d)$ and $|n| = n_1 + \cdots + n_d$. In the case when $W = \Delta^d$, it is readily seen that this plurisubharmonic function takes negative values. With the above notation we have :

THEOREM 1. — *For a continuous linear operator T from some $\mathcal{O}(\Delta^d)$ into $\mathcal{O}(V)$ the following conditions are equivalent* :

(i) *There exists a continuous linear operator $\widetilde{T} : \mathcal{O}(\Delta^d) \to \mathcal{O}(M)$ such that $R \circ \widetilde{T} = T$ where R is the restriction operator from $\mathcal{O}(M)$ onto $\mathcal{O}(V)$.*

(ii) *There exists a negative plurisubharmonic function ρ on M such that $\rho_T \leq p|V$ on V.*

Proof.

(i) $\Rightarrow$ (ii). Let

$$\rho(z) = \overline{\lim_{\xi \to z}} \; \overline{\lim_{|n| \to \infty}} \frac{\ln |\widetilde{T}(z^n)(\xi)|}{|n|}.$$

Then ρ is a plurisubharmonic function on M and in view of the fact that $\widetilde{T}$ is an extension of T one has

$$\rho_T(z) \leq \rho(z) \quad \text{for } z \in V.$$

(ii) $\Rightarrow$ (i). Using multi index notation we set $e_n \doteq z_1^{n_1} \cdots z_n^{n_d}$, $f_n \doteq T(e_n) \in \mathcal{O}(V)$ for $n = (n_1, \ldots, n_d) \in \mathbb{N}^d$. Now choose a negative plurisubharmonic function $\Phi : M \to \mathbb{R}$ with the property that

$$\overline{\lim_{\xi \to z}} \; \overline{\lim_{|n| \to \infty}} \frac{\ln |f_n(\xi)|}{|n|} = \rho_T(z) < \Phi(z)$$

for all $z \in V$. Let

$$\Omega_V = \left\{ (z, w) \in M \times \mathbb{C}^d \,;\; z \in V,\; \max_{1 \leq i \leq d} |w_i| \doteq \|w\| < \mathrm{e}^{-\Phi(z)} \right\}.$$

Fix $(z_0, w_0) \in \Omega_V$ with $\rho_T(z_0) \neq -\infty$, say $\|w_0\| < \mathrm{e}^{-\Phi(z_0) - \delta}$ for some $\delta > 0$. We choose an $\epsilon > 0$ with $2\epsilon < \delta$ and find a neighborhood $\widehat{U}_1$ of z_0 in V such that

(i) $\sup_{\xi\in\widehat{U}_1} \rho_T(z_0) < -\epsilon$,

(ii) $\sup_{\xi\in\widehat{U}_1} \Phi(\xi) \leq \Phi(z_0) + \epsilon$.

Now Hartog's lemma ([8, p. 21], cf. [12]) implies the existence of a neighborhood $U_1 \subset\subset \widehat{U}_1$ of z_0 in V such that

$$\sup_{\xi\in U_1} \frac{\ln|f_n(\xi)|}{|n|} \leq \rho_T(z_0) + \epsilon \quad \text{for } n \text{ large.}$$

Fix a neighborhood U_2 of w_0 in $\mathbb{C}^d$ such that $\sup_{w\in U_2} \|w\| < \mathrm{e}^{-\Phi(z_0)-\delta_\epsilon}$. Now let $U = U_1 \times U_2 \subseteq M \times \mathbb{C}^d$. For $(\xi, w) \in U$ we have

$$\|w\| < \mathrm{e}^{-\Phi(z_0)-\delta+\epsilon} \leq \mathrm{e}^{-\Phi(\xi)+\epsilon-\delta+\epsilon} < \mathrm{e}^{-\Phi(\xi)}$$

so $U \subseteq \Omega_V$. Moreover for large n, we have :

$$\sup_{(\xi,w)\in U} \left|f_n(\xi)\right| |w_1^{n_1} \cdots w_d^{n_d}| \leq \mathrm{e}^{|n|\{\rho_T(z_0)-\Phi(z_0)+2\epsilon-\delta\}}.$$

An estimate of this kind can also be easily obtained in the case when $\rho_T(z_0) = -\infty$. It follows that the function F defined by a locally uniformly convergent infinite series via the formula

$$F(z, w) \doteq \sum_{n\in\mathbb{N}^d} f_n(z) w^n$$

is an analytic function on Ω_V. We set :

$$\Omega_M = \left\{(z, w) \in M \times \mathbb{C}^d ;\ \|w\| < \mathrm{e}^{-\Phi(z)}\right\}.$$

Then Ω_M is a Stein space (see [5, Thm 5.4]) and Ω_V is a closed analytic subvariety of Ω_M.

In view of Cartan theorem B, there exists an analytic function G on Ω_M such that G restricted to Ω_V is F. This function can be represented in the usual way, as a convergent (uniformly on compacta of Ω_M) infinite series via the formula

$$G(z, w) = \sum_{n\in\mathbb{N}^d} a_n(z)\ w_1^{n_1} \cdots w_d^{n_d}$$

where

$$a_n(z) = \frac{1}{(2\pi i)^d} \int\cdots\int_{|\xi|=r} \frac{G(z, \xi_1, \ldots, \xi_d)}{\prod \xi_j^{n_j+1}}\, \mathrm{d}\xi_1 \cdots \mathrm{d}\xi_d,$$

with $0 < r < e^{-\Phi(z)}$ and $n \in \mathbb{N}^d$. Since for $z \in V$, one has

$$\sum_n a_n(z)\, w^n = \sum_n f_n(z) w^n$$

on the polydisc $\Delta(0, e^{-\Phi(z)})$, we conclude that $a_n(z) = f_n(z)$ for all $z \in V$ and $n \in \mathbb{N}^d$; in other words the analytic function $a_n \in \mathcal{O}(M)$ is an extension of $f_n \in \mathcal{O}(V)$ for each $n \in \mathbb{N}^d$.

Moreover, in view of the Cauchy inequalities applied to $G(z, \cdot)$, $z \in M$ we have :

$$\overline{\lim_{|n|\to\infty}} \frac{\ln |a_n(z)|}{|n|} \leq \Phi(z). \tag{1}$$

Now fix a compact set K of M and choose another compact subset $\widehat{K}$ of M, such that $K \subset \widehat{K}$. Set

$$\max_{z\in\widehat{K}} \Phi(z) \doteq -\alpha.$$

We fix an $\beta > 0$, with $\beta < \alpha$. In view of Hartog's lemma and (1) above for $|n|$ large enough we have :

$$\sup_{z\in K} \frac{\ln |a_n|}{|n|} \leq -\alpha + \beta.$$

It follows that for every compact subset K of M there exists an $R(K) < 1$ and a $C > 0$ such that :

$$\sup_{z\in K} |a_n(z)| \leq C \sup_{\|z\|\leq R(K)} |e_n(z)|. \tag{2}$$

But this means that the linear operator $\widetilde{T}$ defined from $\mathcal{O}(\Delta^d)$ into $\mathcal{O}(M)$ by the formula $\widetilde{T}(e_n) \doteq a_n$, for $n \in \mathbb{N}^d$, is a continuous operator satisfying $R \circ \widetilde{T} = T$. This finishes the proof of the THEOREM 1. □

The above result can also be interpreted as giving a description of the kernel of the operator δ appearing in the long exact sequence (1). Our next result gives a necessary and sufficient condition for this operator to be the zero operator. But first we need a lemma on the structure of plurisubharmonic functions on Stein spaces.

LEMMA 1. — *Let X be a Stein space and ρ a plurisubharmonic function on X. Then there exists a sequence $\{f_n\}_n$ of holomorphic functions on X and a sequence of integers $\{c_n\}_n$ such that*

$$\rho(z) = \overline{\lim_n} \frac{\ln|f_n(z)|}{c_n}, \qquad z \in X.$$

Proof. — First we will show that the possibility of approximating a continuous plurisubharmonic function on compact subsets by Hartog's type functions, which is well known for domains of holomorphy in $\mathbb{C}^N$, (see [9, p. 55]), is also valid for Stein spaces. To this end let us fix a continuous plurisubharmonic function ψ on X, and a holomorphically convex compact subset $K \subseteq X$. Choose a Oka-Weil domain $\mathcal{P}$, such that $K \subseteq \mathcal{P} \subset\subset X$, and fix a holomorphic mapping $\Phi : X \to \mathbb{C}^N$ such that Φ restricted to $\mathcal{P}$ is a biholomorphism onto a closed subvariety V of the unit polydisc $\Delta^N \subseteq \mathbb{C}^N$. We can think of ψ as a plurisubharmonic function on V. Arguing as in the proof of Theorem 5.3.1 of [5] we find a Stein domain Ω of Δ^N containing V and a plurisubharmonic function $\widetilde{\psi}$ on Ω such that $\widetilde{\psi}_{|V} = \psi$. Although $\widetilde{\psi}$ need not be continuous on Ω representing it on compacta as a pointwise limit of a decreasing sequence of continuous plurisubharmonic functions and observing that on $K_1 \doteq \psi(K)$ the convergence is uniform, in view of [9, p. 55] for a given $\epsilon > 0$, we can find analytic functions $f_1, \ldots, f_s$ near K, and integers $c_1, \ldots, c_s$ such that :

$$\psi(z) - \epsilon \leq \max_{1 \leq i \leq s} \frac{\ln|f_i(z)|}{c_i} \leq \psi(z) + \epsilon, \qquad \forall z \in K.$$

Now fix a point $z_0 \in K$ and choose an f_j and c_j such that :

$$\psi(z_0) - \epsilon \leq \frac{\ln|f_j(z_0)|}{c_i} \leq \psi(z_0) + \epsilon.$$

Since ψ is continuous we can find a ball U around z_0 such that :

$$\mathrm{e}^{c_j(\psi(z)-2\epsilon)} < |f_j(z)| \qquad \text{for } z \in U. \tag{3}$$

By approximating f_j on the holomorphically convex compact set $K \cup \overline{U}$ uniformly by global analytic functions we can find an $F \in \mathcal{O}(X)$ such that (3) holds with f_j replaced by F and also

$$\psi(z) + 2\epsilon \geq \log \frac{|F(z)|}{c_j}, \qquad z \in K.$$

Now cover K with balls constructed above to get for a given $\epsilon > 0$ analytic functions $F_1, \ldots, F_k$ on X and integers $c_1, \ldots, c_k$ such that :

$$\psi(z) - 2\epsilon < \max_{1 \le j \le k} \left\{ \frac{\ln |F_j(z)|}{c_j} \right\} \le \psi(z) + 2\epsilon, \qquad z \in K$$

Hence Proposition 2 of [9] is valid also for Stein spaces.

Now let ρ be a given plurisubharmonic function on X. In view of Theorem 5.5 of [5] there exists a sequence of continious plurisubharmonic functions $\{\rho_n\}$ that decrease pointwise to ρ. Choose an exhaustion of X by holomorphically convex compact sets $\{K_n\}_n$. Fix a sequence of positive numbers $\{\epsilon_n\}_n$ that decrease to zero. For each n there exists analytic functions $F_1^n, \ldots, F_{\rho(n)}^n$ and integers $c_1^n, \ldots c_{\rho(n)}^n$ such that :

$$\rho_n(z) - \epsilon_n \le \max_{1 \le i \le \rho(n)} \frac{\ln |F_i^n(z)|}{c_i^n} \le \rho_n(z) + \epsilon_n \qquad \forall z \in K_n.$$

We enumerate $\{F_i^n\}_{i,n}$ (similarly $\{c_i^n\}_{i,n}$) as

$$\left\{F_1', \ldots, F_{\rho(1)}', \ldots, F_1^n, \ldots, F_{\rho(n)}^n \cdots\right\}$$

and denote the resulting sequence by $\{G_\alpha\}_\alpha$, (similarly $\{c_\alpha\}_\alpha$). Set :

$$\gamma_\alpha(z) = \frac{\ln |G_\alpha(z)|}{c_\alpha}.$$

Now fix a point $z \in X$, say $z \in K_N$. Let $n > N$ and

$$k = \sum_{i=1}^{n-1} \rho(i) + 1.$$

Since $K_N \subset K_n$ we have

$$\rho_n(z) - \epsilon_n \le \max_{1 \le i \le p(n)} \frac{\ln |F_i^n(z)|}{c_i^n} \le \rho_n(z) + \epsilon_n.$$

Hence

$$\rho(z) - \epsilon_n \le \sup_{\alpha > k} \gamma_\alpha(z)$$

and so

$$\rho(z) - \epsilon_n \le \inf_s \sup_{\alpha > s} \gamma_\alpha(z). \tag{4}$$

On the other hand choose any α with $\alpha > k$, with k as above, then

$$\gamma_\alpha(z) = \frac{\ln|F_i^s(z)|}{c_i^s}$$

for some $s \geq n$. So we have $\gamma_\alpha(z) \leq \rho_s(z) + \epsilon_s = \rho_n(z) + \epsilon_n$; hence $\sup_{\alpha>k} \gamma_\alpha(z) \leq \rho_n(z) + \epsilon_n$. It follows that :

$$\inf_t \sup_{\alpha>t} \gamma_\alpha(z) \leq \inf_n(\rho_n(z) + \epsilon_n) = \rho(z). \tag{5}$$

So combining (4) and (5) and setting $f_n \doteq G_n$ we get :

$$\rho(z) = \overline{\lim}_n \frac{\ln|f_n(z)|}{c_n}.$$

This finishes the proof of the Lemma. □

COROLLARY 1. — *Let M be a Stein space and V a closed subvariety of M. Then the following are equivalent :*

(i) *For every Stein space W for which $\mathcal{O}(W)$ is a finite type power series space and for every continuous linear operator $T : \mathcal{O}(W) \to \mathcal{O}(V)$ there exists a continuous linear operator $\widehat{T} : \mathcal{O}(W) \to \mathcal{O}(M)$ such that $R \circ \widehat{T} = T$ where R is the restriction operator from $\mathcal{O}(M)$ into $\mathcal{O}(V)$.*

(ii) *For every negative plurisubharmonic function ρ on V there exists a negative plurisubharmonic function $\hat{\rho}$ on M such that $\rho \leq \hat{\rho}_{|V}$.*

Proof. — In view of THEOREM 1 we only need to prove the implication (i) $\Rightarrow$ (ii). To this end we fix a negative plurisubharmonic function ρ on V. In view of the LEMMA we can find a sequence $\{f_n\}_n$ of analytic functions on V, and a sequence of positive integers $\{c_n\}_n$, with $c_n \uparrow \infty$ such that :

$$\rho(z) = \overline{\lim}_n \frac{\ln|f_n(z)|}{c_n}.$$

In view of Hartog's lemma for every compact set K in V there exists a negative number α and a constant $c > 0$ such that, for all n,

$$\sup_{z\in K} |f_n(z)| \leq c\, \mathrm{e}^{\alpha c_n}. \tag{6}$$

Hence the assignment

$$T(z^n) = \begin{cases} 0 & \text{if } n \notin \{c_k\}_k, \\ f_{c_s} & \text{if } n = c_s \text{ for some } s \end{cases}$$

defines, in view of (6), a continuous linear operator $T : \mathcal{O}(\Delta) \to \mathcal{O}(V)$. We fix a $\widehat{T} : \mathcal{O}(\Delta) \to \mathcal{O}(M)$ with $\widehat{T}_{|V} = T$ and let as usual

$$\rho_{\widehat{T}}(z) = \overline{\lim_{\xi \to z}} \; \overline{\lim_{n}} \; \frac{\ln |\widehat{T}(z^n)(\xi)|}{n}.$$

Since $\rho = \rho_T$, the argument given in (i) $\Rightarrow$ (ii) of THEOREM 1 shows that $\rho \leq \rho_{\widehat{T}|V}$. This finishes the proof of COROLLARY 1. □

The above corollary can be used to characterize among the *hyperconvex varieties* V of a Stein space M (i.e. the varieties V such that $\mathcal{O}(V)$ is a finite type power series space, see [2]) the ones which admit a continuous linear extension operator $\mathcal{E} : \mathcal{O}(V) \to \mathcal{O}(M)$. Recall that for a Stein space X and a compact set $K \subset X$ the plurisubharmonic function :

$$w_K^X(z) \doteq \overline{\lim}_{\xi \to z} \sup\Big\{ u(\xi) : u \in \mathrm{PSH}(X), \; u \leq 0 \text{ on } X \text{ and } u \leq -1 \text{ on } K \Big\}$$

is called the *plurisubharmonic measure* ($\mathcal{P}$-measure) of K relative to X (see eg. [4], [13], [18]). These functions are natural complex counterparts of harmonic measures of classical potential theory. Since any negative plurisubharmonic function on a Stein space is dominated by a constant multiple of a $\mathcal{P}$-measure one can reexpress the condition (ii) above using $\mathcal{P}$-measures to obtain :

COROLLARY 2. — *Let M be a Stein space and V a hyperconvex subvariety of M. Then the following conditions are equivalent* :

(i) *There exists a continuous linear extension operator*

$$\mathcal{E} : \mathcal{O}(V) \longrightarrow \mathcal{O}(M).$$

(ii) *There exists compact sets $K \subseteq V$, $S \subseteq M$ with non empty interiors and a constant $C > 0$ such that* :

$$w_K^V \leq C \, w_S^M{}_{|V}.$$

REMARKS.

(i) Although we have chosen to treat the case when $\mathcal{O}(W)$ is isomorphic to an infinite type power series space by making use of some general considerations, we note that the line of reasoning given in the proof of THEOREM 1 can also be used in this case. Indeed the existence of

an operator $\widehat{T} : \mathcal{O}(W) \to \mathcal{O}(M)$ with $R \circ \widehat{T} = T$ for any $T : \mathcal{O}(W) \to \mathcal{O}(V)$ can be deduced, in this case, from the fact that for any plurisubharmonic function ρ on V there exists a plurisubharmonic function $\hat{\rho}$ on M such that $\rho \leq \hat{\rho}_{|V}$.

(ii) In the case when $\mathcal{O}(M)$ is isomorphic to an infinite type power series space and when W is hyperconvex, THEOREM 1 characterizes the operators T for which such a $\widehat{T}$ exists as the ones for which $\sup_{z \in V} \rho_T(z) < 0$. This family is precisely the family of all *compact operators* from $\mathcal{O}(W)$ into $\mathcal{O}(V)$. This can also be derived from the general extention properties of compact operators and the fact that every continuous operator from a finite type power series space into an infinite type power series space is compact.

(iii) For a smoothly bounded relatively compact domain D with C^2 boundary in a Stein manifold and a negative plurisubharmonic function ρ on D one has that

$$\rho(z) < C\{-d(z, \partial D)\}, \qquad z \in D$$

for some constant $C > 0$ where $d(z, \partial D)$ is the distance of z from ∂D (see [10, Lemma 3.2]). Hence in the case when D is given by $D = \{z : u(z) < 0\}$, for some C^2 plurisubharmonic function u defined in a neighborhood of $\overline{D}$, we have that any negative plurisubharmonic function on D is dominated by a positive constant multiple of u, since $-d(\,\cdot\,, \partial D)$ is dominated by a positive constant multiple of u. This property remains valid for submanifolds of D of the form $D \cap M'$ where M' is a closed complex submanifold in a neighborhood of $\overline{D}$ which intersects ∂D transversally since in this case $D \cap M' = \{z \in M' : u(z) < 0\}$. Now combining Corollary 5 of [2] with Corollary 2 above we obtain the following slight generalization of Theorem 4.2 of [10].

COROLLARY 3. — *Let M be a Stein manifold and $D \subset\subset M$ a smoothly bounded domain in M of the form $D = \{z : u(z) < 0\}$ for some C^2 plurisubharmonic function defined in a neighborhood of $\overline{D}$. For a complex manifold M' in a neighborhood of $\overline{D}$ which intersects ∂D transversally there exists a continuous linear extension operator $\mathcal{E} : \mathcal{O}(D \cap M') \to \mathcal{O}(D)$.*

Even if we drop the transversality condition in the above corollary we can still get some information about the class of continuous linear operators $T : \mathcal{O}(\Delta^d) \to \mathcal{O}(D \cap M')$ which admit a continuous linear extension operator, namely these are preciselsy the operators for which $\rho_T \leq Cu$ on $D \cap M'$ for some $C > 0$. This observation can be used in constructing concrete operators for which no such $\widehat{T}$ exists. For example following

Example 5.3 of [10], let

$$D = \{(z\,;w) \in \mathbb{C}^2\,;\ |z|^2 + |w-1|^2 < 1\}$$

and

$$M' = \{(z,w) \in \mathbb{C}^2\,;\ w = z^2\}.$$

Then the operator $T : \mathcal{O}(\Delta) \to \mathcal{O}(D \cap M')$ defined as $T(f)(z,w) \doteq f(\mathrm{e}^{-z^3})$ admits no extension operator $\widehat{T} : \mathcal{O}(\Delta) \to \mathcal{O}(D)$, since, an easy computation shows the impossibility of finding a $C > 0$ satisfying

$$\rho_T(z,w) = \ln|\mathrm{e}^{-z^3}| \leq C\{|z|^2 + |w-1|^2 - 1\}.$$

BIBLIOGRAPHIE

[1] AYTUNA (A.). — *On the vector valued Cousin Problem*, Preprint.

[2] AYTUNA (A.). — *On Stein manifolds M for which $\mathcal{O}(M)$ is isomorphic to $\mathcal{O}(\Delta^n)$ as Fréchet spaces*, Manuscripta Math., t. **62**, 1988, p. 297–315.

[3] AYTUNA (A.). — *Stein spaces M for which $\mathcal{O}(M)$ is isomorphic to a power series space*, Advances in the theory of Fréchet spaces (ed. T. Terzioğlu), Kluwer Academic Publishers, 1989, p. 115–149.

[4] BEDFORD (E.) and TAYLOR (B.A.). — *The Dirichlet problem for a complex Monge-Ampère equation*, Invent. Math., t. **37**, 1976, p. 1–44.

[5] FORNAESS (J.E.) and NARASIMHAN (R.). — *The Levi problem on complex spaces with singularities*, Math. Ann., t. **248**, 1980, p. 47–72.

[6] GRAUERT (H.) and REMMERT (R.). — *Coherent Analytic Sheaves*. — Springer-Verlag, Berlin-Heidelberg-New York, Tokyo, 1984.

[7] GUNNING (R.) and ROSSI (H.). — *Analytic Functions of Several Complex Variables*. — Englewood Cliffs, 1965.

[8] HÖRMANDER (L.). — *An Introduction to Complex Analysis in Several Variables*. — North-Holland Publishing Company, 1973.

[9] LELONG (P.). — *Plurisubharmonic Functions and Positive Differential Forms*. — New York, Gordon and Breach, 1969.

[10] MITIAGIN (B.S.) and HENKIN (G.M.). — *Linear problems of complex analysis*, Russian Math. Surveys, t. **26**, 1971, p. 99–164.

[11] PALAMODOV (V.P.). — *Homological methods in the theory of locally convex spaces*, Russian Math. Surveys, t. **26,1**, 1971, p. 1–64.

[12] SADULLAEV (A.). — *An estimate for polynomials on analytic sets*, Math. USSR Izvestiya, t. **20**, 1983, p. 493–502.

[13] SADULLAEV (A.). — *Plurisubharmonic measures and capacities on complex manifolds*, Russian Math. Surveys, t. **36,4**, 1981, p. 61–119.

[14] VOGT (D.). — *Sequence space representations of spaces of test functions and distributions*, Advances in Functional Analysis, Holomorphy and Approximation Theory (ed. G.I. Zapata). New York–Basel 1983, p. 405–443.

[15] VOGT (D.). — *On the functors* $\mathrm{Ext}^1(E, F)$ *for Fréchet spaces*, Studia Math., t. **85**, 1987, p. 163–197.

[16] VOGT (D.) and WAGNER (M.J.). — *Charakterisierung der Unterräume und Quotientenräume der nuklearen stabilen Potenzreihenräume von unendlichem Typ*, Studia Math., t. **70**, 1981, p. 65–80.

[17] ZAHARIUTA (V.P.). — *Spaces of analytic functions on algebraic varieties in* $\mathbb{C}^n$, Izv. Severo-Kavkaz. Nauchn. Tsentia Vyssh. Shkoly Ser. Estestv. Nauk, t. **4**, 1977, p. 52–55.

[18] ZAHARIUTA (V.P.). — *External plurisubharmonic functions, Hilbert scales, and the isomorphism of spaces of analytic functions of several variables*, I, II. Theor. Funkeii Funkcional Anal. i. Prilozen. Vyp., t. **19**, 1974, p. 133–157 and t. **21**, 1974, p. 65–83 (Russian).