ANNALES DE L'INSTITUT FOURIER

EDOARDO BALLICO

A splitting theorem for the Kupka component of a foliation of $\mathbb{CP}^n$, $n \geq 6$. Addendum to an addendum to a paper by Calvo-Andrade and Soares

Annales de l'institut Fourier, tome 49, n° 4 (1999), p. 1423-1425

<http://www.numdam.org/item?id=AIF_1999__49_4_1423_0>

© Annales de l'institut Fourier, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (http://annalif.ujf-grenoble.fr/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Ann. Inst. Fourier, Grenoble
49, 4 (1999), 1423–1425

A SPLITTING THEOREM FOR THE KUPKA COMPONENT OF A FOLIATION OF $\mathbb{CP}^n$, $n \geq 6$.

Addendum to an addendum to a paper by Calvo-Andrade and Soares

by Edoardo BALLICO

A codimension one singular holomorphic foliation F of $\mathbb{CP}^n$ is given by $\omega \in H^0(\mathbb{CP}^n, \Omega(k))$ (for some k) with $\omega \neq 0$, ω not vanishing on a hypersurface. The Kupka subset $K(F) := \{P \in \mathbb{CP}^n : \omega(P) = 0, d\omega(P) \neq 0\}$ of the singular set $S(F) := \{P \in \mathbb{CP}^n : \omega(P) = 0\}$ of F has remarkable properties (e.g. if not empty it is a smooth submanifold of pure codimension 2 with strong stability properties with respect to deformations of F). For much more on this topic, see [GL] and [GS]. Let $K \neq \emptyset$ be a Kupka component of F, i.e. ([CS]) a connected component of $K(F)$. It was proved in [CL] that if K is a complete intersection, then F has a meromorphic first integral. Motivated by this result in [CS] it was conjectured and proved in some cases that every Kupka component, K, is a complete intersection. This conjecture was proved in [B] under the assumption that $\deg(K)$ is not a square. Here we will remove this restrictive asssumption and prove the following result.

THEOREM. — *Let F be a codimension 1 singular holomorphic foliation of $\mathbb{CP}^n$, $n \geq 6$, induced by $\omega \in H^0(\mathbb{CP}^n, \Omega^1(k))$ and such that the*

Keywords: Singular foliations – Codimension 1 foliations – Kupka component – Complete intersection – Unstable vector bundle – Rank 2 vector bundle – Splitting of a vector bundle – Meromorphic first integral – Barth-Lefschetz theorems.
Math. classification: 58F18 – 14F05 – 14M07.

codimension 2 component of the singular set of F contains a Kupka component K. Then K is a complete intersection and F has a meromorphic first integral.

Let N_K be the normal bundle of K in $\mathbb{CP}^n$. By [CS], Corollary 3.5, N_K is the restriction $E|K$ to K of a rank 2 vector bundle E on $\mathbb{CP}^n$. K is a complete intersection if and only if E is the direct sum of two line bundles ([OSS]). If $n \geq 6$ every line bundle on K is the restriction of a line bundle on $\mathbb{CP}^n$ (see [FL]). Hence, by a very nice result of Faltings ([F]) if $n \geq 6$ and N_K is the direct sum of two line bundles, K is a complete intersection. Now we use the notations of [GL], page 320. There are two complex numbers (not both zero) such that the linear part of the foliation near each point of K depends from these two complex numbers. Up to a normalization we may assume that one of these numbers is 1; call μ the other one. Il $\mu \neq \pm 1$, then N_K splits into the direct sum of two line bundles ([GL]), Remark 2 at p. 320). If $\mu = -1$, then there is a two-to-one unramified covering $\pi : Z \to K$ such that $\pi^*(N_K)$ is the direct sum of two isomorphic line bundles (see [GL], the two lines before eq. (2.6)). Since K is simply connected ([FL]), Cor. 6.3), N_K splits into the direct sum of two isomorphic line bundles. Now assume $\mu = 1$. This is the only difference with respect to [B]. By [GL], eq. (2.12) and the two following lines, N_K is an extension of a line bundle L_1 by itself. Since $n \geq 6$ we have $H^1(K, \mathbb{C}) \cong \mathbb{C}$ by a theorem of Barth's (see e.g. [FL], eq. (**) at p. 83). Hence by standard Hodge theory ([GH], bottom of p. 105) we have $H^1(K, \mathcal{O}_K) = 0$. Thus any such extension of L_1, by itself splits and hence $N_K \cong L_1 \oplus L_1$, concluding the proof of the theorem.

The author was partially supported by MURST and GNSAGA of CNR (Italy).

BIBLIOGRAPHY

[B] E. BALLICO, A splitting theorem for the Kupka component of a foliation of $\mathbb{CP}^n$, $n \geq 6$. Addendum to a paper by Calvo-Andrade and Soares, Ann. Inst. Fourier, 45–4 (1995), 1119–1121.

[CS] O. CALVO-ANDRADE, M. SOARES, Chern numbers of a Kupka component, Ann. Inst. Fourier, 44–4 (1994), 1219–1236.

[CL] D. CERVEAU, A. LINS, Codimension one foliations in $\mathbb{CP}^n$ $n \geq 3$, with Kupka components, in Complex analytic methods in dinamical systems, Astérisque, (1994), 93–133.

[F] G. FALTINGS, Ein Kriterium für vollständige Durchsnitte, Invent. Math., 62 (1981), 393–401.

[FL] W. FULTON, R. LAZARSFELD, Connectivity in algebraic geometry, in Algebraic Geometry, Proceedings Chicago 1980, Lect. Notes in Math. 862, Springer-Verlag (1981), 26–92.

[GN] X. GOMEZ-MONT, A. LINS-NETO, A structural stability of foliations with a meromorphic firs integral, Topology, 30 (1990), 315–334.

[GH] Ph. GRIFFITHS, J. HARRIS, Principles of Algebraic Geometry, John Wiley & Sons, 1978.

[OSS] Ch. OKONEK, M. SCHNEIDER, H. SPINDLER, Vector Bundles on Complex Projective spaces, Progress in Math., 3, Birkhäuser, Basel, 1978.

Manuscrit reçu le 1er octobre 1998,
accepté le 12 février 1999.

Edoardo BALLICO,
Department of Mathematics
University of Trento
38050 Povo (NT) (Italy).
ballico@science.unitn.it