ANNALES DE L'INSTITUT FOURIER

LENNART CARLESON

Maximal functions and capacities

Annales de l'institut Fourier, tome 15, n° 1 (1965), p. 59-64

<http://www.numdam.org/item?id=AIF_1965__15_1_59_0>

© Annales de l'institut Fourier, 1965, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (http://annalif.ujf-grenoble.fr/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Ann. Inst. Fourier, Grenoble
15, 1 (1965), 59-64.

MAXIMAL FUNCTIONS AND CAPACITIES

by Lennart CARLESON

---

1. Let $f(x)$ be periodic with period 2π and assume $f(x) \in L^p(-\pi, \pi)$, some $p \geqslant 1$. The maximal function $f^*(x)$ associated with $f(x)$ was introduced by Hardy and Littlewood through the definition

$$f^*(x) = \sup_t \frac{1}{t} \int_x^{x+t} f(u)\, du. \tag{1.1}$$

The inequalities

$$\int_{-\pi}^{\pi} |f^*(x)|^p\, dx \leqslant A_p \int_{-\pi}^{\pi} |f(x)|^p\, dx, \qquad p > 1, \tag{1.2}$$

and

$$m\{x | f^*(x) \geqslant \lambda\} \leqslant \frac{A}{\lambda} \int_{-\pi}^{\pi} |f(x)|\, dx \tag{1.3}$$

are basic in the theory of differentiation. (1.2) can alternatively be given as a theorem on harmonic functions. Assume $f > 0$ and let $u(z)$ be harmonic in $|z| < 1$ with boundary values $f(\theta)$. Then clearly

$$\text{const.} f^*(\theta) \leqslant \sup_r u(re^{i\theta}) \leqslant \text{const.} f^*(\theta). \tag{1.4}$$

The inequality (1.2) follows if we can characterize those non-negative measures μ for which

$$\iint_{|z|<1} u(z)^p\, d\mu(z) \leqslant A_p \int_{-\pi}^{\pi} f(x)^p\, dx. \tag{1.5}$$

It is sufficient to consider $p = 2$ and the complete solution was given in [3]: a necessary and sufficient condition on μ, is

$$\mu(S) \leqslant \text{const.}\, s$$

for every set $S: 1 - s < |z| < 1$, $|\arg(z) - \alpha| < s$.

The corresponding linear problem, i.e., to describe those μ for which

$$\iint u(z)\, d\mu(z) \tag{1.6}$$

is bounded for $f \in L^p$ is clearly much simpler and the solution is that

$$\varphi(\theta) = \iint \frac{1 - |z|^2}{|e^{i\theta} - z|^2} \, d\mu(z) \tag{1.7}$$

belongs to L^q.

Although this result is in principle sufficient for differentiation purposes, it is of little help since no simple geometric characterization of μ seems to be available.

We shall now consider the corresponding problem for the class of functions $f(x)$,

$$f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx},$$

such that

$$\|f\|_K^2 = \Sigma |c_n|^2 \lambda_{|n|} < \infty.$$

Here $\{\lambda_n\}$ is a positive sequence such that

$$K(x) \sim \sum_{0}^{\infty} \frac{\cos nx}{\lambda_n}$$

is a convex function $\in L^1$. The following theorem is quite easy to prove.

THEOREM 1.— *If* $\lambda_n = (n+1)^{1-\alpha}$, $0 \leqslant \alpha < 1$, (1.6) *is bounded if and only if*

$$E_\alpha(\mu) = \iint \frac{d\mu(a)\, d\mu(b)}{|1 - \bar{a}b|^\alpha} < \infty, \qquad 0 < \alpha < 1,$$

$$E_0(\mu) = \iint \log \left| \frac{1}{1 - \bar{a}b} \right| d\mu(a)\, d\mu(b) < \infty, \qquad \alpha = 0.$$

The bound of (1.6) *is* $\leqslant$ const. $\sqrt{E_\alpha}$.

If we specialize $d\mu$ to have the form $d\sigma(\theta)$ placed at a point on the radius from 0 to $e^{i\theta}$ we find using (1.4) and observing that $E_\alpha(\mu)$ essentially increases if we push the masses out to $|z| = 1$

$$\left(\int f^*(x)\, d\sigma(x) \right)^2 \leqslant A_\alpha \|f\|_K^2 I_\alpha(\sigma) \tag{1.8}$$

where I_α is the energy of σ with respect to the kernel $|x|^{-\alpha}$, resp. $\log \frac{1}{|x|}$. This inequality implies easily the existence of derivatives

and boundary-values except on sets of capacities zero. This is a result by Beurling [1] and Broman [2].

The proof of Theorem 1 in the case $\alpha = 0$ is particularly simple. Consider first the case when μ has its support strictly inside $|z| < 1$. Consider the harmonic function

$$u_0(z) = \iint \log|1 - z\bar{\zeta}|\, d\mu(\zeta)$$

and let (u, v) denote scalar product in the space of harmonic functions with finite Dirichletintegral and with $u(0) = 0$. Then by Poisson's formula

$$(u, u_0) = \int_{|z|=1} u \frac{\partial u_0}{\partial n}\, ds = 2\pi \iint u(z)\, d\mu(z).$$

Hence

$$2\pi \left| \iint u\, d\mu \right| \leqq \| u_0 \| \,.\, \| u \|$$

with equality if $u = u_0$, and the linear functional (1.6) has norm $(2\pi)^{-\frac{1}{2}}\sqrt{E_0(\mu)}$. The case of a general μ follows immediately.

The restriction $u(0) = 0$, i.e., $\int f\, dx = 0$, is clearly inessential. Let us also observe that we here (as well as in Section 2) also may restrict ourselves to $f > 0$ since $|f(x)|$ has a smaller norm than f (see (2.1)).

In the case $0 < \alpha < 1$ we write

$$\int u(a)\, d\mu(a) = \int f(\theta)\, d\theta \frac{1}{2\pi} \int \frac{1 - |a|^2}{|e^{i\theta} - a|^2}\, d\mu(a) = \int f(\theta) g(\theta)\, d\theta.$$

The function $v(r, \theta)$ harmonic in $|z| < 1$ with boundary values $g(\theta)$ is

$$v(r, \theta) = \frac{1}{2\pi} \int \frac{1 - |a|^2 r^2}{|e^{i\theta} - ar|^2}\, d\mu(a) = \Sigma\, b_n r^{|n|} e^{in\theta}. \tag{1.9}$$

We wish to prove

$$\iint v(r, \theta)^2 (1 - r)^{-\alpha}\, dr\, d\theta < \infty, \tag{1.10}$$

since this inequality is equivalent to $\Sigma |b_n|^2 (|n| + 1)^{\alpha - 1} < \infty$. Inserting (1.9) in (1.10) we see that (1.10) holds if $E_\alpha(\mu) < \infty$.

2. It is clearly possible to use the same method for general kernels $K(x)$ and corresponding weights λ_n. However the formulas become so involved that they cannot be used to deduce inequalities of the form (1.8). Of particular interest is the case

$$\lambda_n = (\log(n + 2))^\alpha, \qquad 0 < \alpha < \infty.$$

For functions f with corresponding $\|f\|_K$ finite and $0 < \alpha < 1$, nothing is known on convergence of Fourier series and no better result on existence of derivatives than Lebesgue's theorem. The kernel K_α that is associated with this sequence is

$$K_\alpha(x) \sim \frac{1}{|x|(\log 1/|x|)^{1+\alpha}}, \qquad x \to 0.$$

The following theorem holds

THEOREM 2. — *There is a constant* B_α *such that*

$$C_{K_\alpha}\left[\{x|f^*(x) \geqslant \lambda\}\right] \leqslant \frac{B_\alpha}{\lambda^2}\|f\|^2_{K_\alpha}, \qquad 0 < \alpha < \infty.$$

By standard methods this implies that the primitive function of f has a derivative except on a set of K_α-capacity zero. It is interesting to compare this result with what is known on convergence of Fourier series. It has been proved by Temko [4], that if $\|f\|_{K_{\alpha+1}} < \infty$ then the Fourier series converges except on a set of K_α-capacity zero, while we here get a stronger result on existence of boundary values.

In the proof we use the equivalent norm

$$(2.1) \qquad \int\int_{-\pi}^{\pi} \frac{|f(x) - f(y)|^2}{\varphi(x-y)}\,dx\,dy, \qquad \varphi(t) = |t|\left(\log\frac{8}{|t|}\right)^{1-\alpha}$$

and the following potential theoretic lemma:

LEMMA. — *If* σ *is an interval of length* d *on* $(-\pi, \pi)$, *denote by* $T\sigma$ *an interval of length* $3d$ *and having the same midpoint as* σ. *We assume that* $\{\sigma_\nu\}$ *are disjoint and denote by* $E = \cup\sigma_\nu$ *and* $E' = \cup T\sigma_\nu$. *Then there is a constant* Q *only depending on* K *such that*

$$C_K(E') \leqslant QC_K(E)$$

provided $K(x) = O(K(2x))$, $x \to 0$.

In an outline, the proof of theorem 2 proceeds as follows. Let $\sigma_{\nu n}$ denote the 2^n disjoint intervals of length $2\pi \cdot 2^{-n}$ on $(-\pi, \pi)$. Let λ be given and denote by $M_\alpha(f)$ the mean value of f over the interval α. We choose intervals $\sigma_1, \sigma_2, \ldots$, such that

$$(2.2) \qquad M\sigma_\nu(f) \geqq \lambda$$

by first choosing those $\sigma_{\nu 1}$ that satisfy (2.2), then $\sigma_{\mu 2}$ disjoint from those chosen before, etc. It follows easily from the lemma that it is sufficient to prove $C\{\cup\sigma_\nu\} \leqslant$ const. $\|f\|^2 \cdot \lambda^{-2}$.

Let τ_ν be intervals such that $T\tau_\nu = \sigma_\nu$. We want to construct $f_1(x)$ such that $\|f_1\| \leqslant$ const. $\|f\|$ and $f_1(x) \equiv M\sigma_\nu(f)$, $x \in \tau_\nu$. We first modify f on each σ_ν according to the following rule where we have normalized σ_ν to $(-1, 1)$:

$$f_2(x) = \begin{cases} f(2x), & -\tfrac{1}{2} < x < \tfrac{1}{2} \\ f(-x - \tfrac{3}{2}), & -\tfrac{3}{4} < x < -\tfrac{1}{2} \\ f(x), & -1 < x < -\tfrac{3}{4} \\ \text{analogously on } (\tfrac{1}{2}, 1). \end{cases}$$

Outside $\cup\sigma_\nu$ we define $f_2(x) = f(x)$. From (2.1) it follows that $\|f_2\|_K \leqslant$ const. $\|f\|_K$.

Let 4δ be the length of the shortest of the intervals σ_ν. We have the following picture:



where we construct α_i and β_i until their length $< \delta$. α_i and β_i have lengths $= 3^{-i-1}$ (length σ_ν). We define

$$f_1(x) = \begin{cases} M_{\tau_\nu}(f_2) = M_{\sigma_\nu}(f), & x \in \tau_\nu ; \\ M_{\alpha_i}(f_2), & x \in \alpha_i ; \\ M_{\beta_i}(f_2), & x \in \beta_i ; \\ \text{linear between the intervals.} \end{cases}$$

We do the same construction on each σ_ν and each complementary interval. A computation in (2.1) shows that $\|f_1\| <$ const. $\|f_2\|$.

To complete the proof, let μ be a distribution of unit mass on $E'' = \cup\tau_\nu$. Then

$$\lambda \leqslant \int_{E''} f_2(x)\, d\mu(x) \leqslant \|f_2\|_K \cdot I_K(\mu)^{\frac{1}{2}} \leqslant \text{const.} \|f\|_K \cdot I_K(\mu)^{\frac{1}{2}}.$$

The lemma now yields theorem 2.

BIBLIOGRAPHY

[1] A. BEURLING, Ensembles exceptionnels, *Acta Math.*, 72 (1940), 1-13.
[2] A. BROMAN, On two classes of trigonometrical series, Thesis, Uppsala (1947).
[3] L. CARLESON, Interpolations by bounded analytic functions and the Corona problem, *Ann. of Math.*, 76 (1962) 547-559.
[4] K. V. TEMKO, Convex capacity and Fourier series, *Dokl. Akad. Nauk*, 110 (1956).

Lennart CARLESON,
Department of Mathematics,
Uppsala University,
Sysslomansgatan 8,
Uppsala (Suède).