ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

PAUL H. RABINOWITZ

On nontrivial solutions of a semilinear wave equation

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 8, n° 4 (1981), p. 647-657

<http://www.numdam.org/item?id=ASNSP_1981_4_8_4_647_0>

© Scuola Normale Superiore, Pisa, 1981, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (http://www.sns.it/it/edizioni/riviste/annaliscienze/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

On Nontrivial Solutions of a Semilinear Wave Equation (*).

PAUL H. RABINOWITZ

The question of the existence of nontrivial time periodic solutions of autonomous or forced semilinear wave equations has been the object of considerable recent interest [1-12]. These papers study the equation

$$u_{tt} - u_{xx} + f(x, u) = 0\,, \qquad 0 < x < l \tag{1.1}$$

(or its analogue where f also depends on t in a time periodic fashion) together with boundary conditions in x and periodicity conditions in t. In particular the following result was proved in [11, Theorem 3.37 and Corollary 4.14]:

THEOREM 1.2. *Let* $f \in C([0, l] \times \boldsymbol{R}, \boldsymbol{R})$ *and satisfy*

(f_1) $f(x, 0) = 0$ *and* $f(x, r)$ *is strictly monotone increasing in* r,

(f_2) $f(x, r) = o(|r|)$ *at* $r = 0$,

(f_t) *there are constants* $\bar{r} > 0$ *and* $\mu > 2$ *such that*

$$0 < \mu F(x, r) \leqslant r f(x, r)$$

for $|r| \geqslant \bar{r}$ *and* $x \in [0, l]$ *where*

$$F(x, r) = \int_0^r f(x, s)\, ds\,.$$

(*) This research was sponsored in part by the Office of Naval Research under Contract No. N00014-76-C-0300.

Pervenuto alla Redazione il 22 Settembre 1980.

Then for any T which is a rational multiple of l, equation (1.1) *possesses a non trivial continuous weak solution u satisfying*

$$\begin{cases} u(0, t) = 0 = u(l, t) , \\ u(x, t + T) = u(x, t) . \end{cases} \tag{1.3}$$

Furthermore $f \in C^k$ implies $u \in C^k$.

As part of the proof of Theorem 1.2, it was shown that the functional

$$I(u) = \int_0^T \int_0^l [\tfrac{1}{2}(u_t^2 - u_x^2) - F(x, u)]\, dx\, dt \tag{1.4}$$

defined on the class of functions satisfying (1.3) (and of which (1.1) is formally the Euler equation) has a positive critical value. Therefore (f_1) and the form of I imply that $u_t \not\equiv 0$ for the corresponding critical point u. Thus u is nonconstant in x and must depend explicitly on t. It was further observed in [11] (Theorem 5.24 and Remark 5.25) that if g satisfies (f_1)-(f_3) the equation

$$u_{tt} - u_{xx} - g(x, u) = 0 , \qquad 0 < x < l \tag{1.5}$$

together with (1.3) also possesses a nontrivial weak solution. Indeed the arguments of Theorem 1.2 go through with minor modifications to establish this fact. However the functional one studies for this case is

$$J(u) = \int_0^T \int_0^l [\tfrac{1}{2}(u_x^2 - u_t^2) - G(x, u)]\, dx\, dt \tag{1.6}$$

where G is the primitive of g. Again the positivity of $J(u)$ for a critical point u implies u is nonconstant but we can no longer conclude that u depends explicitly on t. In fact it is known [13, 14] that as a consequence of (f_2)-(f_3), the ordinary differential equation boundary value problem

$$-\frac{d^2u}{dx^2} = g(x, u) , \qquad u(0) = 0 = u(l) \tag{1.7}$$

has an unbounded sequence of solutions which can be characterized by the number of zeros they possess in $(0, l)$.

Our goal in this paper is to show that if (f_3) is strengthened somewhat, (1.5), (1.3) possesses infinitely many time dependent solutions. More precisely we will prove:

THEOREM 1.8. *Let $g \in ([0, l]\times \boldsymbol{R}, \boldsymbol{R})$ and suppose g satisfies (f_1)-(f_2) and $(\tilde{f}_3)$. There is a constant $\mu > 0$ such that*

$$0 < \mu F(x, r) \leqslant rf(x, r)$$

for all $r \neq 0$.

Then for any $T \in l\boldsymbol{Q}$ there is a $k_0 \in \boldsymbol{N}$ such that for all $k \geqslant k_0$, (1.5), (1.3) possesses a solution u_k which is kT periodic in t and $\partial u_k/\partial t \not\equiv 0$. Moreover infinitely many of the functions u_k are distinct.

REMARK 1.9. We have no estimate for the size of k_0 and do not know if the result is false in general for $k = 1$. Note also that since (1.5) is an autonomous equation with respect to t, whenever $u(x, t)$ is a solution, so is $u(x, t + \theta)$ for any $\theta \in \boldsymbol{R}$. The above statement about the u_k's being distinct means in particular that they do not differ by merely a translation in time.

The proof of Theorem 1.8 draws on several results from [11] and ideas from [12]. For convenience we will take $l = \pi$ and $T = 2\pi$. Choosing $k \in \boldsymbol{N}$, we seek a solution u_k of (1.5) which is $2\pi k$ periodic in t and $\partial u_k/\partial t \not\equiv 0$. Making the change of time scale $\tau = t/k$, the period becomes 2π again and the problem to be solved is

$$\begin{cases} U_{\tau\tau} - k^2(U_{xx} + g(x, U)) = 0\,, & 0 < x < \pi \\ U(0, \tau) = 0 = U(l, \tau) \\ U(x, \tau + 2\pi) = U(x, \tau) \end{cases} \tag{1.10}$$

with $U(x, \tau) = u(x, t)$.

For the convenience of the reader and to set the stage for a key estimate, the argument used in [11] to establish the existence of nontrivial solutions of (1.10) will be sketched quickly. Solutions are obtained by an approximation argument. To begin (1.10) is modified in two ways. The wave operator $\partial^2/\partial\tau^2 - k^2(\partial^2/\partial x^2)$ possesses an infinite dimensional null space in the class of functions satisfying the boundary and periodicity conditions of (1.10) and given by

$$N = \operatorname{span}\{\sin jx \sin kj\tau, \sin jx \cos kj\tau \,|\, j \in \boldsymbol{N}\}\,.$$

The fact that N is infinite dimensional complicates the analysis of (1.10) and to introduce some compactness to the problem in N, we perturb (1.10) by adding a βV_{tt} term to the left hand side of the equation. Here $\beta > 0$ and V is the (L^2 orthogonal) projection of U onto N. A second difficulty in treating (1.10) arises due to the unrestricted rate of growth of $g(x, r)$ as

$|r| \to \infty$. We get around this by truncating g. More precisely $g(x, r)$ is replaced by $g_K(x, r)$ which coincides with g for $|r| \leqslant K$ and grows cubically at ∞ [11]. Thus (1.10) is replaced by the modified problem

$$\text{(1.11)} \qquad \begin{cases} U_{\tau\tau} + \beta V_{\tau\tau} - k^2(U_{xx} + g_K(x, U)) = 0, & 0 < x < \pi \\ U(0, \tau) = 0 = U(l, \tau) \\ U(x, \tau + 2\pi) = U(x, \tau) \end{cases}$$

where g_K satisfies (f_1), (f_2), $(\bar{f}_3)$ with a new constant $\bar{\mu} = \min(\mu, 4)$.

Letting G_K denote the primitive of g_K, in a formal fashion (1.11) can be interpreted as the Euler equation arising from the functional

$$\text{(1.12)} \qquad J(U; k, \beta, K) = \int_0^{2\pi} \int_0^{\pi} \left[\frac{k^2}{2} U_x^2 - \frac{1}{2} U_\tau^2 - \frac{\beta}{2} V_\tau^2 - k^2 G_K(x, U) \right] dx\, d\tau .$$

Let

$$E_m \equiv \operatorname{span}\{\sin jx \sin n\tau, \sin jx \cos b\tau \,|\, 0 \leqslant j,\ n \leqslant m\} .$$

The strategy pursued in [11] was to find a critical point U_{mk} of $J|_{E_m}$, let $m \to \infty$, and then let $\beta \to 0$ to get a solution U_k of (1.10) with g replaced by g_K. Then L^∞ bounds for U_k independent of K show if we choose $K(k)$ sufficiently large $g_K(x, U_k) = g(x, U_k)$ so (1.10) obtains. A separate comparison argument is required to prove that $U_k \not\equiv 0$.

The first step in carrying out the details of the above argument involves obtaining an upper bound M_k for $c_{mk} \equiv J(U_{mk}; k, \beta, K)$ with M_k independent of m, β, and K. For the current problem which also depends on k, it is crucial to know the behavior of M_k as a function of k. Thus we will take a closer look at c_{mk} and use a variant of an argument of [12]. By Lemma 1.13 of [11], c_{mk} can be characterized in a minimax fashion. We will not write down this characterization explicitly but will note a consequence of it which in turn provides an upper bound for c_{mk}. Set

$$\Phi_{mk} = \operatorname{span}\{\sin jx \sin n\tau,\ \sin jx \cos n\tau \,|\, 0 \leqslant j,\ n \leqslant m \text{ and } n^2 \geqslant j^2 k^2\}$$

and

$$\psi_k = a_k \sin x \sin(k-1)\tau$$

where $a_k = \sqrt{2}/\pi$ so $\|\psi_k\|_{L^2} = 1$. Set $\Psi_{mk} = \Phi_{mk} \oplus \operatorname{span} \psi_k$. Then by Lemma 1.3. of [11]

$$\text{(1.13)} \qquad 0 < c_{mk} \leqslant \max_{u \in \Psi_{mk}} J(u; k, \beta, K) .$$

Inequality (1.13) will lead to a suitable choice for M_k. Note that by $(\bar{f}_3)$ (or even (f_3)), there are constants α_1, $\alpha_2 \geqslant 0$ and independent of K such that

$$G_K(x, r) \geqslant a_1 |r|^{\bar{\mu}} - a_2 \tag{1.14}$$

for all $r \in \boldsymbol{R}$, $x \in [0, \pi]$. Consequently $J \to -\infty$ as $u \to \infty$ in Ψ_{mk} (under $\|\cdot\|_{L^2}$) so there is a point $z \equiv Z_{mk}$ at which the maximum in (1.13) is achieved. Writing

$$z = \|z\|_{L^2}(\gamma\xi + \delta\psi_k) \tag{1.15}$$

where $\xi \in \Phi_{mk}$, $\|\xi\|_{L^2} = 1$, and $\gamma^2 + \delta^2 = 1$ and substituting (1.15) into (1.13) gives

$$k^2 \int_0^{2\pi}\int_0^{\pi} G(x, z)\, dx\, dt \leqslant \frac{1}{2}\int_0^{2\pi}\int_0^{\pi}(k^2 z_x^2 - z_\tau^2)\, dx\, d\tau \leqslant \tag{1.16}$$

$$\leqslant \frac{\delta^2}{2}\|z\|_{L^2}^2 \int_0^{2\pi}\int_0^{\pi}[k^2(\psi_k)_x^2 - (\psi_k)_\tau^2]\, dx\, d\tau \leqslant k\|z\|_{L^2}^2 .$$

Combining (1.14) and (1.16) shows that

$$k^2(\alpha_1\|z\|_{L^{\bar{\mu}}}^{\bar{\mu}} - \alpha_3) \leqslant k\|z\|_{L^2}^2 . \tag{1.17}$$

Applying the Hölder inequality yields

$$\|z\|_{L^2} \leqslant A \tag{1.18}$$

where A is a constant independent of m, k, β, K. Hence by (1.13), (1.18), and the form of J,

$$c_{mk} \leqslant Mk \tag{1.19}$$

for a constant M independent of m, k, β, K.

Letting $m \to \infty$ and then $\beta \to 0$, and formalizing what we have just shown gives:

LEMMA 1.20. *Under the hypotheses of Theorem* 1.8 *(with* $l = \pi$ *and* $T = 2\pi$*), for all* $k \in \boldsymbol{N}$*, there exists a solution* U_k *of* (1.10) *satisfying*

$$c_k \equiv J(U_k;\, k, 0, K) \leqslant Mk \tag{1.21}$$

with M *independent of* k *and* K.

It remains to show that for all k sufficiently large, $\partial U_k/\partial t \not\equiv 0$ and infinitely many of the functions $u_k(x,t) = U_k(x,\tau)$ are distinct. If U_k is independent of τ for any subsequence of k's tending to ∞, $U_k = U_k(x)$ is a classical solution of (1.7). Thus by (1.21) with $K = K(k)$ suitably large,

$$c_k = 2\pi k^2 \int_0^\pi \left[\frac{1}{2}\left|\frac{dU_k}{dx}\right|^2 - G(x, U_k)\right] dx\,. \tag{1.22}$$

By (1.7),

$$\int_0^\pi \left|\frac{dU_k}{dx}\right|^2 dx = \int_0^\pi U_k(x)\, g(x, U_k(x))\, dx\,. \tag{1.23}$$

Combining (1.21)-(1.23) yields

$$\int_0^\pi [\tfrac{1}{2} U_k g(x, U_k) - G(x, U_k)]\, dx \to 0 \tag{1.24}$$

as $k \to \infty$ along this subsequence. Moreover by $(\bar{f}_3)$,

$$\int_0^\pi \left[\frac{1}{2} U_k g(x, U_k) - G(x, U_k)\right] dx \geqslant \int_0^\pi \left(\frac{1}{2} - \frac{1}{\bar{\mu}}\right) U_k g(x, U_k)\, dx\,. \tag{1.25}$$

Thus $U_k g(x, U_k) \to 0$ in L^1. From (1.23) again we conclude that $dU_k/dx \to 0$ in L^2 which easily implies $U_k \to 0$ in L^∞. By (f_2), for any $\varepsilon > 0$, there is a $\delta > 0$ such that $|r| \leqslant \delta$ implies $|g(x,r)| \leqslant \varepsilon r$. Choosing $\varepsilon < 1/\pi$ and k large enough so that $\|U_k\|_{L^\infty} \leqslant \delta$, (1.23) then shows

$$\left\|\frac{dU_k}{dx}\right\|_{L^2}^2 \leqslant \varepsilon \|U_k\|_{L^2}^2 \leqslant \pi\varepsilon \left\|\frac{dU_k}{dx}\right\|_{L^2}^2 < \left\|\frac{dU_k}{dx}\right\|_{L^2}^2, \tag{1.26}$$

a contradiction. Consequently U_k depends on τ for all large k.

To prove the second assertion of Theorem 1.8, suppose two functions $U_k(x,\tau)$, $U_j(x,\tau)$ correspond to the same function of x, t modulo a translation in time (keeping Remark 1.9 in mind). Thus $U_k(x,\tau) = U_k(x, t/k) \equiv U(x,t)$ and $U_j(x,\tau) = U_j(x, t/j) = U(x, t+\theta)$ for some $\theta \in \boldsymbol{R}$ or $U_k(x,\tau) = U(x, k\tau)$, $U_j(x,\tau) = U(x, j\tau + \theta)$. Since U must be both $2\pi k$ and $2\pi j$ periodic in t, letting σ denote the greatest common divisor of j and k, we have $j = \sigma\bar{j}$,

$k = \sigma\bar{k}$ and U has period $2\pi\sigma$ in t. Furthermore

$$c_k = \int_0^{2\pi}\int_0^{\pi}\left[\frac{k^2}{2}U_{kx}^2 - \frac{1}{2}U_{k\tau}^2 - k^2 G(x, U_k)\right]dx\,d\tau = \tag{1.27}$$

$$= k\int_0^{2\pi k}\int_0^{\pi}\left[\frac{1}{2}(U_x^2 - U_t^2) - G(x, U)\right]dx\,dt =$$

$$= \frac{k^2}{\sigma}\int_0^{2\pi\sigma}\int_0^{\pi}\left[\frac{1}{2}(U_x^2 - U_t^2) - G(x, U)\right]dx\,dt \equiv \frac{k^2}{\sigma}b$$

and similarly

$$c_j = \frac{j^2}{\sigma}b\,. \tag{1.28}$$

Consequently if there were a sequence of solutions U_{k_i} of (1.10) corresponding to the same function U (up to a translation in t), by (1.27)-(1.28) we have

$$c_{k_i} = \frac{k_j^2}{\sigma}b \tag{1.29}$$

and $c_{k_i} \to \infty$ like k_i^2 along this sequence contrary to (1.19). Thus at most finitely many functions $U_k(x, \tau)$ correspond to the same solution $u_k(x, t)$ of (1.5), (1.3) and infinitely many of the functions u_k must be time dependent solutions of (1.5), (1.3). The proof of Theorem 1.8 is complete.

REMARK 1.30. Both the existence assertions from [11] and the arguments given above use hypothesis (f_2) which requires that g vanish more rapidly than linearly at 0. However this condition can be weakened. The simplest such generalization would be to replace $g(x, r)$ by $\alpha r + g(x, r)$ with α a constant and for this case we have:

THEOREM 1.31. *Let g satisfy (f_1), (f_2), $(\bar{f}_3)$ and let $\alpha > 0$. Then for all $T \in l\boldsymbol{Q}$, there exists a $k_0 \in \boldsymbol{N}$ such that for all $k \geqslant k_0$, the problem*

$$\begin{cases} u_{tt} - u_{xx} - \alpha u - g(x, u) = 0 \qquad 0 < x < l \\ u(0, t) = 0 = u(l, t) \\ u(x, t + kT) = u(x, t) \end{cases} \tag{1.32}$$

has a continuous weak solution u_k which is kT periodic in t and $\partial u_k/\partial t \not\equiv 0$. Moreover infinitely many of these functions are distinct.

PROOF. For convenience we again take $l = \pi$, $T = 2\pi$. It was shown in [11] that Theorem 1.2 carries over to (1.32) for $\alpha > 0$. It is also easy to see that the argument of Lemma 1.20 will give (1.21) for this setting. Likewise (1.27)-(1.29) are unaffected by the α term. Thus we get Theorem 1.31 provided that we can show $U_k(x, \tau)$ depends on τ for all large k. If not, the analogues of (1.22)-(1.23) here are

$$c_k = 2\pi k^2 \int_0^\pi \left[\frac{1}{2}\left|\frac{dU_k}{dx}\right|^2 - \frac{1}{2}\alpha U_k^2 - G(x, U_k)\right] dx \tag{1.33}$$

and

$$\int_0^\pi \left|\frac{dU_k}{dx}\right|^2 dx = \int_0^\pi (\alpha U_k^2 + U_k g(x, U_k))\, dx\,. \tag{1.34}$$

Thus (1.19), (1.33)-(1.34), and $(\tilde{f}_3)$ show that $U_k g(x, U_k) \to 0$ in L^1 as $k \to \infty$ as in (1.24)-(1.25). Since

$$\|g(x, U_k)\|_{L^1} \leqslant \pi \max_{0 \leqslant x \leqslant \pi, |r| \leqslant 1} |g(x, r)| + \|U_k g(x, U_k)\|_{L^1} \tag{1.35}$$

and the right hand side of (1.35) is uniformly bounded in k, it follows from (1.7) that the functions d^2U_k/dx^2 are uniformly bounded in L^1. The boundary conditions $U_k(0) = 0 = U_k(\pi)$ imply that there is $x_k \in (0, \pi)$ such that $(dU_k/dx)(x_k) = 0$. Hence

$$\frac{dU_k}{dx} = \int_{x_k}^x \frac{d^2U_k(\xi)}{d\xi^2}\, d\xi$$

which implies that

$$\left\|\frac{dU_k}{dx}\right\|_{L^\infty} \leqslant \left\|\frac{d^2U_k}{dx^2}\right\|_{L^1}. \tag{1.36}$$

Thus the functions U_k, dU_k/dx are bounded in L^∞ and by (1.7) again, so are d^2U_k/dx^2. It follows that a subsequence of U_k converges (in $\|\cdot\|_{C^2}$) to a solution U of (1.7) as $k \to \infty$. But (f_1) and $\|U_k g(x, U_k)\|_{L^1} \to 0$ as $k \to \infty$ imply $U \equiv 0$.

Next observe that (1.7) can be written as

$$U_k(x) = \int_0^\pi H(x, y)\big(\alpha U_k(y) + g(y, U_k(y))\big)\, dy \tag{1.37}$$

where H is the Green's function for $-d^2/dx^2$ under the boundary condition $U(0) = 0 = U(\pi)$. Dividing (1.37) by $\|U_k\|_{C^1}$ gives

$$\frac{U_k(x)}{\|U_k\|_{C^1}} = \int_0^\pi H(x, y)\left(\alpha \frac{U_k(y)}{\|U_k\|_{C^1}} + \frac{g(x, U_k(y))}{\|U_k\|_{C^1}}\right)dy . \tag{1.38}$$

By (f_2), the arguments of the integral operator are uniformly bounded in C^1. Hence since this operator is compact from C^1 to C^2, by (f_2) again a subsequence of $U_k/\|U_k\|_{C^1}$ converge to V satisfying $\|V\|_{C^1} = 1$ and

$$V(x) = \alpha \int_0^\pi H(x, y) V(y)\, dy \tag{1.39}$$

or equivalently

$$-V'' = \alpha V \qquad 0 < x < \pi; \qquad V(0) = 0 = V(\pi) . \tag{1.40}$$

If α is not an eigenvalue of $-d^2/dx^2$ under these boundary conditions we have a contradiction and the proof is complete. Thus suppose α is an eigenvalue. Consider the eigenvalue problems:

$$-z'' = \lambda \alpha z , \qquad 0 < x < \pi; \qquad z(0) = 0 = z(\pi) \tag{1.41}$$

$$-y'' = \mu\left(\alpha + \frac{g(x, \varphi)}{\varphi}\right) y , \qquad 0 < x < \pi; \qquad y(0) = 0 = y(\pi) \tag{1.42}$$

where φ is C^1 on $[0, \pi]$. Let λ_j (resp. $\mu_j(\varphi)$) denote the j-th eigenvalue of (1.41) (resp. (1.42)), the eigenvalues being ordered according to increasing magnitude. As is well known any eigenfunction corresponding to λ_m or $\mu_m(\varphi)$ belongs to

$$S_m = \{\varphi \in C^1([0, \pi], \boldsymbol{R}) \,|\, \varphi(0) = 0 = \varphi(\pi),\ \varphi \text{ has exactly } m - 1 \text{ zeros in } (0, \pi), \text{ and } \varphi' \neq 0 \text{ at all zeros of } \varphi \text{ in } [0, \pi]\} .$$

(Indeed the eigenvalues of (1.41) are $\lambda_m = m^2\alpha^{-1}$ and corresponding eigenfunctions are multiples of $\sin mx$). Since $g(x, \varphi)\varphi^{-1} \geqslant 0$ via (f_1), we have $\lambda_j \geqslant \varphi_j(\varphi)$ for all $j \in \boldsymbol{N}$ and $\varphi \in C^1$, $\varphi \not\equiv 0$ via a standard comparison theorem [15, Chapter 6]. By (1.40), 1 is an eigenvalue of (1.41), say $1 = \lambda_m$ and $V \in S_m$. Thus $\mu_m(\varphi) \leqslant 1$ and since S_m open (in the C^1 topology) and $U_k/\|U_k\|_{C^1} \to V$ in C^1 along some subsequence, it follows that $U_k/\|U_k\|_{C^1}$ and therefore U_k belongs to S_m for all large k in this subsequence. Writ-

ing (1.7) as

$$(1.43) \qquad -U_k'' = \left(\alpha + \frac{g(x, U_k)}{U_k}\right) U_k, \qquad 0 < x < \pi; \qquad U_k(0) = 0 = U_k(\pi),$$

we see $\mu_m(U_k) = 1$. By (f_1) again, $g(x, U_k)U_k^{-1} > 0$ except at the $m+1$ zeros of U_k. An examination of the proof of the Sturm Comparison Theorem [16, pp. 208-209] then shows U_k has a zero between each pair of successive zeros of V. Consequently $U_k \in S_{m+1}$, a contradiction. Thus Theorem 1.31 is established.

REMARK 1.44. In [5], Brezis, Coron and Nirenberg study (1.1), (1.3) replacing (f_3) by

$$(f_4) \quad \tfrac{1}{2}rf(r) - V(r) \geqslant \beta|f(r)| - \gamma$$

and

$$(f_5) \quad f(r)/r \to \infty \quad \text{as } |r| \to \infty$$

(and with no analogue of (f_2)). If we use (f_4)-(f_5) with x dependent f in place of (f_3), it is not difficult to see that the proof of [11] carries over for this case as does Lemma 1.20 and (1.27)-(1.29). Thus we obtain a variant of Theorem 1.8 for this case once it is established that $U_k(x, \tau)$ depends on τ for large k. To do this, we argue as in the proof of Theorem 1.8. Assume (f_4) holds with $\gamma = 0$. Then by (1.25) and (f_4), $\|g(x, U_k)\|_{L^1} \to 0$ as $k \to \infty$. This in turn implies $\|U_k\|_{L^\infty} \to 0$ via (1.7) and (1.36). Hence (1.26) again provides a contradiction.

It is also possible for us to drop (f_2) and even the requirement that $f(x, 0) = 0$ in (f_1) but then a new existence mechanism is required and we shall not carry out the details here.

REFERENCES

[1] H. AMANN, *Saddle points and multiple solutions of differential equations*, Math. Z., **169** (1979), pp. 127-166.

[2] H. AMANN - E. ZEHNDER, *Nontrivial solutions for a class of nonresonance problems and applications to nonlinear differential equations*, Ann. Scuola Norm. Sup. Pisa, 4, 7 (1980), pp. 539-603.

[3] A. BAHRI - H. BREZIS, *Periodic solutions of a nonlinear wave equation*, to appear Proc. Roy. Soc. Edinburgh.

[4] H. BREZIS - J. M. CORON, *Periodic solutions of nonlinear wave equations and Hamiltonian systems*, preprint.

[5] H. BREZIS - J. M. CORON - L. NIRENBERG, *Free vibrations for a nonlinear wave equation and a theorem of P. Rabinowitz*, preprint.

[6] H. BREZIS - L. NIRENBERG, *Characterizations of the ranges of some nonlinear operators and applications to boundary value problems*, Ann. Scuola Norm. Sup. Pisa, 4, **5** (1978), pp. 225-326.

[7] H. BREZIS - L. NIRENBERG, *Forced vibrations for a nonlinear wave equation*, Comm. Pure Appl. Math., **31** (1978), pp. 1-30.

[8] J. M. CORON, *Résolution de l'équation $Au + Bu = f$ où A est linéaire auto-adjoint et B déduit d'un potential convexe*, to appear Ann. Fac. Sc. Toulouse.

[9] G. MANCINI, *Periodic solutions of some semi-linear autonomous wave equations*, Boll. Un. Mat. Ital. (5), **15**-B (1978), pp. 649-672.

[10] P. H. RABINOWITZ, *Some minimax theorems and applications to nonlinear partial differential equations*, Nonlinear Analysis: A collection of papers in honor of Erich H. Rothe, Academic Press (1978), pp. 161-177.

[11] P. H. RABINOWITZ, *Free vibrations for a semilinear wave equation*, Comm. Pure Appl. Math., **31** (1978), pp. 31-68.

[12] P. H. RABINOWITZ, *Subharmonic solutions of a forced wave equation*, to appear Amer. J. Math.

[13] R. E. L. TURNER, *Superlinear Sturm-Liouville problems*, J. Differential Equations, **13** (1973), pp. 157-171.

[14] P. H. RABINOWITZ, *Some aspects of nonlinear eigenvalue problems*, Rocky Mountain J. Math., **3** (1973), pp. 161-202.

[15] R. COURANT - D. HILBERT, *Methods of Mathematical Physics*, vol. I, Interscience (1953).

[16] E. A. CODDINGTON - N. LEVINSON, *Theory of Ordinary Differential Equations*, McGraw-Hill (1955).

University of Wisconsin
Department of Mathematics
Madison, Wisconsin 53706