COMPOSITIO MATHEMATICA

H. JACQUET
J. A. SHALIKA

Hecke theory for $GL(3)$

Compositio Mathematica, tome 29, n° 1 (1974), p. 75-87
<http://www.numdam.org/item?id=CM_1974__29_1_75_0>

© Foundation Compositio Mathematica, 1974, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (http://http://www.compositio.nl/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

COMPOSITIO MATHEMATICA, Vol. 29, Fasc. 1, 1974, pag. 75–87
Noordhoff International Publishing
Printed in the Netherlands

HECKE THEORY FOR *GL*(3)

H. Jacquet [1] and J. A. Shalika [2]

In view of the recent results of Gelfand, Kajdan, and one of the authors ([1], [6]), it appears likely that the results of [2] – the Hecke theory – will extend to all groups $GL(p)$. In this note, we present nearly complete results for the case $p = 3$. To avoid technical difficulties we restrict ourselves to the case of a function field.

1. Global computations

Let F be a commutative field, G (resp. G') the group $GL(p)$ (resp. $GL(p-1)$) regarded as an algebraic group defined over F. We regard G' as imbedded into G by the map

$$g \to \begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix}.$$

We denote by Z (resp. Z') the center of G (resp. G') and by ${}^t g$ the transpose of a matrix g. The entries of a matrix g in G are written as g_{ij}, the first index being the column index.

In this section we take F to be a function field whose field of constants has cardinality Q. We let $\mathbb{A}$ be the ring of adeles of F and $\mathbb{I}$ the group of ideles. Then $G_{\mathbb{A}} = GL(p, \mathbb{A})$ is a locally compact group of which $G_F = GL(p, F)$ is a discrete subgroup. A cusp form on $G_{\mathbb{A}}$ is a complex-valued function ϕ, which satisfies the following three conditions:

(1.1) for all $g \in G_{\mathbb{A}}$, $a \in Z_{\mathbb{A}} = \mathbb{I}$, and $\gamma \in G_F$, $\phi(a\gamma g) = \omega(a)\phi(g)$,

where ω is a quasi-character of $\mathbb{I}/F^\times$,

(1.2) the function ϕ is invariant on the right by a compact open subgroup of $G_{\mathbb{A}}$,

(1.3) if P is a proper F-parabolic subgroup of G and U its unipotent radical, then

$$\int_{U_F \backslash U_{\mathbb{A}}} \phi(ug)\,du = 0, \qquad \text{for all } g \in G_{\mathbb{A}}.$$

[1] Research partially supported by NSF-GP27952.
[2] Research partially supported by NSF-GP31348.

Let N be the group of p by p upper triangular matrices whose diagonal entries are one. Let $N' = N \cap G'$ be the corresponding subgroup of G'. Choose a non-trivial character ψ of $\mathbb{A}/F$ and define a character θ of $N_{\mathbb{A}}$ by the formula

$$\theta(n) = \prod_{1 \leqq i \leqq p-1} \psi(n_{i+1,i}). \tag{1.4}$$

Let ϕ be a cusp form on $G_{\mathbb{A}}$. The function

$$W(g) = \int_{N_F \backslash N_{\mathbb{A}}} \phi(ng)\bar{\theta}(n)\,\mathrm{d}n \tag{1.5}$$

is called the 'Whittaker function' attached to ϕ. It transforms on the left according to the formula

$$W(ng) = \theta(n)W(g), \qquad \text{for } n \in N_{\mathbb{A}},$$

and the form ϕ has the following Fourier expansion:

$$\phi(g) = \sum_{\gamma \in N_F \backslash G_F} W(\gamma g). \tag{1.6}$$

Now let ϕ' be a cusp form on $G'_{\mathbb{A}}$ and let us compute the integral

$$\int_{G'_F \backslash G'_{\mathbb{A}}} \phi(g)\phi'(g)|\det g|^s \,\mathrm{d}g, \tag{1.7}$$

where $s \in \mathbb{C}$ and $\mathrm{d}g$ is an invariant measure on $G'_F \backslash G'_{\mathbb{A}}$. Replacing ϕ by (1.6), we obtain

$$\begin{aligned}
\int_{G'_F \backslash G'_{\mathbb{A}}} \sum_{N'_F \backslash G'_F} W(\gamma g)\phi'(g)|\det g|^s \,\mathrm{d}g &= \int_{G'_F \backslash G'_{\mathbb{A}}} \sum_{N'_F \backslash G'_F} W(\gamma g)\phi'(\gamma g)|\det(\gamma g)|^s \,\mathrm{d}g \\
&= \int_{N'_F \backslash G'_{\mathbb{A}}} W(g)\phi'(g)|\det g|^s \,\mathrm{d}g \\
&= \int_{N'_{\mathbb{A}} \backslash G'_{\mathbb{A}}} |\det g|^s \,\mathrm{d}g \int_{N'_F \backslash N'_{\mathbb{A}}} W(ng)\phi'(ng)\,\mathrm{d}n \\
&\quad \int_{N'_{\mathbb{A}} \backslash G'_{\mathbb{A}}} |\det g|^s W(g)\,\mathrm{d}g \int_{N'_F \backslash N'_{\mathbb{A}}} \theta(n)\phi'(ng)\,\mathrm{d}n.
\end{aligned}$$

Let W' be the Whittaker function attached to ϕ' and ε the $p-1$ by $p-1$ matrix defined by

$$\varepsilon_{ij} = \delta_{i,j}(-1)^i.$$

Since $\theta(\varepsilon n \varepsilon^{-1}) = \theta(n^{-1})$, in the last line the inner integral is actually $W'(\varepsilon g)$. Thus (1.7) is equal to

$$\int_{N'_{\mathbb{A}}\backslash G'_{\mathbb{A}}} W(g)W'(\varepsilon g)|\det g|^s \mathrm{d}g. \tag{1.8}$$

More precisely, for Res sufficiently large, all the above integrals converge and are equal.

Since ϕ and ϕ' are compactly supported modulo $G_F Z_{\mathbb{A}}$ and $G'_F Z'_{\mathbb{A}}$ respectively, the integral (1.7) is always convergent. Hence, the integral (1.8), which converges only in a half-space, is, as a function of s, a polynomial in Q^{-s}, Q^s.

Now, let w be the element of G_F defined by

$$w_{ij} = 0 \quad \text{if } i+j \neq p+1, \; w_{p+1-i,i} = (-1)^{i-1} \quad (1 \leqq i, j \leqq p)$$

and let w' be the corresponding element of G'_F. Clearly

$$wNw^{-1} = {}^tN, \qquad \theta(w{}^tn^{-1}w^{-1}) = \theta(n).$$

In particular, the functions $\tilde{\phi}$ and $\tilde{\phi}'$ defined by

$$\tilde{\phi}(g) = \phi(w{}^tg^{-1}) = \phi({}^tg^{-1}), \qquad \tilde{\phi}(g) = \phi'(w'{}^tg^{-1}) = \phi'({}^tg^{-1})$$

are automorphic forms on $G_{\mathbb{A}}$ and $G'_{\mathbb{A}}$ respectively, whose Whittaker functions are the functions $\tilde{W}$ and $\tilde{W}'$ given by

$$\tilde{W}(g) = W(w{}^tg^{-1}), \qquad \tilde{W}'(g) = W'(w'{}^tg^{-1}).$$

Changing g into ${}^tg^{-1}$ in the integral (1.7), we easily obtain that (1.7) is also equal to the following integral:

$$\int_{G_F\backslash G'_{\mathbb{A}}} \tilde{\phi}(g)\tilde{\phi}'(g)|\det g|^{-s} \mathrm{d}g, \tag{1.9}$$

and conclude that (1.8) and the integral

$$\int_{N'_{\mathbb{A}}\backslash G'_{\mathbb{A}}} \tilde{W}(g)\tilde{W}'(g)|\det g|^{-s} \mathrm{d}g \tag{1.10}$$

are equal in the sense of analytic continuation – as polynomials in Q^{-s}, Q^s.

2. Local conjectures

From now on and until Section 5, the field F will be a non-archimedean local field whose residual field has q elements. We denote by $|x|$ or $\alpha_F(x)$ or simply $\alpha(x)$ the module of an x in F. We denote by $M(p \times r, F)$ the space of matrices with p columns and r rows whose entries belong to F, and by $\mathscr{S}(p \times r, F)$ the space of locally constant compactly supported complex-valued functions on $M(p \times r, F)$.

Fix a non-trivial character ψ of F and let θ be the character of N_F defined by (1.4). Let π be an irreducible admissible representation of G_F. We say that π is non-degenerate if it can be realized by right translations in a space $\mathcal{W}$ of functions W on G_F satisfying

$$W(ng) = \theta(n)W(g), \qquad \text{for all } n \text{ in } N_F,\ g \text{ in } G_F.$$

If π is non-degenerate the space $\mathcal{W}$ is unique and denoted by $\mathcal{W}(\pi, \psi)$. It is called the Whittaker model of π ([1], [6]).

Since ${}^tg$ and g are always conjugate in G_F, the representation $\tilde{\pi}$ contragredient to π is equivalent to the representation $g \to \pi({}^tg^{-1})$ ([1]). In particular, if π is non-degenerate so is $\tilde{\pi}$. More precisely, if W belongs to $\mathcal{W}(\pi, \psi)$ then the function $\tilde{W}$ defined by

$$\tilde{W}(g) = W(w\,{}^tg^{-1})$$

belongs to $\mathcal{W}(\tilde{\pi}, \psi)$.

Let π and π' be non-degenerate representations of G_F and G'_F respectively. For $W \in \mathcal{W}(\pi, \psi)$ and $W' \in \mathcal{W}(\pi', \psi)$ and $s \in \mathbb{C}$, we set

$$\Psi(s, W, W') = \int_{N'_F \backslash G'_F} W(g)W'(\varepsilon g)|\det g|^{s-\frac{1}{2}}\,dg.$$

The global computations of Section 1 lead to the formulation of the following conjectures.

(2.1) For Res sufficiently large the integrals $\Psi(s, W, W')$ and $\Psi(s, \tilde{W}, \tilde{W}')$ are absolutely convergent.

(2.2) They are rational functions of q^{-s}. More precisely, for $W \in \mathcal{W}(\pi, \psi)$ and $W' \in \mathcal{W}(\pi', \psi)$ the integrals $\Psi(s, W, W')$ span a fractional ideal $\mathbb{C}[q^{-s}, q^s]L(s, \pi \times \pi')$ of the ring $\mathbb{C}[q^{-s}, q^s]$; the factor $L(s, \pi \times \pi')$ has the form $1/P(q^{-s})$ where $P \in \mathbb{C}[X]$ and $P(0) = 1$.

There is a similar factor $L(s, \tilde{\pi} \times \tilde{\pi}')$.

(2.3) There is a factor $\varepsilon(s, \pi \times \pi', \psi)$ of the form cq^{-ns} such that

$$\Psi(1-s, \tilde{W}, \tilde{W}')/L(1-s, \tilde{\pi} \times \tilde{\pi}') = \varepsilon(s, \pi \times \pi', \psi)\omega'(-1)^p\Psi(s, W, W')/L(s, \pi \times \pi')$$

for $W \in \mathcal{W}(\pi, \psi)$ and $W' \in \mathcal{W}(\pi', \psi)$, where ω' is the quasi-character of $F^\times$ such that $\pi'(a1_{p-1}) = \omega'(a) \cdot 1$.

Let us emphasize that in the above statements the pairs (π, π') and $(\tilde{\pi}, \tilde{\pi}')$ play symmetrical roles, since, in fact, $\tilde{W}$ (resp. $\tilde{W}'$) belongs to $\mathcal{W}(\tilde{\pi}, \psi)$(resp. $\mathcal{W}(\tilde{\pi}', \psi)$) if W (resp. W') belongs to $\mathcal{W}(\pi, \psi)$(resp. $\mathcal{W}(\tilde{\pi}', \psi)$).

These conjectures have been proved for $p = 2$ (see below), and, in [1], for all p under the additional assumption that π is supercuspidal. Indeed,

under this assumption, the restriction of a W in $\mathcal{W}(\pi, \psi)$ to G'_F belongs to the space $C_c^\infty(G', \theta)$ of all maps f from G'_F to $\mathbb{C}$ which, on the left, transform according to

$$f(ng) = \theta(n)f(g), \qquad n \in N'_F,$$

and are locally constant, and compactly supported modulo N'_F. In fact, the map $W \to W|G'_F$ is a bijection of $\mathcal{W}(\pi, \psi)$ onto $C_c^\infty(G', \theta)$. It follows that the integrals are convergent for all s and satisfy (2.2) with $L(s, \pi \times \pi') = L(s, \tilde{\pi} \times \tilde{\pi}') = 1$. In order to prove (2.3) one need only show that there is a constant c such that, for all W in $\mathcal{W}(\pi, \psi)$ and W' in $\mathcal{W}(\pi', \psi)$,

$$\int_{N'_F \backslash G'_F} \tilde{W}(g)\tilde{W}'(\varepsilon g)\,\mathrm{d}g = c \int_{N'_F \backslash G'_F} W(g)W'(\varepsilon g)\,\mathrm{d}g.$$

Indeed, the left hand side defines a bilinear form B on the product $C_c^\infty(G', \theta) \times W(\pi', \psi)$. It satisfies the following invariance condition:

$$B(\rho(g)f, \pi'(g)W') = B(f, W'), \qquad \text{where } (\rho(g)f)(h) = f(hg).$$

Therefore, it must of the form

$$B(f, W') = \int_{N'_F \backslash G'_F} \lambda(\pi'(g)W')f(g)\,\mathrm{d}g,$$

where λ is a linear form on $\mathcal{W}(\pi', \psi)$ such that

$$\lambda(\pi'(n)W') = \bar{\theta}(n)\lambda(W'), \qquad \text{for } n \in N'_F.$$

The uniqueness of the Whittaker model shows that λ must be proportional to the linear form $W' \to W'(\varepsilon)$ and our assertion follows.

In the next sections we call the factor $L(s, \pi \times \pi')$ the 'g.c.d.' of the integrals $\Psi(s, W, W')$ and set

$$\varepsilon'(s, \pi \times \pi', \psi) = \varepsilon(s, \pi \times \pi', \psi)L(1-s, \tilde{\pi} \times \tilde{\pi}')/L(s, \pi \times \pi').$$

We shall use also the same notation and terminology for other integrals which have similar properties.

3. The case $p = 2$

In this section we review the case $p = 2$. Accordingly, $G = GL(2)$, $G' = GL(1)$, $G'_F = F^\times$,

$$w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

$w' = 1$. Moreover, $w\,{}^t g^{-1} w^{-1} = \det g^{-1} \cdot g$ for g in G_F.

Let π and π' be as above. Let ω be the quasi-character of $F^\times$ such that

$$\pi\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} = \omega(a)\cdot 1.$$

Then $\tilde{\pi}$ is equivalent to the representation $\pi \otimes \omega^{-1}$ (i.e. the representation $g \to \pi(g)\omega^{-1}(\det g)$). Moreover, for W in $\mathcal{W}(\pi, \psi)$ the function $\tilde{W}$ is given by

$$\tilde{W}(g) = W(gw)\omega^{-1}(\det g).$$

On the other hand, the representation π' is just a quasi-character of $F^\times$ and the function W' coincides with the function π'. Therefore the integral $\Psi(s, W, W')$ reduces to the integral

$$\int_{F^\times} W\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} |a|^{s-\frac{1}{2}}\pi'(a)\mathrm{d}^\times a$$

and the integral $\Psi(s, \tilde{W}, \tilde{W}')$ to

$$\int_{F^\times} W\left[\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} w\right] |a|^{s-\frac{1}{2}}\omega^{-1}\pi'^{-1}(a)\mathrm{d}^\times a.$$

Hence, the results of [2] show that (2.1) to (2.3) are true with

$$L(s, \pi\times\pi') = L(s, \pi\otimes\pi'), \qquad \varepsilon(s, \pi\times\pi', \psi) = \varepsilon(s, \pi\otimes\pi', \psi).$$

We shall need the fact that these factors are related to other integrals as well. Indeed, let V be the space of π, $\tilde{V}$ the space of $\tilde{\pi}$, and $\langle\cdot,\cdot\rangle$ the invariant bilinear form on $V\times\tilde{V}$. A coefficient of π is a function f of the form

$$f(g) = \langle\pi(g)v, \tilde{v}\rangle$$

where v is in V and $\tilde{v}$ in $\tilde{V}$. The function $\check{f}$ defined by $\check{f}(g) = f(g^{-1})$ is a coefficient of $\tilde{\pi}$.

Let f be a coefficient of π, Φ a function in $\mathcal{S}(2\times 2, F)$ and s a complex number; we set

$$Z(\Phi, s, f) = \int_{G_F} f(x)|\det x|^s\Phi(x)\mathrm{d}^\times x,$$

where $\mathrm{d}^\times x$ is a multiplicative Haar measure on G_F. These integrals converge for Res large, the 'g.c.d.' of the integrals $Z(\Phi, s+\frac{1}{2}, f)$ is the factor $L(s, \pi) = L(s, \pi\times 1)$, and the integrals satisfy the functional equation

$$Z(\hat{\Phi}, \tfrac{3}{2}-s, \check{f}) = \varepsilon'(s, \pi, \psi)Z(\Phi, s+\tfrac{1}{2}, f) \tag{3.1}$$

where $\hat{\Phi}$ denotes the Fourier transform of Φ:

$$\hat{\Phi}(x) = \int_{M(2,F)} \Phi(y)\psi(Tr(yx))\mathrm{d}y,$$

Tr denoting the trace of an element of $M(2\times 2, F)$ and $\mathrm{d}y$ being the self-dual Haar measure on $M(2\times 2, F)$.

At least when π is supercuspidal, this can be derived from the following lemma combined with the fact that on $V\times\tilde{V} = \mathcal{W}(\pi,\psi)\times\mathcal{W}(\tilde{\pi},\psi)$ the invariant bilinear form is given by

$$\langle W_1, W_2\rangle = \int_{F^\times} W_1\begin{pmatrix} a & 0\\ 0 & 1\end{pmatrix} W_2\begin{pmatrix} -a & 0\\ 0 & 1\end{pmatrix}\mathrm{d}^\times a.$$

LEMMA (3.2): *Suppose Φ is in $\mathcal{S}(2\times 2, F)$. Then*

$$\int_{F^3}\Phi\left[\begin{pmatrix}1 & x\\ 0 & 1\end{pmatrix}\begin{pmatrix} a & 0\\ b & -1\end{pmatrix}\right]\psi(-b)\psi(-x)\mathrm{d}a\,\mathrm{d}b\,\mathrm{d}x$$

$$= \int_F \hat{\Phi}\left[w^{-1}\begin{pmatrix}1 & x\\ 0 & 1\end{pmatrix}\right]\psi(x)\mathrm{d}x.$$

The lemma itself is a simple consequence of the Fourier theorem in one variable.

Finally, we shall need the results of [3] that we now state in a form appropriate to our purposes. Let π_1 and π_2 be non-degenerate representations of $G_F = GL(2, F)$. For $\Phi\in\mathcal{S}(1\times 2, F)$, $W_1\in\mathcal{W}(\pi_1,\psi)$, and $W_2\in\mathcal{W}(\pi_2,\psi)$, we set

$$\Psi(s, W_1, W_2, \phi) = \int_{N_F\backslash G_F} W_1(g)W_2(\varepsilon g)\phi[(0,1)g]|\det g|^s\mathrm{d}g$$

where

$$\varepsilon = \begin{pmatrix}-1 & 0\\ 0 & 1\end{pmatrix}.$$

Then again, the integrals are convergent for Res large, have a g.c.d. $L(s, \pi\times\pi')$, and satisfy a functional equation

(3.3) $\Psi(1-s, \tilde{W}_1, \tilde{W}_2, \hat{\phi}) = \omega_2(-1)\varepsilon'(s, \pi_1\times\pi_2, \psi)\Psi(s, W_1, W_2, \phi)$

where $\hat{\phi}$ is the Fourier transform of ϕ:

$$\hat{\phi}(x, y) = \int_{F^2}\phi(u, v)\psi(ux+vy)\mathrm{d}u\,\mathrm{d}v,$$

and ω_i, for $i = 1, 2$, is the quasi-character of $F^\times$ defined by

$$\pi_i\begin{pmatrix} a & 0\\ 0 & a\end{pmatrix} = \omega_i(a)\cdot 1.$$

4. The case $p = 3$

In this section $p = 3$; accordingly, $G = GL(3)$, $G' = GL(2)$,

$$w = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \qquad w' = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \qquad \varepsilon = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

In addition, we let U be the subgroup of matrices of the form

$$\begin{pmatrix} 1 & 0 & u \\ 0 & 1 & v \\ 0 & 0 & 1 \end{pmatrix}.$$

Fix an admissible irreducible representation σ of G'_F and a quasi-character μ of $F^\times$. Let $\mathscr{H}$ be the space of σ and denote by $\mathscr{V}$ the space of all locally constant mappings f from G_F to $\mathscr{H}$, such that

$$f\left[\begin{pmatrix} m & 0 \\ u & a \end{pmatrix} g\right] = \sigma(m)|\det m|^{-\frac{1}{2}}\mu\alpha(a)f(g)$$

for all m in $G'_F = GL(2, F)$, a in $F^\times$, and g in G_F. We call π_0 the representation of G_F in $\mathscr{V}$ defined by right translations. From the results of [4], one can prove that any irreducible admissible representation of G_F which is not supercuspidal is a component of π_0, for a suitable choice of σ and μ. Furthermore, according to a result of Rodier, if π_0 has a non-degenerate component, then σ itself is non-degenerate. So, from now on, we assume σ to be so. We also let $\mathscr{H}$ be the space $\mathscr{W}(\sigma, \psi)$. Then, one can show that π_0 has exactly one non-degenerate component, π say.

In order to show that (2.1) to (2.3) are true for any non-degenerate representation of G_F, it suffices to show that they are true for such a π. In these notes, we assume in addition that π_0 is irreducible and sketch a proof of the fact that $\pi = \pi_0$ is then non-degenerate and that the conjectures are true for π and $\tilde{\pi}$.

Let $\mathscr{S} = \mathscr{S}(3 \times 2, F) \otimes \mathscr{H}$ be the space of all locally constant compactly supported maps from $M(3 \times 2, F)$ *to* $\mathscr{H}$. If Φ is in $\mathscr{S}$, then the function f defined by

$$f(g) = \mu\alpha(\det g) \int_{G'_F} \mu\alpha^{\frac{3}{2}}(\det m)\sigma(m^{-1})\Phi(vmg)\mathrm{d}^\times m, \tag{4.1}$$

where

$$v = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix},$$

is an element of $\mathscr{V}$ – provided the integral converges. For $g \in G_F$, $f(g)$ is

a function on G'_F whose value at $h \in G'_F$ is denoted by $f(g; h)$. A similar notation is used for Φ. Accordingly, $f(g; e)$ is simply

$$f(g; e) = \mu\alpha(\det g) \int_{G'_F} \mu\alpha^{\frac{3}{2}}(\det m)\Phi(vmg; m^{-1})\mathrm{d}^{\times}m.$$

Set now

$$W(g) = \int_{U_F} f(ug; e)\overline{\theta}(u)\mathrm{d}u. \tag{4.2}$$

Then, when W is defined, it satisfies

$$W(ng) = \theta(n)W(g), \qquad \text{for } n \in N_F.$$

Suppose now that Φ has the form

$$\Phi\left[\begin{pmatrix} a & b & x \\ c & d & y \end{pmatrix}; g\right] = \phi_1\begin{pmatrix} a & b \\ c & d \end{pmatrix}\phi_2(x, y)W_1(g), \tag{4.3}$$

where ϕ_1 is in $\mathscr{S}(2\times 2, F)$, ϕ_2 in $\mathscr{S}(2\times 1, F)$, and W_1 in $\mathscr{W}(\sigma, \psi)$. Then, for $g \in G'_F$,

$$\begin{aligned} W(g) &= \int_{U_F} f(gg^{-1}ug; e)\overline{\theta}(u)\mathrm{d}u \\ &= \alpha^{\frac{1}{2}}(\det g)\int_{U_F} f(u; g)\overline{\theta}(gug^{-1})\mathrm{d}u \\ &= \alpha^{\frac{1}{2}}(\det g)\int_{G'_F\cdot F^2} \phi_1(m)\phi_2[(x, y)^t m]W_1(gm^{-1})\psi[(0, -1)g^t(x, y)] \\ &\qquad \times u\alpha^{\frac{3}{2}}(\det m)d^{\times}m\mathrm{d}x\mathrm{d}y. \end{aligned}$$

Let ϕ be the co-Fourier-transform of ϕ_2:

$$\phi(u, v) = \int_{F^2} \phi_2(x, y)\psi[-(u, v)^t(x, y)]\mathrm{d}x\mathrm{d}y.$$

Then, we may write the above integral as

$$W(g) = \alpha^{\frac{1}{2}}(\det g)\int_{G'_F} \phi_1(m)\phi[(0, 1)gm^{-1}]W_1(gm^{-1})\mu\alpha^{\frac{1}{2}}(\det m)\mathrm{d}^{\times}m, \tag{4.4}$$

an integral which converges 'better' than (4.2). It can be used to define W even though (4.2) might not converge. Then, it can be shown that π is non-degenerate and W belongs to $\mathscr{W}(\pi, \psi)$. Actually, since the space $\mathscr{S}$ is spanned by functions of the form (4.3), the space $\mathscr{W}(\pi, \psi)$ is spanned by functions W for which $W(g)$, $g \in G'_F$, is given by (4.4).

We also need an integral representation for $\tilde{W}(g)$, $g \in G'_F$. Now, if g is in G'_F,

$$\tilde{W}(g) = W(w\,{}^t g^{-1}) = \int_{U_F} f(uw\,{}^t g^{-1}; e)\bar{\theta}(u)\,\mathrm{d}u$$

$$= \mu\alpha(\det g^{-1}) \int_{G'_F \cdot U_F} \mu\alpha^{\frac{3}{2}}(\det m)\Phi[vmuw\,{}^t g^{-1}; m^{-1}]\bar{\theta}(u)\,\mathrm{d}^\times m\,\mathrm{d}u.$$

More explicitly,

$$u = \begin{pmatrix} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}, \qquad \theta(u) = \psi(b), \qquad \mathrm{d}u = \mathrm{d}a\,\mathrm{d}b,$$

and $vmuw\,{}^t g^{-1}$ is the 3 by 2 matrix obtained by the juxtaposition of the 2 by 2 matrix

$$m \begin{pmatrix} a & 0 \\ b & -1 \end{pmatrix} {}^t g^{-1}$$

and the 1 by 2 matrix $m\binom{1}{0}$. Hence, the above integral may be written in the form

$$\mu\alpha(\det g^{-1}) \int_{G'_F \cdot F^2} \mu\alpha^{\frac{3}{2}}(\det m)\phi_1 \left[m \begin{pmatrix} a & 0 \\ b & -1 \end{pmatrix} {}^t g^{-1} \right] \phi_2[(1,0)\,{}^t m] \times W_1(m^{-1})\psi(-b)\,\mathrm{d}a\,\mathrm{d}b\,\mathrm{d}^\times m.$$

The integral on G'_F can be realized as an integral on N'_F and then on G'_F/N'_F. In this way, one obtains

$$\mu\alpha(\det g^{-1}) \int_{G'_F/N'_F} \mu\alpha^{\frac{3}{2}}(\det m)\phi_2[(1,0)\,{}^t m]W_1(m^{-1})\,\mathrm{d}^\times m \times \int \phi_1 \left[mn \begin{pmatrix} a & 0 \\ b & -1 \end{pmatrix} {}^t g^{-1} \right] \psi(-b)\,\mathrm{d}a\,\mathrm{d}b\bar{\theta}(n)\,\mathrm{d}n.$$

Using Lemma (3.2) to transform the inner integral, one arrives at

$$\alpha\mu^{-1}(\det g) \int_{G'_F/N'_F} \mu\alpha^{-\frac{1}{2}}(\det m)\phi_2[(1,0)\,{}^t m]W_1(m^{-1})\,\mathrm{d}^\times m \times \int_{N'_F} \hat{\phi}_1[{}^t g w'^{-1} n m^{-1}]\theta(n)\,\mathrm{d}n.$$

At this point, one puts back together the integrations on G'_F/N'_F and N'_F and changes m into ${}^t m^{-1}\,{}^t g w'^{-1}$ to arrive at the following formula:

$$\tilde{W}(g) = \alpha^{\frac{1}{2}}(\det g) \int_{G'_F} \mu^{-1}\alpha^{\frac{1}{2}}(\det m)\hat{\phi}_1\,{}^t(m)\hat{\phi}[(0,1)gm^{-1}] \times \tilde{W}_1(gm^{-1})\,\mathrm{d}^\times m. \tag{4.5}$$

Here $\hat{\phi}_1{}^t(m) = \hat{\phi}_1({}^t m)$ and $\hat{\phi} = \phi_2$ is the Fourier transform of ϕ.

Now let π' be a non-degenerate representation of G_F'. For W' in $\mathscr{W}(\pi', \psi)$, the integral $\Psi(s, W, W')$ may be written as a double integral with respect to $g \in N_F' \backslash G_F'$ and $m \in G_F'$. After changing g into gm we find that

$$\Psi(s, W, W') = \int_{G_F'} \phi_1(m) \mu \alpha^{s+\frac{1}{2}}(\det m) f(s, m) \mathrm{d}^\times m \tag{4.6}$$

where

$$f(s, m) = \Psi(s, W_1, \pi'(m)W', \phi).$$

Since, for a fixed s, the function $m \to f(s, m)$ is a coefficient of π', it is not too hard to see that (2.1) and (2.2) are true with

$$L(s, \pi \times \pi') = L(s, \pi' \times \sigma) L(s, \pi' \otimes \mu).$$

Similarly, one finds that

$$\Psi(s, \tilde{W}, \tilde{W}') = \int_{G_F'} \hat{\phi}_1(m) \mu^{-1} \alpha^{s+\frac{1}{2}}(m) f'(s, {}^t m) \mathrm{d}^\times m \tag{4.7}$$

where

$$f'(s, m) = \Psi(s, \tilde{W}_1, \tilde{\pi}'(m)\tilde{W}', \hat{\phi}).$$

Now one observes that

$$\tilde{\pi}'(m)\tilde{W}' = (\pi'({}^t m^{-1})W')^{\sim}$$

and therefore that, by (3.3),

$$f'(1-s, {}^t m) = \omega'(-1)\varepsilon'(s, \pi \times \pi', \psi) f(s, m^{-1}).$$

Combining this with (3.1), one arrives at the functional equation (2.3) with

$$\varepsilon(s, \pi \times \pi', \psi) = \varepsilon(s, \pi' \times \sigma, \psi)\varepsilon(s, \pi' \otimes \mu, \psi).$$

Hence (2.1) to (2.3) are now completely proved for $p = 3$. For instance, if μ, π, and π' are 'unramified' – contain the trivial representation of a maximal compact subgroup, the various factors have the following form:

$$L(s, \mu) = (1 - cq^{-s})^{-1}, \qquad L(s, \sigma) = (1 - aq^{-s})^{-1}(1 - bq^{-s})^{-1},$$

$$L(s, \pi') = (1 - a'q^{-s})^{-1}(1 - b'q^{-s})^{-1}, \tag{4.5}$$

$$L(s, \pi \times \pi')^{-1} = (1 - a'cq^{-s})(1 - b'cq^{-s})(1 - aa'q^{-s})(1 - ba'q^{-s})(1 - ab'q^{-s}) \times (1 - bb'q^{-s}).$$

If, in addition, the character ψ is of order zero, then the ε factor is one.

5. Conclusion

Let us go back to the situation and the notations of Section 1. Let π be an admissible irreducible representation of $G_{\mathbb{A}}$. Then, π is, in a certain sense, an infinite tensor product $\otimes \pi_v$, where, for each place v, π_v is an admissible irreducible representation of $G_v = GL(3, F_v)$; almost all the representations π_v are unramified. We shall assume that π is non-degenerate (i.e. that each π_v is non-degenerate) and also that there is a character ω of $\mathbb{I}/F^\times$ such that

$$\pi\begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix} = \omega(a) \cdot 1 \qquad \text{for } a \text{ in } \mathbb{I}.$$

Let also π' be a representation of $G'_{\mathbb{A}}$ satisfying the same assumptions. Then, we define

$$L(s, \pi \times \pi') = \prod_v L(s, \pi_v \times \pi'_v), \qquad \varepsilon(s, \pi \times \pi') = \prod_v \varepsilon(s, \pi_v \times \pi'_v, \psi_v).$$

The first product is an infinite Euler product whose factors are almost all of the form (4.5); if, for instance, the representations are pre-unitary, then it converges for Res sufficiently large. In the second factor, we have set $\psi(x) = \prod \psi_v(x_v)$; the product has only a finite number of terms $\neq 1$ and is actually independent of the choice of ψ.

Suppose now that π and π' are contained in the corresponding space of cusp-forms. They are then both pre-unitary and non-degenerate. The factor $L(s, \pi \times \pi')$ is a polynomial in Q^{-s}, Q^s, and satisfies the functional equation

$$L(s, \pi \times \pi') = \varepsilon(s, \pi \times \pi') L(1-s, \tilde{\pi} \times \tilde{\pi}').$$

The proof follows step by step the proof of (11.1) in [3].

REFERENCES

[1] I. M. GELFAND and D. A. KAJDAN: Representations of $GL(n, k)$ where k is a local field. *Institute for Applied Mathematics, No. 942*, 1971.

[2] H. JACQUET and R. P. LANGLANDS: Automorphic forms on $GL(2)$. *Lecture Notes in Math., No. 114*. Springer-Verlag, Berlin and New York, 1970.

[3] H. JACQUET: Automorphic Forms on $GL(2)$, II. *Lecture Notes in Math., No. 278*. Springer-Verlag, Berlin and New York, 1972.

[4] H. JACQUET and R. GODEMENT: Zeta Functions of Simple Algebras. *Lecture Notes in Math., No. 260*. Springer-Verlag, Berlin and New York, 1972.

[5] F. RODIER: Whittaker models for admissible representations of reductive p-adic split groups. *Harmonic Analysis on Homogeneous Spaces, Proc. of Symp. in Pure Math.*, Vol. XXVI, A.M.S., Providence, R.I., 1973.

[6] J. A. SHALIKA: The multiplicity one theorem for GL_n. *Ann. of Math.*, Vol. 100, No. 1, 1974.

(Oblatum 16–I–1974 & 11–III–1974)

Columbia University
in the city of New York
The Johns Hopkins University