Groupe d'étude de théories stables

Anand Pillay

The models of a non-multidimensional ω-stable theory

Groupe d'étude de théories stables, tome 3 (1980-1982), exp. n° 10, p. 1-22

<http://www.numdam.org/item?id=STS_1980-1982__3__A10_0>

© Groupe d'étude de théories stables
(Secrétariat mathématique, Paris), 1980-1982, tous droits réservés.

L'accès aux archives de la collection « Groupe d'étude de théories stables » implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

Numdam
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

Théories Stables
(B. POIZAT)
Vol. 3, 1980-1982, n° 10, 22 p.

THE MODELS OF A NON-MULTIDIMENSIONAL ω-STABLE THEORY

by Anand PILLAY (*)
[University of Manchester]

I give a (self-contained) account of the classification of the models of a non-multidimensional ω-stable theory. This result is the generalisation of the Baldwin-Lachlan-Morley classification of the models of an $\aleph_1$-categorical theory, and includes of course the possible spectra that can occur. (Remember that the spectrum of a theory T is given by the function $I(-, T)$, where for κ a cardinal, $I(\kappa, T)$ is the number of models of T of power κ, up to isomorphism.) The crude idea is that, instead of a model of T being determined by the cardinalfty of one indiscernible set (as when T is $\aleph_1$-categorical), a model of T is now determined by the cardinalities of each member of a fixed "independent" family of indiscernible sets.

I assume the basic facts about stability, forking, definability, rank, etc., which can be found in [4] or even [5].

T will be a countable complete ω-stable theory. The ω-stability of T furnishes us with several nice properties. The most important of these will be :

(i) for any subset A of a model M of T, there is a (real) prime model of $Th(M, a)$, $a \in A$,

(ii) if $M \models T$ and $p \in S(M)$, then there is a finite $A \subset M$ such that p is definable over A (thus p does not fork over A and $p \restriction A$ is stationary),

(iii) all types over arbitrary subsets are ranked by Morley rank.

I will also follow the usual practice of working in a large sufficiently saturated model of T.

I. Strongly regular types.

Strongly regular types are generalisations of types of Morley rank 1, degree 1. If $p \in S(M)$, I denote by $M(p)$, the model which is prime over $M \cup \{\bar{a}\}$, where $tp(\bar{a}/M) = p$. This model might also be denoted by $M(\bar{a})$, and is unique up to M-isomorphism.

Definition 1.1. - Let $p \in S_1(M)$, p not algebraic and $\varphi(x) \in p$ (φ might countain parameters from M). The pair (p, φ) is said to be strongly regular if whenever $b \in M(p)$, $b \notin M$ and $M(p) \models \varphi(b)$, then $tp(b/M) = p$. p is said to be

---

(*) Anand PILLAY, Dept of Mathematics, The University, MANCHESTER, M13 9PL (Grande-Bretagne).

strongly regular if there is $\varphi \in p$ such that (p , φ) is strongly regular.

LEMMA 1.2. - Suppose that p and $q \in S(M)$, p is strongly regular and q is realised in $M(p)$. Then p is realised in $M(q)$, (We assume q is not algebraic).

Proof. - Suppose that (p , φ) is strongly regular. Let a realise p, and $\bar{b} \in M(a)$ such that $\bar{b}$ realises q. It is clear that a and $\bar{b}$ are note independant over M $(\bar{b} \notin M)$. Thus there is a formula $\alpha(x , \bar{y})$ over M such that $M(a) \models \alpha(a , \bar{b})$, but $M(a) \models \neg \alpha(m , \bar{b})$ for all $m \in M$. Note that " $\varphi(x) \wedge \alpha(x , \bar{b})$ " is consistent. Now $M(q) = M(\bar{b}) \prec M(a)$, and let $c \in M(\bar{b})$ such that $M(b) \models \varphi(c) \wedge \alpha(c , \bar{b})$. Then $c \notin M$, $c \in M(a)$ and $M(a) \models \varphi(c)$. Thus $tp(c/M) = p$, and so p is realised in $M(q)$.

Definition 1.3. - Let p and q be strongly regular types over M such that q is realised in $M(p)$. Then we say that p and q are equivalent, $p \sim q$.

(By lemma 1.2, this definition makes sense).

The next lemma shows that "enough" strongly regular types exist.

LEMMA 1.4. - Suppose that $M \prec N$, the $L(M)$ formula $\varphi(x)$ is "augmented" in N, and a is chosen in $\varphi^N - M$ such that $tp(a/M)$ has least possible Morley rank. Then $tp(a/M)$ is strongly regular.

Proof. - (Let $R(-)$ denote Morley rank). Let $R(tp(a/M)) = \alpha$, and pick $L(M)$-formula $\psi(x)$ such that $\vdash \psi(x) \longrightarrow \varphi(x)$, $N \models \psi(a)$ and $R(\psi(x)) = \alpha$, and degree $(\psi(x)) = 1$. Now $M(a) \prec N$, and so it is clear that $(tp(a/M) , \psi)$ is strongly regular.

Definition 1.5. - Let $p(\bar{x})$ and $q(\bar{y})$ be types over M. p and q are said to be perpendicular $(p \perp q)$ if $p(\bar{x}) \cup q(\bar{y})$ determines a complete $\bar{x} \hat{} \bar{y}$ type over M.

Note. - If $p(\bar{x})$ and $q(\bar{y})$ are types over a model M, then $p \perp q$ if, and only if, whenever $\bar{a}$ and $\bar{b}$ realise p and q respectively, then $\bar{a}$ and $\bar{b}$ are independent over M.

Fact 1.6. - Let $\bar{a}$ and $\bar{b}$ be independent over M. Let A be a atomic over $M \cup \{\bar{a}\}$ and B atomic over $M \cup \{\bar{b}\}$. Then A and B are independent over M.

Now the proof of lemma 1.2 actually implies that if $tp(a/M)$ is strongly regular and $\bar{b} \in M(a) - M$, then $tp(a/M \cup \{\bar{b}\})$ is isolated. A simple consequence of this and fact 1.6 is the following :

Observation 1.7. - Let p_1, p_2, q_1, q_2 be all strongly regular types over M such that $p_1 \sim p_2$ and $q_1 \sim q_2$. Then $p_1 \perp q_1$ if, and only if, $p_2 \perp q_2$.

PROPOSITION 1.8. - Let p and q be strongly regular types over M. Then p and q are perpendicular if, and only if, p and q are not equivalent.

Proof. - It is clear that if p and q are equivalent then they are not perpendicular. Conversely, assume that p and q are not equivalent. We wish to show that they are perpendicular. By 1.7, we can assume that $R(p) = \alpha$ is minimal among strongly regular types over M equivalent to p, and similarly for q, with $R(q) = \beta$. So we can find formulae $\varphi(x)$ and $\psi(x)$, both of degree 1, and of rank α and β respectively, such that (p , φ) and (q , ψ) are strongly regular. Suppose (without loss) that $\alpha \leqslant \beta$. Now if p and q are not perpendicular, then there are realisations a and b of p and q respectively, such that a and b are not independent over M. As in the proof of 1.2, if follows that $\varphi(x)$ is "augmented" in $M(b)$ (i. e. $\varphi^{M(b)} - M$ is nonempty). By lemma 1.4, there is $c \in \varphi^{M(b)} - M$, such that $tp(c/M)^*$ is strongly regular. Clearly, $tp(c/M)$ is equivalent to q, and $R(tp(c/M)) \leqslant \alpha$. If $R(tp(c/M)) = \alpha$, then clearly $tp(c/M) = p$, which contradicts the non-equivalence of p and q. On the other hand, if $R(tp(c/M)) < \alpha$, then we contradict the minimal choice of $R(q)$. Thus the proposition is proved.

PROPOSITION 1.9. - Let $M \prec M'$, $p \in S_1(M)$ and p' the nonforking extension (or heir) of p over M'. Then p is strongly regular if, and only if, p' is strongly regular.

Proof. - First suppose that p' is strongly regular. Then there is an $L(M)$ formula $\varphi(x)$ such that (p' , φ) is strongly regular (Any $L(M')$ formula $\varphi(x) \in p'$ such that degree $(\varphi) = 1$, and $R(\varphi) = R(p')$ will suffice. But $p \subset p'$, and $R(p) = R(p')$. Thus φ can be chosen over M). We show that (p , φ) is strongly regular. Let a realise p'. So $tp(a/M) = p$, and $M(p) = M(a) \prec M'(a)$. Let $b \in M(a)$, $b \notin M$ and b satisfy φ. Now b and a are not independent over M. Thus $b \notin M'$. But then $tp(b/M') = p'$ (by strong regularity of (p' , φ) .) Thus $tp(b/M) = p$. So (p , φ) is strongly regular.

Conversely, suppose that $\varphi(x) \in p$, and (p , φ) is strongly regular. Let a realise p'. If (p' , φ) is not strongly regular, then there is b in $\varphi^{M'(a)} - M'$ such that $tp(b/M') \neq p$. Now p' is definable by a schema d, over M (where d also defines p), and also a and b are not independent over M'. Thus there are L-formulae $\psi(y , \bar{z})$ and $\alpha(x , y , \bar{w})$, and $\bar{c}$ and $\bar{d}$ in M' such that

$$M'(a) \models (\exists\, y)(\varphi(y) \wedge \psi(y , \bar{c}) \wedge \neg\, d(\psi)(\bar{c}) \wedge \alpha(a , y , \bar{d})) ,$$

where the formula $\alpha(x , y , \bar{w})$ is not represented in p' (so neither in p). But $tp(a/M')$ is the heir of $tp(a/M)$. Thus we can find $\bar{c}'$ and $\bar{d}'$ in M such that

$$M(a) \models (\exists\, y)(\varphi(y) \wedge \psi(y , \bar{c}') \wedge \neg\, d(\psi)(\bar{c}') \wedge \alpha(a , y , \bar{d}')) .$$

If we let b' be such a y in $M(a)$, then $b' \notin M$, $M(a) \models \varphi(b')$ and $tp(b'/M) \neq p$. This contradicts the fact that (p , φ) is strongly regular, and completes the proof.

Definition 1.10. - Let A be a subset (of the big model), $p \in S_1(A)$ a stationary type, and $\varphi(x) \in p$. We call (p , φ) strongly regular if there is a model M containing A and nonforking extension p' of p over M such that (p',φ) is strongly regular. Again p will be called strongly regular if there is $\varphi(x)$ such that (p , φ) is strongly regular.

Note. - If follows immediately from 1.9 that for $p \in S_1(A)$, p is strongly regular if, and only if, for all M extending A and nonforking extension p' of p to M, p' is strongly regular.

PROPOSITION 1.11. - Let p and q be strongly regular types over M, and let p' and q' be their respective heirs over $M' \prec M$. Then $p \perp q$ if, and only if, $p' \perp q'$.

Proof. - Suppose that p dans q are not perpendicular. Then there are realisations a and b of p and q respectively, such that a and b are not independent over M. Let $a' \hat{} b'$ realise the heir of $tp(a \hat{} b/M)$ over M'. Then $tp(a'/M') = p'$, $tp(b'/M') = q'$, and a' and b' are not independent over M'. Thus $p' \not\perp q'$.

Conversely, suppose that p and q are perpendicular. We may again suppose that p and q are chosen with minimal rank in their equivalence classes. So we have (p , φ) strongly regular, with $R(p) = R(\varphi) = \alpha$, and (q , ψ) strongly regular, with $R(q) = R(\psi) = \beta$, and suppose $\alpha \leqslant \beta$. So (p' , φ) and (q' , ψ) are strongly regular. If p' and q' are not perpendicular, then again if follows that $\varphi(x)$ is augmented in $M'(q')$. As q' is the heir of q, it is easy to prove that $\varphi(x)$ is augmented in $M(q)$, but this will again contradict the minimal choice of $R(q)$. So the proposition is proved.

By propositions 1.8 and 1.11, we have :

COROLLARY 1.12. - Let p and q be strongly regular types over M, and $M \prec M'$, and p', q' the heirs of p and q over M'. Then $p \sim q$ if, and only if, $p' \sim q'$.

Definition 1.13. - Let $p(\bar{x})$ and $q(\bar{y})$ be in $S(A)$, where A is an arbitrary subset. Then p and q are said to be orthogonal if for all $B \supset A$ and nonforking extensions p' and q' of p and q over B, $p'(\bar{x}) \cup q'(\bar{y})$ determines a complete type over B.

PROPOSITION 1.14. - Let p and q be strongly regular types over A. Then the following are equivalent :

(i) p and q are orthogonal,

(ii) for some $M \supset A$, (M a model), and nonforking extensions p', q' of p, q over M, p' and q' are perpendicular.

Proof. - By proposition 1.11.

Note. - It was shown in [3] that if p and q are any types over M, and p', q' their heirs over some $M' \succ M$, then $p \perp q$ if, and only if, $p' \perp q'$. It follows that proposition 1.14 holds without the hypothesis that p and q are strongly regular. However 1.14 in its present form will suffice for our needs.

Given strongly regular types p and q over A, we will call p and q equivalent if they are not orthogonal. By 1.8 and 1.14, this is consistent with def. 1.3.

I complete this section with a couple of observations which will be of use later on.

LEMMA 1.15. - Let $\{p_i ; i \in I\}$ be a set of stationary pairwise orthogonal types over A. For each $i \in I$, let $\{\bar{a}^i_j ; j < \gamma_i\}$ be an independent set of realisations of p_i over A. Then $\{\bar{a}^i_j ; i \in I, j < \gamma_i\}$ is independent over A.

Proof. - It suffices to show that if $\bar{a}_1, \bar{a}_2, \ldots, \bar{a}_n$ is independent over A, and $tp(\bar{b}/A)$ and $tp(\bar{a}_i/A)$ are orthogonal for $i = 1, \ldots, n$, then $\{\bar{a}_1, \bar{a}_2, \ldots, \bar{a}_n, \bar{b}\}$ is independent over A. This we show by induction. So suppose that we already have $\{\bar{a}_1, \ldots, \bar{a}_r, \bar{b}\}$ is independent over A, where $r < n$. Thus $tp(\bar{b}/\{\bar{a}_1, \ldots, \bar{a}_r\} \cup A)$ does not fork over A, and we know anyway that $tp(\bar{a}_{r+1}/\{\bar{a}_1, \ldots, \bar{a}_r\} \cup A)$ does not fork over A. Thus by the orthogonality of $tp(\bar{b}/A)$ and $tp(\bar{a}_{r+1}/A)$, $\bar{a}_{r+1}$ and $\bar{b}$ are independent over $A \cup \{a_1, \ldots, a_r\}$. Thus $\{\bar{a}_1, \ldots, \bar{a}_r, \bar{a}_{r+1}, \bar{b}\}$ is independent over A.

LEMMA 1.16. - Let M be a model, and $\{p_i ; i \in I\}$ a maximal collection of pairwise orthogonal strongly regular types over M. Let $A \subset M$ be such that each p_i is definable over A, and for each $i \in I$, let $\{a^i_j ; j < \gamma_i\}$ be a maximal independent set of realisations of $p_i \restriction A$ in M. Then $J = \{a^i_j ; i \in I, j < \gamma_i\}$ is independent over A, and moreover M is minimal over $A \cup J$.

Proof. - By 1.14, the types $p_i \restriction A$ are strongly regular and pairwise orthogonal. Thus the independence of J over A follows by 1.15.

Suppose that M were not minimal over $A \cup J$. Then there would be a model N such that $A \cup J \subset N \not\supseteq M$. By 1.4, we can find $a \in M - N$ such that $tp(a/N)$ is strongly regular. Let $p = tp(a/N)$, and let p' be the heir of p over M. So p' is strongly regular (1.9), and by the choice of the p'_i s there is $s \in I$ such that p' and p_s are not orthogonal. But p_s does not fork over N, and so $p_s \restriction N$ is strongly regular and not orthogonal to p (by prop. 1.9 and prop. 1.11). Thus $p_s \restriction N$ and p are equivalent, and so $p_s \restriction N$ is realised in $N(a)$, where we can assume that $N(a) \prec M$. Let $c \in N(a)$ realise $p_s \restriction N$. Then, as $p_s \restriction N$ does not fork over A, if follows that c and $\{a^s_j ; j < \gamma_s\}$ are independent over A. But this contradicts the maximal choice of the independent set $\{a^s_j ; j < \gamma_s\}$ of realisations of $p_s \restriction A$ in M. So the lemma is proved.

II. Dimension.

Let M be a model, $A \subset M$ and $p \in S(A)$. A set I of tuples from M will be called a basis for p in M, if I is a set of realisations of p in M, independent over A, and maximal such (Note that if p is stationary, then I is also indiscernible over A).

PROPOSITION II.1. - Suppose that $p \in S(A)$, $A \subset M$ and p has some infinite basis in M. Then all bases for p in M have the same cardinality.

Proof. - If not, then it is clear that there are bases I and J of p in M with J infinite and $|I| < |J|$. As I is maximal, for each $\bar{c} \in J$, $tp(\bar{c}/I \cup A)$ forks over A. So there is some finite $I_{\bar{c}} \subset I$ such that $tp(\bar{c}/I_{\bar{c}} \cup A)$ forks over A.

So by the cardinality difference, there is finite $I' \subset I$ and $\bar{c}_n \in J$ for $n<\omega$, such that $tp(\bar{c}_n/I' \cup A)$ forks over A, for each $n < \omega$. But then, as the c_n' s are independent over A, we have for each $n < \omega$, $tp(\bar{c}_{n+1}/\{\bar{c}_0,\ldots,\bar{c}_n\} \cup I' \cup A)$ forks over $A \cup \{\bar{c}_0, \ldots, \bar{c}_n\}$, and thus $tp(I'/\{\bar{c}_0, \ldots, \bar{c}_n, \bar{c}_{n+1}\} \cup A)$ forks over $\{\bar{c}_0, \ldots, \bar{c}_n\} \cup A$. But this contradicts superstability.

Definition II.2. - If all bases of p in M have the same cardinality, then we define $\dim(p, M)$ to be this cardinality.

Note. - We will see later on that if $p \in S(A)$ is strongly regular and $A \subset M$, then $\dim(p, M)$ is always defined.

Let I be an infinite indiscernible set (maybe of tuples), and B an arbitrary set. Recall that $Av(I/B)$ is defined as follows : for $\bar{b} \in B$, $\varphi(\bar{x}, \bar{b}) \in Av(I/B)$ if, for cofinitely many $\bar{c}$ in I, we have $\models \varphi(\bar{c}, \bar{b})$. Then $Av(I/B)$ is a complete and consistent type over B. Moreover, suppose that p is a stationary type over A, and I is an infinite independent set of realisations of p over A (so I is indiscernible over A), and $B \supset A$. Then $Av(I/B)$ is precisely p' the nonforking extension of p over B.

LEMMA II.3.

(i) Let I be an infinite indiscernible set over A, and M prime over $A \cup I$. Then I is a maximal indiscernible set over A in M.

(ii) Let $I \cup \{c\}$ be an infinite indiscernible set over A, and let M be prime over $A \cup I$. Then $tp(c/A \cup I) \vdash Av(I/M)$.

(iii) Let p be a stationary type over A, and I an independent set of realisations of p over A. Let I_1 be an infinite subset of I, and M be prime over $A \cup I_1$, and let p' denote the nonforking extension of p over M. Then $I - I_1$ is an independent (over M) set of realisations of p'.

Proof.

(i) If I is not maximal indiscernible over A in M, extend it by c in M. Now $tp(c/A \cup I)$ is isolated by a formula $\alpha(x, \bar{a}, \bar{d})$, where $\bar{a} \in A$ and $\bar{d} \subset I$. In particular, $M \models \alpha(x, \bar{a}, \bar{d}) \rightarrow x \neq c'$ for all $c' \in I$ (as $c \notin I$). But $M \models \alpha(c, \bar{a}, \bar{d})$, $I \cup \{c\}$ is indiscernible over A and I is infinite. Thus we can find c' in I such that $M \models \alpha(c', \bar{a}, \bar{d})$, and this is a contradiction.

(ii) I show that if $I \cup \{c\}$ is indiscernible over A, then $tp(c/M) = Av(I/M)$ (where M is prime over $A \cup I$). So let $\varphi(x, \bar{m}) \in Av(I/M)$, where $\bar{m} \in M$. I will show that this formula is satisfied by c. Now $tp(\bar{m}/A \cup I)$ is isolated by a formula $\varphi(\bar{y}, \bar{a}, \bar{d})$ where $\bar{a} \in A$ and $\bar{d} \subset I$. Now as $\varphi(x, \bar{m})$ is satisfied by cofinitely many members of I, there is $c' \in I$, $c' \notin d'$ such that $M \models \varphi(c', \bar{m})$. Thus $M \models \forall \bar{y}(\varphi(\bar{y}, \bar{a}, \bar{d}) \rightarrow \varphi(c', \bar{y}))$. But $tp(c \hat{} \bar{d}/\bar{a}) = tp(c \hat{} \bar{d}/\bar{a})$. So we have $\models \forall \bar{y}(\varphi(\bar{y}, \bar{a}, \bar{d}) \rightarrow \varphi(c, \bar{y}))$, whereby $\models \varphi(c, \bar{m})$, and we finish.

(iii) Let $c_1, \ldots, c_n$ be in $I - I_1$. We must show that $c_1, \ldots, c_n$ is an independent set of realisations of p' over M. Let p^n denote $tp(c_1 \hat{} \ldots \hat{} c_n/A)$. Then I can be considered (by partitioning it into n-tuples) as an independent set of realisations over A of p^n. But then by (ii) and the remarks preceding this lemma, $tp(c_1 \hat{} \ldots \hat{} c_n/M)$ does not fork over A, and this is just what we want.

LEMMA II.4.

(i) *Let p and q be equivalent strongly regular types over a model M, and let $N \succ M$. If I is a basis of p in N, then there is a basis J of c in N with $|I| \leqslant |J|$.*

(ii) *Let p and q be equivalent strongly regular types over a set A, and let $N \supset A$. If p has an infinite basis in N, then so does q, and moreover $\dim(p, M) = \dim(q, M)$.*

Proof.

(i) Let I be a basis of p in N, and write I as $\{a_\alpha ; \alpha < \kappa\}$. Define models M_α in N for $\alpha < \kappa$, and elements b_α for $\alpha < \kappa$, as follows: $M_0 = M$, $M_{\alpha+1} = M_\alpha(a_\alpha)$, and $M_\delta = \bigcup_{\alpha<\delta} M_\alpha$. Clearly $tp(a_\alpha/M_\alpha)$ is the heir of p over M_α and so strongly regular and equivalent to a_α, the heir of c over M_α (by 1.12). Thus c_α is realised in $M_{\alpha+1}$, and let b_α be such a realisation. By fact 1.6, $\{b_\alpha ; \alpha < \kappa\}$ is an independent set of realisations of q over M, and so can be extended to a maximal such set in N.

(ii) It is enough by II.1 and symmetry to show that if p has an infinite basis I in N, then q has a basis J in N with $|I| \leqslant |J|$. So let I be an infinite basis of p in N. Partition I as $I_1 \cup I_2$, where I_1 is infinite and $|I| = |I_2|$. Let N' be an elementary substructure of N which is prime over

$A \cup I_1$. Let p' be the nonforking extension of p over M' . Then by II.3 (iii), I_2 is a basis of p' in N . But p' is strongly regular and equivalent to q' , the (strongly regular) nonforking extension of q over M' . So by (i) there is a basis J' of q' in N , with $|J'| \geq |I_2| = |I|$. But J' is clearly an independent set of realisations of q in N and so can be extended to a basis J of q in N , and clearly $|I| \leq |J|$.

Let p be a stationary type over A , and let $\bar{a}_1$, $\bar{a}_2$, ... , $\bar{a}_n$ an independent set of realisations of p over A . Then I will denote $tp(\bar{a}_1 \hat{} \ldots \hat{} \bar{a}_n/A)$ by p^n .

PROPOSITION II.5. - Let p and q be strongly regular types over a set A , and suppose that, for all n , $m < \omega$, $p^n(\bar{x}) \cup q^m(\bar{y})$ determines a complete type over A . Then p and q are orthogonal.

Proof. - So suppose that, for all n , $m < \omega$, $p^n(\bar{x}) \cup q^m(\bar{y})$ is complete. It follows that if I is an independent set of realisations of p over A , and J is an independent set of realisations of q over A , then I and J are independent over A . Now pick I and J as in the last sentence, and such that both are infinite and $|I| < |J|$. Let M be prime over $A \cup J \cup I$. I assert that I is a basis for p in M . Note first that I is indiscernible over $A \cup J$. Now if c were a realisation of p in M such that $I \cup \{c\}$ were independent over A , then by our hypothesis, $I \cup \{c\}$ and J would be independent over A , and thus $I \cup \{c\}$ would be indiscernible over $A \cup J$, contradicting lemma II.3 (i). Thus I is a basis for p in M , and so $\dim(p , M) = |I|$. But clearly $\dim(q , M) \geq |J| > |I|$. So by lemma II.4, p and q are not equivalent, that is, p and q are orthogonal.

Note. - Proposition II.5 is actually true without the restriction that p and q be strongly regular (although we will not need this here). This fact, together with lemma 1.15 characterises orthogonality for types over sets.

LEMMA II.6. - Let $p \in S_1(A)$ and (p , φ) strongly regular. Suppose that $B \supset A$ and that p' and q are in $S_1(B)$, where p' is the nonforking extension of p over B , $q \neq p'$, and $\varphi \in q$. Then p' and q are orthogonal.

Proof. - It is enough to prove this in the case where B is a model, say M , and in this case it is enough to show that p' and q are perpendicular. So let a and b be realizations of p' and q respectively. I show that a and b are independent over M . Now as $q \neq p'$, there is some $L(M)$ formula $\psi(x)$ such that $\psi(x) \in q$ but $\neg \psi(x) \in p$. Suppose that $\alpha(x , y)$ is an $L(M)$-formula such that $\models \alpha(b , a)$. Thus $\models (\exists x)(\varphi(x) \wedge \alpha(x , a))$. So

$$M(a) \models (\exists x)(\varphi(x) \wedge \varphi(x) \wedge \alpha(x , a)) .$$

Let $c \in M(a)$ be such that $M(a) \models \varphi(c) \wedge \psi(c) \wedge \omega(c , a)$. So c satisfies $\varphi(x)$ but c does not realise p' . Thus $c \in M$ (by strong regularity of (p' , φ)). Thus we have shown that $tp(a/M \cup \{b\})$ is the heir of p' , whereby a and b are independent over M .

If follows from lemma II.6 that if p is strongly regular then p is regular ($p \in S(A)$ is said to be regular if whenever $B \supset A$, p' is the nonforking extension of p over B and q is a forking extension of p over B , then p' and q are orthogonal). Now for regular types the "nonforking" notion of independence on realisations of such types satisfies the familiar exchange principle. Namely : let $p \in S(A)$ be regular, $A \subset M$, and let $\bar{a}_i$, for $i < n$, and $\bar{b}$ realise p in M , where $\{\bar{a}_i ; i < n\}$ is a basis for p in M . Let $\bar{a}_m$ be the first element such that $tp(\bar{b}/\{\bar{a}_i ; i < m\})$ forks over M . Then $\{\bar{a}_0 , \dots , \bar{a}_{m-1} , \bar{b} , \bar{a}_{m+1} , \dots , \bar{a}_{n-1}\}$ is a basis for p in M . (This is a simple consequence of regularity and the basis properties of forking). Thus we have :

PROPOSITION II.7. - Let $p \in S_1(A)$ be strongly regular, and $A \subset M$. Then all bases for p in M have the same cardinality (and thus we can speak of $\dim(p , M)$)

PROPOSITION II.8. - Let p and q be equivalent strongly regular types over a model M , and let $N > M$. Then $\dim(p , N) = \dim(q , N)$.

Proof. - By lemma II.4 and proposition II.8.

I recall the following :

Fact II.9. - Let $p \in S(M)$ and $\varphi(\bar{x}) \in p$. Then p does not fork over $\cup \varphi^M$.

LEMMA II.10. - Let $p \in S_1(A)$, (p , φ) strongly regular, and $A \subset M \prec N$. Let p' denote the nonforking extension of p over M . Let I_1 be a basis for p in M , and let I_2 be an independent over M set of realisations of p' in N , and finally let $c \in N$ and $tp(c/I_1 \cup I_2 \cup A)$ is the nonforking extension of p over $I_1 \cup I_2 \cup A$. Then $tp(c/I_2 \cup M)$ does not fork over A (and thus $I_2 \cup \{c\}$ is an independent set of realisations of p' in N , over M .)

Proof. - It is enough to show that $tp(\{c\} \cup I_2/M)$ does not fork over A . By fact II.9, it is enough to show that $tp(\{c\} \cup I_2/\varphi^M \cup A)$ does not fork over A . Now, by hypothesis, $tp(I_2/\varphi^M \cup A)$ does not fork over A , and thus it suffices to prove that $tp(c/I_2 \cup \varphi^M \cup A)$ does not fork over $I_2 \cup A$. But $I_1 \subset \varphi^M$, and we know that $tp(c/I_2 \cup I_1 \cup A)$ does not fork over $I_2 \cup A$. So this leaves us having to prove that

$(*)$ $\qquad tp(c/I_2 \cup \varphi^M \cup A)$ does not fork over $I_2 \cup I_1 \cup A$.

Let $\bar{d} \subset \varphi^M$, and $d \in \varphi^M$, and suppose that we already know that $tp(c/I_2 \cup I_1 \cup \bar{d} \cup A)$ does not fork over $I_2 \cup I_1 \cup A$. Now it is clear that $tp(d/I_2 \cup I_1 \cup \bar{d} \cup A) \neq tp(c/I_2 \cup I_1 \cup \bar{d} \cup A)$ (either $tp(d/A) \neq p$, or d and I_1 are dependent over A). But d satisfies $\varphi(x)$. So by strong regularity of (p, φ) and lemma II.6, c and d are independent over $I_2 \cup I_1 \cup \bar{d} \cup A$. Thus $tp(c/I_2 \cup I_1 \cup \bar{d} \hat{} d \cup A)$ does not fork over $I_2 \cup I_1 \cup A$. So (*) is proved, and so also the lemma.

PROPOSITION II.11. - Let $p \in S(A)$ be strongly regular, $A \subset M \prec N$, and p' the nonforking extension of p over M. Then $\dim(p, N) = \dim(p, M) + \dim(p', N)$.

Proof. - By lemma II.10, if I_1 is a basis for p in M, and I_2 is a basis for p' in N, then $I_1 \cup I_2$ is a basis for p in N.

III. Non-multidimensional theories.

Definition III.1.

(i) Let M be a model of T. Then $\mu(M)$ denotes the maximum number of pairwise orthogonal strongly regular types over M.

(ii) T will be said to be multidimensional if for any λ there is a model M of T with $\mu(M) \geq \lambda$. Otherwise T is said to be non-multidimensional.

I now give some background on material to come. Firstly, if p_1 is a type over a finite set $\bar{a}$, then p_1 can be written in the form $p(\bar{x}, \bar{a})$ (so $p(\bar{x}, \bar{y})$ is a type over $\emptyset$). Moreover, if $tp(\bar{a}) = tp(\bar{b})$, then $p(\bar{x}, \bar{b})$ is in $S(\bar{b})$, and, for example, $p(\bar{x}, \bar{a})$ is strongly regular if, and only if, $p(\bar{x}, \bar{b})$ is strongly regular.

Secondly, suppose that $p \in S(A)$, and $q \in S(B)$ (A and B subsets of the big model). Then, because p and q are not types over the same set it does not make immediate sense to speak of, for example, p and q being orthogonal or not orthogonal. However we can interpret this to mean that for some C which includes A and B, any nonforling extensions of p and q over C are orthogonal (or not orthogonal, as the case might be). (We assume p and q to be stationary). Then by the results in section I, p and q will be orthogonal if, and only if, for any $C \supset A \cup B$ n the nonforking extensions of p and q over C are orthogonal.

Finally, we assume familiarity with the notion of strong type (denoted stp). The important facts are the following assuming ω-stability. If $p \in S_n(A)$, then there is $E \in FE_n(A)$ (that is, $E(\bar{x}, \bar{y})$ is an equivalence relation on n-tuples, definable over A, and with a finite number of classes), such that if $\bar{a}$ and $\bar{b}$ realise p then $\bar{a}$ and $\bar{b}$ have the same strong type over A ($stp(\bar{a}/A) = stp(\bar{b}/A)$) if, and only if, $\models E(\bar{a}, \bar{b})$. Also, if I is independent over A, and all

elements of I have the same strong type over A, then I is indiscernible over A. Moreover, if I and J are two such sets, and the elements of I and J have the same type over A, then $tp(I/A) = tp(J/A)$. (In the cases in which we shall be interested, A will be the empty set and so will be omitted.) (Also $stp(\bar{a}/A) = stp(\bar{b}/B)$ implies $tp(\bar{a}/A) = tp(\bar{b}/A)$.)

PROPOSITION III.2. - The following are equivalent (for the theory T).

(i) For all M, $\mu(M) \leq \aleph_0$.

(ii) T is non-multidimensional.

(iii) If $p(x, \bar{a}) \in S(\bar{a})$ is strongly regular, and $stp(\bar{a}) = stp(\bar{b})$, then $p(x, \bar{a})$ and $p(x, \bar{b})$ are not orthogonal (that is equivalent).

Proof. -

(i) implies (ii) is immediate.

(ii) $\Rightarrow$ (iii) : Suppose that $p(x, \bar{a}) \in S(\bar{a})$ is strongly regular, $stp(\bar{a}) = stp(\bar{b})$, but $p(x, \bar{a})$ and $p(x, \bar{b})$ are orthogonal. First we can assume that $\bar{a}$ and $\bar{b}$ are independent (For if not, then choose $\bar{c}$ such that $\bar{c}$ and $\bar{a} \frown \bar{b}$ are independent, and $stp(\bar{c}) = stp(\bar{a}) = stp(\bar{b})$. Then $p(x, \bar{a})$ and $p(x, \bar{c})$ are orthogonal). Let λ be any cardinal, and let $\{\bar{a}_\alpha ; \alpha < \lambda\}$ be an independent set of realisations of $tp(\bar{a})$, such that $\bar{a}_0 = \bar{a}$, $\bar{a}_1 = \bar{b}$, and, for all $\alpha < \lambda$, $stp(\bar{a}_\alpha) = stp(a)$. So $\{\bar{a}_\alpha ; \alpha < \lambda\}$ is indiscernible, and, for $\alpha < \beta < \lambda$, $p(x, \bar{a}_\alpha)$ and $p(x, \bar{a}_\beta)$ are orthogonal, (and strongly regular). Let M be a model containing all the $\bar{a}_\alpha$. For each $\alpha < \lambda$, let p_α be the nonforking extension of $p(x, \bar{a}_\alpha)$ over M. Then the p_α are pairwise orthogonal strongly regular types over M. Thus T is multidimensional.

(iii) $\Rightarrow$ (i) : Let M be a model, and $q \in S(M)$ strongly regular. There is finite $\bar{a}$ in M such that q is definable over $\bar{a}$. So $p = q \restriction \bar{a}$ is strongly regular, and q is the unique nonforking extension of p over M. Thus it suffices to show that there are at most $\aleph_0$ pairwise orthogonal strongly regular types over finite sets. Now there are only $\aleph_0$ many possible types of finite sets. Moreover for any $\bar{a}$, there are at most $\aleph_0$ types in $S_1(\bar{a})$. Also for any $\bar{a}$ and strongly regular $p(x, \bar{a}) \in S_1(\bar{a})$, there can be only finitely many pairwise orthogonal types of the form $p(x, \bar{b})$, where $tp(\bar{b}) = tp(\bar{a})$ (by (iii) and the paragraph preceding this proposition). Thus we finish.

PROPOSITION III.3. - Let T be non-multidimensional and N a model of T. Then there is a countable $M \prec N$, and a set $J \subset N$, J independent over M such that N is minimal over $M \cup J$.

Proof. - By III.2, $\mu(N)$ is countable. So we can find countable $M \prec N$ such that each of some maximal collection of pairwise orthogonal strongly regular types over N, is definable over M. Now use lemma 1.16.

T will be maid to be unidimensional if, for each $M \models T$, $\mu(M) = 1$.

PROPOSITION III.4. - T is unidimensional if, and only if, T is $\aleph_1$-categorical.

Proof. - Suppose that T is not unidimensional and let M be a model and p, q orthogonal strongly regular types over M. Assume that p and q are chosen with least possible Morley ranks in their respective equivalence classes, say $R(p)=\alpha$, $R(q) = \beta$, $\alpha \leqslant \beta$, and (p, φ) is strongly regular, where $R(\varphi) = \alpha$. As in the proof of 1.8, $\varphi(x)$ is not augmented in $M(q)$, and this, as is well known contradicts $\aleph_1$-categoricity.

Conversely, suppose that T is unidimensional. Let M_0 be the prime model of T. Then there is a strongly regular type p over M_0. If N is any model of T, then M_0 is elementarily embedded in N, and p' the heir of p over N, is strongly regular, and so is essentially the only strongly regular type over N. So N is prime over M_0 and a basis for p' in N. Such a basis is just a Morley sequence of p over M, and its type is determined. Thus if $|N_1| = |N_2| = \lambda > \aleph_0$, then N_1 is prime over $M_0 \cup I$ and N_2 is prime over $M_0 \cup J$, where I and J must both have cardinality λ, and have the same type over M_0. So $N_1 \cong N_2$.

PROPOSITION III.5. - Let T be non-multidimensional, and $p(x, \bar{a})$ a strongly regular type in $S(\bar{a})$. Suppose that $stp(\bar{b}) = stp(\bar{a})$ and M contains $\bar{a}$ and $\bar{b}$. Then $\dim(p(x, \bar{a}), M) = \dim(p(x, \bar{b}), M)$.

Proof. - Suppose first that $\bar{a}$ and $\bar{b}$ are independent (over $\emptyset$). Let $M_1 \prec M$ be prime over $\bar{a} \frown \bar{b}$, and let p_1, q_1 be the nonforking extensions of $p(x, \bar{a})$ and $p(x, \bar{b})$ over M_1. Now $tp(\bar{a} \frown \bar{b}) = tp(\bar{b} \frown \bar{a})$, and thus $(M_1, \bar{a}, \bar{b}) \equiv (M_1, \bar{b}, \bar{a})$, whereby $\dim(p(x, \bar{a}), M_1) = \dim(p(x, \bar{b}), M_1)$. By III.2, p_1 and q_1 are equivalent, and thus $\dim(p_1, M) = \dim(q_1, M)$. Thus by II.11, $\dim(p(x, \bar{a}), M) = \dim(p(x, \bar{b}), M)$.

Now in the general case, let $\bar{c}$ be such that $stp(\bar{c}) = stp(\bar{a}) = stp(\bar{b})$, and $\bar{c}$ and $\bar{a} \frown \bar{b}$ are independent (over $\emptyset$). Let $M' = M(\bar{c})$, and p', q' the nonforking extensions of $p(x, \bar{a})$ and $p(x, \bar{b})$ over M. Then $\dim(p', M') = \dim(q', M')$ (as p' and q' are strongly regular and equivalent), and both these dimensions are finite (otherwise $M' - M$ contains an infinite independent set over M, each element of which is dependent on $\bar{c}$ over M; which contradicts superstability). But by the first part of the proof,

$$\dim(p(x, \bar{c}), M') = \dim(p(x, \bar{a}), M') = \dim(p(x, \bar{b}), M'),$$

and we know that

$$\dim(p(x , \bar{a}) , M') = \dim(p(x , \bar{a}) , M) + \dim(p' , M')$$

and

$$\dim(p(x , \bar{b}) , M') = \dim(p(x , \bar{b}) , M) + \dim(q' , M') \qquad (II.11) .$$

Thus $\dim(p(x , \bar{a}) , M) = \dim(p(x , \bar{b}) , M)$, and we finish.

I now proceed to show that in the non-multidimensional case, all strongly regular types can be taken as being definable over the prime model of T (and thus in proposition III.3, M can be taken to be M_0 the prime model of T).

LEMMA III.6. - Let T be non-multidimensional. Let $M \prec M' \not\prec N$ be models. Then there is $c \in N - M'$, such that $tp(c/M')$ is strongly regular, and $tp(c/M')$ does not fork over M .

Proof. - Choose $c \in N - M'$ such that $tp(c/M)$ is of least possible Morley rank. Thus clearly there is $\bar{a} \in M$ and $\varphi(x , \bar{a}) \in tp(c/M)$, and for all $d \in (\varphi(x , a))^N - M'$, $tp(d/M) = tp(c/M)$. Let us denote $tp(c/M)$ by p . Now if $tp(c/M')$ does not fork over M (and so is the nonforking extension of p), then it is clear that $(tp(c/M') , \varphi)$ is strongly regular, and we finish. So let us assume that $tp(c/M')$ forks over M , and we seek a contradiction. Now, as $tp(c/M')$ forks over M (by our assumption), $R(tp(c/M')) < R(p)$. We can clearly assume that c has been chosen also to satisfy $R(tp(c/M'))$ being as small as possible (among those x in $N - M'$ for which $tp(x/M) = p$). So $tp(c/M')$ is strongly regular (I.4). Now let $\bar{b}_0$ be chosen in M' such that $tp(c/M')$ is definable over $\bar{b}_0$, and let $q(x , \bar{b}_0)$ denote $tp(c/\bar{b}_0)$. Thus $q(x , \bar{b}_0)$ is strongly regular. Now let $\bar{b}_1$ be such that $tp(\bar{b}_1/M) = tp(\bar{b}_0/M)$ and $\bar{b}_0$ and $\bar{b}_1$ are independent over M .

Thus $stp(\bar{b}_0) = stp(\bar{b}_1)$ (this is easy), and so by III.2, $q(x , \bar{b}_0)$ and $q(x , \bar{b}_1)$ are equivalent. Let q_0 and q_1 be the nonforking extensions f , $q(x , \bar{b}_0)$ and $q(x , \bar{b}_1)$ respectively over $M \cup \{\bar{b}_0 , \bar{b}_1\}$. (So in particular $q_0 \restriction M \cup \bar{b}_0 = tp(c/M \cup \bar{b}_0)$.) So q_0 and q_1 are strongly regular types over the same set which are not orthogonal. Thus by II.5, there are $n , m < \omega$ such that $q_0^n(\bar{x}) \cup q_1^m(\bar{y})$ is not a complete type over $M \cup \{\bar{b}_0 , \bar{b}_1\}$. Thus (as q_0 and q_1 are stationary), there are $c_1 , \ldots , c_n$ independent realisations of q_0 over $M \cup \{\bar{b}_0 , \bar{b}_1\}$, and $d_1 , \ldots , d_m$ independent realisations of q_1 over $M \cup \{\bar{b}_0 , \bar{b}_1\}$ such that $\{c_1 , \ldots , c_n\}$ and $\{d_1 , \ldots , d_m\}$ are not independent over $M \cup \{\bar{b}_0 , \bar{b}_1\}$. By minimalising m , we can assume that $\{c_1 , \ldots , c_n\}$ and $\{d_1 , \ldots , d_{m-1}\}$ are independent over $M \cup \{\bar{b}_0 , \bar{b}_1\}$. Let us denote $\langle c_1 , \ldots , c_n \rangle$ by $\bar{c}$ and $\langle d_1 , \ldots , d_{m-1} \rangle$ by $\bar{d}$. I assert that

(*) $\bar{b}_0 \hat{} \bar{c}$ and $\bar{b}_1 \hat{} \bar{d}$ are independent over M .

First note that $tp(\bar{d}/\bar{b}_0 \cup \bar{b}_1 \cup M)$ does not fork over $\bar{b}_1 \cup M$, and that $tp(\bar{b}_1/\bar{b}_0 \cup M)$ does not fork over M. Thus $tp(\bar{b}_1 \hat{} \bar{d}/\bar{b}_0 \cup M)$ does not fork over M, and so

(i) $tp(\bar{b}_0/\bar{b}_1 \hat{} \bar{d} \cup M)$ does not fork over M.

Also $tp(\bar{c}'/\bar{b}_0 \cup \bar{b}_1 \hat{} \bar{d} \cup M)$ does not fork over $M \cup \bar{b}_0$. This together with (i) yields $tp(\bar{b}_0 \hat{} \bar{c}'/\bar{b}_1 \hat{} \bar{d} \cup M)$ does not fork over M, which means (*)

Note also that $tp(c_n/\{c_1, \ldots, c_{n-1}\} \cup \bar{d} \cup \bar{b}_0 \cup \bar{b}_1 \cup M)$ does not fork over $M \cup \{\bar{b}_0, \bar{b}_1\}$, but that

(**) $tp(c_n/\{c_1, \ldots, c_{n-1}\} \cup \bar{d} \hat{} d_n \cup \bar{b}_0 \cup \bar{b}_1 \cup M)$ does not fork over $M \cup \{\bar{b}_0, \bar{b}_1\}$

Now $tp(c_n/M \cup \bar{b}_0) = tp(c/M \cup \bar{b}_0)$. So <u>we can assume that</u> $c_n = c$ (leave $\bar{b}_0$ fixed but shift around the other c_i's, the d_i's and $\bar{b}_1$ so as to preserve the type of everything over M), <u>let us denote</u> d_m by d. So $tp(d \hat{} \bar{b}_1/M) = tp(c \hat{} \bar{b}_0/M)$, whereby $tp(d/M) = p$, and $tp(d/M \cup \bar{b}_1)$ forks over M, and so there is finite $\Delta \subset L$ such that

(***) $R(tp(d/M \cup \bar{b}_1), \Delta, 2) < R(p, \Delta, 2) = r$.

Let us now sum up the information obtained; denoting now $\langle c_1, \ldots, c_{n-1} \rangle$ by $\bar{c}$, and as before $\langle d_1, \ldots, d_{m-1} \rangle$ by $\bar{d}$.

(a) c and $\bar{c}$ are independent over $M \cup \bar{b}_0$.

(b) $\bar{b}_0 \hat{} \bar{c} \hat{} c$ and $\bar{b}_1 \hat{} \bar{d}$ are independent over M (by (*)).

(c) There is a formula $\chi(x, \bar{z})$ and $\bar{e} \in M$ such that $\models \chi(c, d \hat{} \bar{d} \hat{} \bar{c} \hat{} \bar{b}_0 \hat{} \bar{b}_1 \hat{} \bar{e})$, but $\chi(x, \bar{z})$ is not in bound $(tp(c/\bar{b}_0))$ (and so $\chi(x, \bar{z})$ is not represented in $tp(c/M')$) (by (**)).

(d) There is an $L(M)$ formula $\psi(x, \bar{w})$ such that d satisfies $\psi(x, \bar{b}_1)$ and $R(\psi(x, b_1), \Delta, 2) < r$ (by (***)).

(Remember for any type q and finite $\Delta \subset L$, there is finite subtype of q, say q' such that $R(q, \Delta, 2) = R(q', \Delta, 2)$.)

Remember that d also satisfies the formula $\varphi(x, \bar{a})$. Thus by (c) and (d), we have

$\models (\exists y)(\varphi(y, \bar{a}) \wedge \chi(c, y \hat{} \bar{d} \hat{} \bar{c} \hat{} \bar{b}_0 \hat{} \bar{b}_1 \hat{} \bar{e}) \wedge \psi(y, \bar{b}_1) \wedge "R(\psi(x, \bar{b}_1), \Delta, 2) < r")$.

By (b) we can find $\bar{b}_1'$ and $\bar{d}'$ in M such that

$\models (\exists y)(\varphi(y, \bar{a}) \wedge \chi(c, y \hat{} \bar{d}' \hat{} \bar{c} \hat{} \bar{b}_0 \hat{} \bar{b}_1' \hat{} \bar{e}) \wedge \psi(y, \bar{b}_1') \wedge "R(\psi(x, \bar{b}_1'), \Delta, 2) < r")$.

Now by (a) and the fact that $tp(c/M')$ is definable over $M \cup \bar{b}_0$, we can find $\bar{c}' \in M'$ such that

$N \models (\exists y)(\varphi(y , \bar{a}) \wedge \chi(c , y\, \bar{d}'^\frown\bar{c}^\frown\bar{b}_0^\frown\bar{b}_1^\frown\bar{e}) \wedge \psi(y , \bar{b}_1') \wedge \text{"}R(\psi(x , \bar{b}_1') , \Delta , 2) < r\text{"})$.

Pick $a \in N$ to be such a y as given above. First note that $a \notin M'$, for if not then $\chi(x , \bar{z})$ would be represented in $tp(c/M')$, contradicting (c). Thus as a satisfies $\varphi(x , \bar{a})$, we must have $tp(a/M) = p$ (by choice of p and $\varphi(x , \bar{a})$). But now, as a satisfies $\psi(x , \bar{b}_1')$ and $R(\psi(x , \bar{b}_1') , \Delta , 2) < r = R(p , \Delta , 2)$, we must have that $tp(a/M) \neq p$. This contradiction proves the lemma.

PROPOSITION III.7. - Let $M < M'$ be models of T, where T is non-multidimensional, and let $p \in S(M')$ be strongly regular. Then there is $q \in S(M')$, such that q is strongly regular, q is equivalent of p, and q does not fork over M.

Proof. - Lemma III.6 gives us c in $M'(p) - M'$ such that $tp(c/M')$ is strongly regular, and does not fork over M. Clearly $tp(c/M')$ is equivalent to p.

COROLLARY III.8. - Let T be non-multidimensional. Let M be a model, A a set, and N prime over $M \cup A$. Then N is minimal over $M \cup A$.

Proof. - If not, there is model M' such that $M \cup A \subset M' \not\prec N$. Lemma III.6 gives us $c \in N - M'$ such that $tp(c/M')$ does not fork over M. But $tp(c/M)$ is not isolated, and $tp(c/M \cup A)$ is isolated, whereby $tp(c/M \cup A)$ forks over M, and so $tp(c/M')$ forks over M. Contradiction.

Let me now state a few obvious things. Let us assume T to be non-multidimensional, and let M_0 be the prime model of T. Let $\{p_i ; i < \mu \leq \aleph_0\}$ be a maximal collection of pairwise orthogonal strongly regular types over M_0. Let N be any model of T. So M_0 is elementarily embedded in N, and let p_i' for $i < \mu$, be the heirs of the p_i over N. Then $\{p_i' ; i < \mu\}$ is a maximal collection of pairwise orthogonal strongly regular types over N. For choose strongly regular $q \in S(N)$. By III.7, q is equivalent of $q \in S(N)$, where p is strongly regular and does not fork over M_0. But there is $i < \mu$ such that $p \restriction M_0$ is equivalent to p_i and so p is equivalent to p_i', and so q is equivalent to p_i'.

IV. The spectrum.

In this section T will be assumed to be non-multidimensional, and M_0 will denote the prime model of T.

First, some more preliminary results.

LEMMA IV.1. - Let M be a model, $\bar{a} \in M$, $p(x , \bar{a}) \in S(\bar{a})$ be strongly regular, and $tp(\bar{a})$ isolated. Suppose that $\bar{b} \in M$, $tp(\bar{b}) = tp(\bar{a})$ and $p(x , \bar{b})$ is equivalent to $p(x , \bar{a})$. Then $\dim(p(x , \bar{a}) , M) = \dim(p(x , \bar{b}) , M)$.

Proof. - Let $M_0 < M$ be a copy of the prime model such that $\bar{a} \in M_0$. It is easy

to find $\bar{c} \in M_0$ such that $stp(\bar{c}) = stp(\bar{b})$. By III.2, $p(x , \bar{b})$ and $p(x , \bar{c})$ are equivalent. Thus $p(x , \bar{a})$ and $p(x , \bar{c})$ are equivalent. Let p_1 and p_2 be the nonforking extensions of $p(x , \bar{a})$ and $p(x , \bar{c})$ over M_0. So p_1 and p_2 are equivalent and strongly regular, and thus by II.8, $dim(p_1 , M) = dim(p_2,M)$. But it is clear that $(M_0 , \bar{a}) \cong (M_0 , \bar{c})$, and so $dim(p(x,\bar{a}),M_0)=dim(p(x,\bar{c}),M_0)$. Thus by II.11, we have

$$dim(p(x , \bar{a}) , M) = dim(p(x , \bar{c}) , M) .$$

But by III.5,

$$dim(p(x , \bar{c}) , M) = dim(p(x , \bar{b}) , M) ,$$

and so we have

$$dim(p(x , \bar{a}) , M) = dim(p(x , \bar{b}) , M) ,$$

as desired.

LEMMA IV.2. (which does not need non-multidimensionality). - Let $p \in S(M_0)$, p definable over $\bar{a} \in M_0$, $p_1 = p \restriction \bar{a}$, and p_1 has an infinite basis in M_0 (thus $dim(p_1 , M_0) = \aleph_0$). Then $M_0(p) \cong M_0$.

Proof. - $M_0(p)$ is countable, and thus it is enough to show that $M_0(p)$ is atomic (i. e. realises only isolated types). Let $\bar{c} \in M_0(p)$ be such that $tp(\bar{c}/M_0) = p$ and $M_0(p)$ is atomic over $M_0 \cup \bar{c}$. It is enough to show that $M_0 \cup \bar{c}$ is atomic. So let $\bar{b} \in M_0$. I show that $tp(\bar{b} \hat{} \bar{c})$ is isolated, in fact that $tp(\bar{a} \hat{} \bar{b} \hat{} \bar{c})$ is isolated. Let $\bar{c}_i$, for $i < \omega$, be a basis for $p_1 = p \restriction \bar{a}$ in M_0. Then by superstability, there must be $i < \omega$ such that $\bar{c}_i$ and $\bar{b}$ are independent over $\bar{a}$. Then clearly $tp(\bar{a} \hat{} \bar{b} \hat{} \bar{c}_i) = tp(\bar{a} \hat{} \bar{b} \hat{} \bar{c})$, and $tp(\bar{a} \hat{} \bar{b} \hat{} \bar{c}_i)$ is isolated, as it is realised in the prime model M_0. So we finish.

Note. - An extension of the above proof shows that if $p \in S(M)$ and for some $\bar{a} \in M$ over which p is definable, $p \restriction \bar{a}$ has an infinite basis in M, then for all $\bar{a} \in M$ over which p is definable $p \restriction \bar{a}$ has an infinite basis in M.

COROLLARY IV. 3. - Let $\{p_i ; i < \kappa (\leq \aleph_0)\}$ be a set of pairwise orthogonal strongly regular types over M_0, such that for each i there is $\bar{a}_i \in M_0$ such that p_i is definable over $\bar{a}_i$, and $dim(p_i \restriction \bar{a}_i , M_0)$ is infinite. For each $i < \kappa$, let J_i be an independent set of realisations of p_i over M_0, such that $|J_i| \leq \omega$. Then $M_0(\bigcup_{i<\kappa} J_i) \cong M_0$.

Proof. - It is easy, using IV.2, induction and fact 1.6, to show that $M_0(J_0)$ is isomorphic to M_0 (let $J_0 = \{c_n ; n < \lambda\}$, let $M_1 = M_0(c_0)$, and in general $M_{n+1} = M_n(c_n)$. Then $tp(c_n/M_n)$ is the heir of p_0 over M_n, and $M_{n+1} \cong M_0$. So $\bigcup_{n<\lambda} M_n$ is isomorphic to M_0, and is also easily see to be the same as

$M_0(J_0)$). Then it is easy to see that $tp(J_1/M_0(J_0))$ does not fork over M_0, and so we can repeat the process to get $M_0(J_0)(J_1) \cong M_0$. Carry on, and putting $M^0 = M_0$, and $M^{n+1} = M^n(J_n)$, we see that $\bigcup_{n<\omega} M^n$ is isomorphic to M_0 and is the same as $M_0(\bigcup_{i<\omega} J_i)$.

LEMMA IV.4. - Let $\{p_i ; i < \omega\}$ be pairwise orthogonal types over a model M, and let for each $i < \omega$, J_i be a set of independent realisations of p_i over M.

Let N be prime over $M \cup \bigcup_{i<\omega} J_i$. Then for each $i < \omega$, J_i is a basis for p_i in N.

Proof. - Consider J_0 for example. Let us define $M_i \prec N$ for $1 \leq i \leq \omega$, such that M_1 is prime over $M \cup J_1$, and for $i \geq 1$, M_{i+1} is prime over $M_i \cup J_{i+1}$ and $M_\delta = \bigcup_{i<\delta} M_i$ for δ limit. Let M' be M_ω. Let p_0^i be the heir of p_0 over M_i for $i \leq \omega$. Then it is easy to show by induction, using the orthogonality of p_0 and the p_i's and fact 1.6, that $p_0 \vdash p_0^i$ for $1 \leq i \leq \omega$. Thus J_0 is a basis for p_0^ω in N if, and only if, J_0 is a basis for p_0 in N, and clearly J_0 is an independent set of realisations of p_0^ω over M' in N. By III.8 for example, N is prime over $M' \cup J_0$, and so J_0 is easily seen to be a basis for p_0^ω in N. So the lemma is proved.

LEMMA IV.5. - Let $p \in S(\bar{a})$ be strongly regular, where $tp(\bar{a})$ is isolated, and for some copy of M_0 which contains $\bar{a}$, $\dim(p, M_0) = 0$. Let A be any countable set which is atomic over $\bar{a}$, and let p' be the nonforking extension of p over $A \cup \bar{a}$. Then $p \vdash p'$.

Proof. - Let A be as given. Then $A \cup \bar{a}$ is an atomic countable set, and we can find a copy of the prime model M_0 such that $A \subset M_0$. By isomorphism, p is not realised in M_0. So by lemma II.11, for any c realising p, $tp(c/M_0)$ does not fork over $\bar{a}$, and thus $tp(c/A \cup \bar{a})$ does not fork over $\bar{a}$. So clearly $p \vdash p'$

We can now begin on the classification. First let μ be the maximum number of pairwise orthogonal strongly regular types over M_0, the prime model of T. (We call μ the number of dimensions of T). Let p_i for $i < \mu$, be pairwise orthogonal and strongly regular types over M_0, and a maximal such collection. Now let N be any model. So $M_0 \prec N$, and (by 1.16, III.7 and remarks at the end of III) N is prime over (in fact minimal over) $M_0 \cup \bigcup_{i<\mu} J_i$ where J_i is a basis for p_i in N, and moreover (by 1.15) $tp(\bigcup_{i<\mu} J_i/M_0)$ is determined just by $\langle \lambda_i ; i < \mu \rangle$ where $\lambda_i = |J_i|$. Conversely, given a sequence $\langle \lambda_i ; i < \mu \rangle$ of cardinals, there is a model N prime over $M_0 \cup \bigcup_{i<\mu} J_i$ where J_i is an independent set of realisations of p_i, and thus by IV.4, a basis for p_i in N. So if we are considering the models of T up to isomorphism over some fixed copy of the prime model M_0 (which we could do by for example adding names for the elements of M_0 to the langage, and replacing T by $Th(M_0)$ in this new language), then

the models would correspond exactly to the possible sequences of cardinals $\langle \lambda_i ; i < \mu \rangle$. However in the general case, one model might contain different copies of M_0 and correspond to different sequences of cardinals. So we have to be more careful in the choices of the p_i , and use some material developed in this section and section III. This we proceed to do, summing up the results later on in a theorem.

First let K_i , for $i < \mu$, be the equivalence classes (or non-rothogonality classes) of strongly regular types over M_0 . We choose, for each $i < \mu$, $p_i \in K_i$ and $\bar{a}_i \in M_0$, such that p_i is definable over $\bar{a}_i$, and also satisfying the following two conditions, where $q_i(x , \bar{a}_i)$ denotes $p_i \restriction \bar{a}_i$ (so $q_i(x , \bar{y}_i)$ is over $\emptyset$) :

(i) $\dim(q_i(x , \bar{a}_i) , M_0)$ is 0 or infinite (i. e. $\aleph_0$), for all $i < \mu$, and

(ii) if $i < j < \mu$, then <u>either</u> $tp(\bar{a}_i) = tp(\bar{a}_j)$ and $q_i(x , \bar{y}_i) = q_j(x , \bar{y}_j)$, <u>or</u> for <u>no</u> $p \in K_j$ is there $\bar{a} \in M_0$ such that p is definable over $\bar{a}$, $tp(\bar{a}) = tp(\bar{a}_i)$ and $p \restriction \bar{a} = q_i(x , \bar{a})$.

(Note that if the second disjunct of (ii) holds, then we also have that for <u>no</u> $p \in K_i$ is there $\bar{a} \in M_0$ such that p is definable over $\bar{a}$, $tp(\bar{a}) = tp(\bar{a}_j)$ and $p \restriction \bar{a} = q_j(x , \bar{a})$.)

This is achieved quite easily. To get (i) for example, suppose p_i has been chosen in K_i , and, for some $\bar{a} \in M_0$, p_i is definable over $\bar{a}$ and $\dim(p_i \restriction \bar{a} , M_0) = n < \omega$. Let $c_1 , \ldots , c_n$ be a basis for $p_i \restriction \bar{a}$ in M_0 , and put $\bar{a}_i = \bar{a} \hat{} \langle c_1 , \ldots , c_n \rangle$. Then clearly $\dim(p_i \restriction \bar{a}_i , M_0) = 0$. (ii) can easily be obtained by defining the p_i and $\bar{a}_i$ inductively.

This having been done, pick some particular $i < \mu$, and let us put $p = p_i$, $a = \bar{a}_i$, and $q(x , \bar{y}) = q_i(x , \bar{y}_i)$. For how many $j < \mu$, do we have $tp(\bar{a}_j) = tp(\bar{a})$ and $q_j(x , \bar{y}_j) = q(x , \bar{y})$ (and thus $p_j \restriction \bar{a}_j = q(x , \bar{a}_j)$) ? I assert that there can be only finitely many such j . For if not, then there is infinite $J \subset \omega$, such that the types $\{q(x , \bar{a}_j) ; j \in J\}$ are pairwise orthogonal, and $tp(\bar{a}_j) = tp(\bar{a})$ for all $j \in J$. Thus (see background at the beginning of section III), there is $j_1 < j_2$ in J such that $stp(\bar{a}_{j_1}) = stp(\bar{a}_{j_2})$. But by III.2, this contradicts the orthogonality of $q(x , \bar{a}_{j_1})$ and $q(x , \bar{a}_{j_2})$. (Remember $q(x , \bar{a})$ is strongly regular). Thus there are only finitely many such j .

Thus by renumbering the q_i and renaming the p_i and $\bar{a}_i$, we have :

LEMMA IV.6. - <u>There is</u> $\mu' \leq \aleph_0$, <u>and for each</u> $i < \mu'$, <u>some finite</u> n_i , <u>and</u> $q(x , \bar{y}_i)$ <u>over</u> $\emptyset$, <u>and for each</u> $i < \mu'$ <u>and</u> $j < n_i$, <u>types</u> p_i^j <u>over</u> M_0 <u>and tuples</u> $\bar{a}_i^j$ <u>in</u> M_0 <u>such that</u>

(i) $\{p_i^j ; i < \mu' , j < n_i\}$ <u>is a maximal collection of pairwise orthogonal strongly regular types over</u> M_0 .

(ii) p_i^j is definable over $\bar{a}_i^j$,

(iii) for any $i < \mu'$, for $j_1 < j_2 < n_i$, $tp(\bar{a}_i^{j_1}) = tp(\bar{a}_i^{j_2}) = r_i$, and, for each $j < n_i$, $p_i^j \restriction a_i^j = q_i(x , \bar{a}_i^j)$,

(iv) for each i and j , $\dim(a_i(x , \bar{a}_i^j) , M_0) = 0$ or $\aleph_0$,

(v) if $i_1 < i_2 < \mu'$, then there are no $\bar{a}_1$, $\bar{a}_2$ in M_0 such that $tp(\bar{a}_1) = r_{i_1}$ and $tp(\bar{a}_2) = r_{i_2}$, and $q_{i_1}(x , \bar{a}_1)$ is equivalent to $q_{i_2}(x , \bar{a}_2)$.

(vi) $\mu' = \aleph_0$ if, and only if, $\mu = \aleph_0$, and $\mu' = 1$ if, and only if, $\mu = 1$.

LEMMA IV.7. - Let N be any model of T , and let $i_1 < i_2 < \mu'$. Then there are no $\bar{a}_1$ and $\bar{a}_2$ in N such that $tp(\bar{a}_1) = r_{i_1}$, $tp(\bar{a}_2) = r_{i_2}$, and $q_{i_1}(x , \bar{a}_1)$ is equivalent of $q_{i_2}(x , \bar{a}_2)$.

Proof. - Suppose that there are $\bar{a}_1$ and $\bar{a}_2$ in N as described, and we get a contradiction. Let M_0 be some copy of the prime model in N . Now both r_{i_1} and r_{i_2} are isolated types, and so it is easy to find $\bar{a}'_1$ and $\bar{a}'_2$ in M_0 such that $stp(\bar{a}'_1) = stp(\bar{a}_1)$ and $stp(\bar{a}'_2) = stp(\bar{a}_2)$. Thus by III.2, $q_{i_1}(x , \bar{a}'_1)$ is equivalent to $q_{i_1}(x , \bar{a}_1)$, and $q_{i_2}(x , \bar{a}'_2)$ is equivalent to $q_{i_2}(x , \bar{a}_2)$. But then $q_{i_1}(x , \bar{a}'_1)^1$ is equivalent to $^2q_{i_2}(x , \bar{a}'_2)$, which contradicts lemma IV.6 (v).

Now we go through the cases depending on the number of dimensions.

Case 1. - μ is finite. So also μ' is finite. Let $A = \bigcup\{\bar{a}_i^j ; i < \mu' , j < n_i\}$ and let q_i^j be the nonforling extension of $q_i(x , \bar{a}_i^j)$ over A . Let λ_i^j for $i < \mu'$ and $j < n_i$ be cardinals chosen arbitrarily subject to the proviso that $\lambda_i^j \geq \aleph_0$ if $\dim(q_i(x , \bar{a}_i^j) , M_0) = \aleph_0$. Let $A(\langle\lambda_i^j ; i < \mu' , j < n_i\rangle)$ denote the model prime over $A \cup \bigcup(I_i^j ; i < \mu' , j < n_i)$, where I_i^j is an independent set of realisations of q_i^j over A of cardinality λ_i^j . Note that $A(\bar{\lambda})$ (where $\bar{\lambda} = \langle\lambda_i^j ; i < \mu' , j < n_i\rangle$) is well defined by 1.15 and uniqueness of prime models.

Observation IV.8.

(i) $\dim(q_i^j , A(\bar{\lambda})) = \lambda_i^j$.

(ii) $\dim(q_i(x , \bar{a}_i^j) , A(\bar{\lambda})) = \lambda_i^j$.

Proof.

(i) Let M be prime over $M_0 \cup \bigcup\{X_i^j ; i < \mu' , j < n_i\}$ where X_i^j is an independent set of realisations of p_i^j over M_0 of cardinality λ_i^j . Then $\dim(p_i^j , M) = \lambda_i^j$, by lemma IV.4. It is easily seen that M is isomorphic (over A) to $A(\bar{\lambda})$, and that (by II.11 and choice of p_i^j and $\bar{a}_i^j$) that $\dim(q_i^j , M) = \lambda_i^j$.

(ii) We use (i). First suppose that $\dim(q_i(x , \bar{a}_i^j) , M_0) = 0$. Then as $tp(A/\bar{a}_i^j)$ is isolated, we have by IV.5 that $q_i(x , \bar{a}_i^j) \vdash q_i^j$, and thus

$$\dim(q_i(x\ ,\ \bar{a}_i^j)\ ,\ A(\bar{\lambda})) = \dim(q_i^j\ ,\ A(\bar{\lambda})) = \lambda_i^j\ .$$

Secondly, suppose that $\dim(q_i(x\ ,\ \bar{a}_i^j)\ ,\ M_0)$ is infinite. Then so must be $\dim(q_i(x\ ,\ \bar{a}_i^j)\ ,\ A(\bar{\lambda}))$. But only finitely many members of a basis for $q_i(x\ ,\ \bar{a}_i^j)$ in $A(\bar{\lambda})$ can be made to fork by $A - \bar{a}_i^j$ (remember that A is finite at the moment). Thus clearly $\dim(q_i(x\ ,\ \bar{a}_i^j)\ ,\ A(\bar{\lambda})) = \dim(q_i^j\ ,\ A(\bar{\lambda})) = \lambda_i^j$.

Conversely we know that <u>any</u> model N of T can be written as (i. e. is isomorphic to) $A(\langle \lambda_i^j\ ;\ i < \mu'\ ,\ j < n_i \rangle)$, where λ_i^j must be infinite if $\dim(q_i(x\ ,\ a_i^j)\ ,\ M_0)$ is infinite (by I. 16, III.7, and remarks at the end of section III). It is also clear that $|A(\bar{\lambda})| = \max(\{\lambda_i^j\ ;\ i < \mu'\ ,\ j < n_i\} \cup \{\aleph_0\})$. When is $A(\lambda) \cong A(\bar{\lambda}^*)$.

<u>Case 1 (i)</u>. - $\mu = 1$. So $\mu' = 1$, and $n_0 = 1$. Also $A = \bar{a}_0^0$. Let us write $\bar{a}_0^0$ as $\bar{a}$ and $q_0(x\ ,\ \bar{y}_0)$ as $q(x\ ,\ \bar{y})$. Now suppose that $M = \bar{a}(\lambda) \cong \bar{a}(\lambda^*)$. Then there is $\bar{a}^* \in M$, $tp(\bar{a}^*) = tp(\bar{a})$, and $M = a^*(\lambda^*)$. So $\dim(q(x\ ,\ \bar{a}), M) = \lambda$, and $\dim(q(x\ ,\ \bar{a}^*)\ ,\ M) = \lambda^*$ (by IV.8 (ii)). But as $\mu = 1$, we must have that $q(x\ ,\ \bar{a})$ and $q(x\ ,\ \bar{a}^*)$ are equivalent, but then by lemma IV.1, we have that $\lambda = \lambda^*$. So we have $\bar{a}(\lambda) \cong \bar{a}(\lambda^*)$ if, and only if, $\lambda = \lambda^*$. Thus in this case

$$I(\kappa\ ,\ T) = 1 \quad \text{if} \quad \kappa > \aleph_0\ .$$

If $\dim(q(x\ ,\ \bar{a})\ ,\ M_0) = 0$, then

$$I(\aleph_0\ ,\ T) = \aleph_0 \quad \text{(as all finite dimensions can occur)},$$

and if $\dim(q(x\ ,\ \bar{a})\ ,\ M_0)$ is infinite, then

$$I(\aleph_0\ ,\ T) = 1\ .$$

<u>Case 1 (ii)</u>. - $\mu > 1$ (but still finite).

Let $\bar{\mu}$ denote $\langle \mu_i^j\ ;\ i < \mu'\ ,\ j < n_i \rangle$ (no connection with μ, the number of dimension). Suppose that $N = A(\bar{\lambda}) \cong A(\bar{\mu})$. Thus there is A^* in N with $tp(A^*) = tp(A)$, and $N = A^*(\bar{\mu})$. Denote by $\bar{a}_i^{j*}$ the copy of $\bar{a}_i^j$ in A^*. Then $\{q_i(x\ ,\ \bar{a}_i^{j*})\ ;\ i < \mu'\ ,\ j < n_i\}$ is a set of pairwise orthogonal strongly regular types, and by IV.8 (ii), $\dim(q_i(x\ ,\ \bar{a}_i^{j*})\ ,\ M_0) = \mu_i^j$. So as the $q_i(x\ ,\ \bar{a}_i^j)$ are a maximal collection of pairwise orthogonal strongly regular types, and by lemma IV.7, there is σ such that for each $i < \mu'$, $\sigma(i\ ,\ -)$ is a permutation of n_i and $q_i(x\ ,\ a_i^{j*})$ is equivalent to $q_i{}'x\ ,\ a_i^{\sigma(i,j)})$. Thus by lemma IV.1, $\mu_i^j = \lambda_i^{\sigma(i,j)}$. Thus $A(\bar{\lambda}) \cong A(\bar{\mu})$ implies that $\bar{\mu} = \sigma(\bar{\lambda})$, where σ is a permutation of the sequence $\bar{\lambda}$ (As the number of dimensions is finite, there can only be finitely many such permutations).

<u>Case 1 (ii) (a)</u>. - For some $i < \mu'$, $j < n_i$, $\dim(q_i(x\ ,\ \bar{a}_i^j)\ ,\ M_0) = 0$. Then all cardinals (including finite ones) are possible for λ_i^j. Thus the number of sequences of cardinals $\leq \aleph_\alpha$, $\langle \lambda_i^j\ ;\ i < \mu'\ ,\ j < n_i \rangle$ at least one member of which is $\aleph_\alpha$, is $|\alpha| + \aleph_0$. (Note that in this case $|A(\bar{\lambda})| = \aleph_\alpha$.) But by the

above there can be only finitely many other sequences $\bar{\mu}$ giving rise to the same model, and thus we have

$$I(\aleph_\alpha , T) = |\alpha| + \aleph_0 , \text{ for all } \alpha \geqslant 0 .$$

Case 1 (ii) (b). - For all $i < \mu'$, $j < n_i$, $\dim(q_i(x , \bar{a}_i^j) , M_0) = \aleph_0$. But then the countable models of T are just models isomorphic to $A(\bar{\lambda}^0)$, and thus T is $\aleph_0$-categorical, i. e. $I(\aleph_0 , T) = 1$. Now suppose that $A(\bar{\lambda}) \cong A(\bar{\mu})$ as above and thus that there is $A^* \subset N = A(\kappa)$, with $N = A^*(\bar{\mu})$, and σ with $q_i(x , \bar{a}_i^{j*})$ equivalent to $q_i(x , \bar{a}_i^{\sigma(i,j)})$. Then as T is $\aleph_0$-categorical, all types are isolated, and thus $tp(A \frown A^*)$ is realised in every model of T . Clearly the fact that $q_i(x , \bar{a}_i^j)$ is equivalent ot $q_i(x , \bar{a}_i^{j'})$, say, depends only on $tp(\bar{a}_i^j \frown \bar{a}_i^{j'})$. So we let G denote the group of permutations σ of μ , induced as above, and clearly $A(\bar{\lambda}) \cong A(\bar{\mu})$ if, and only if, there is $\sigma \in G$ with $\sigma(\bar{\lambda}) = (\bar{\mu})$. By our case hypothesis, only infinite values are possible for the λ_i^j . Let us denote by $(|\alpha + 1|^\mu)^*$ the number of sequences of length μ of ordinals $\leqslant \alpha$, at least one of which is α . Thus it is clear that

$$I(\aleph_\alpha , T) = (|\kappa + 1|^\mu)^*/G , \text{ for all } \alpha \geqslant 0 ;$$

Case 2. - $\mu = \aleph_0$, and so μ' is also $\aleph_0$.

Let me denote by $M_0(\lambda_i^j)_{i,j}$ the model prime over $M_0 \cup U\{I_i^j ; i < \mu' , j < n_i\}$ where I_i^j is an independent set of realisations of p_i^j over M_0 . We know that any λ_i^j can occur. I first want to observe that if $\dim(q_i(x , \bar{a}_i^j) , M_0) = \aleph_0$, then we can assume that λ_i^j is always 0 or uncountable.

LEMMA IV. 9. - Let $N = M_0(\lambda_i^j)_{i,j}$, where, for $(i , j) \in X$, $\dim(q_i(x , \bar{a}_i^j) , M_0) = \aleph_0$, and $\lambda_i^j \leqslant \aleph_0$. Then $N \cong M_0(\lambda_i^{j*})_{i,j}$ where $\lambda_i^{j*} = \lambda_i^j$ if $(i , j) \notin X$, and $\lambda_i^{j*} = 0$ if $(i , j) \in X$.

Proof. - Easy using IV.3 and IV.5.

Thus the models of T are all of the form $M_0(\bar{\lambda})$ where λ_i^j can be anything, if $\dim(q_i(x , a_i^j) , M_0) = 0$, and is 0 or uncountable otherwise. Moreover, it is easy to see, using II.11 and IV.5, that $\dim(q_i(x , \bar{a}_i^j) , M_0(\bar{\lambda})) = \lambda_i^j$, if $\dim(q_i(x , \bar{a}_i^j) , M_0) = 0$, and $= \aleph_0 + \lambda_i^j$ otherwise. It is also clear by IV.4, that $\dim(p_i^j , M_0(\bar{\lambda})) = \lambda_i^j$. Thus, as in case 1, it follows that if $M_0(\bar{\lambda}) \cong M_0(\bar{\mu})$, then there is σ such that for $i < \mu'$, and $j < n_i$, $\sigma(i , j) < n_i$, and for all j_1 , j_2 , $\mu_i^{j_2} = \lambda_i^{\sigma(i,j)}$. But μ' is infinite, and if $i_1 < i_2 < \aleph_0$, we can vary $\lambda_{i_1}^{j_1}$ and $\lambda_{i_2}^{j_2}$ (j_1 , j_2 arbitrary), to get different models. Thus it is clear that $I(\aleph_\alpha , T) = \prod_{i<\aleph_0} \kappa_i$, where $\kappa_i = |\alpha| + \aleph_0$; if $\dim(q_i(x , a_i^0) , M_0) = 0$, and $\kappa_i = |\alpha| + 1$ otherwise.

Thus we have proved :

THEOREM IV.10. - Let T be non-multidimensional ω-stable. Let $I(\aleph_\alpha, T)$ denote the number of models of T of power $\aleph_\alpha$ up to isomorphism. Then there is $\mu \leq \aleph_0$, where μ is called the number of dimensions of T, such that :

1° if $\mu = 1$, then $I(\aleph_\alpha, T) = 1$ for all $\alpha > 0$, and $I(\aleph_0, T) = 1$ or $\aleph_0$.

2° If $\mu > 1$ but finite, then either $I(\aleph_\alpha, T) = |\alpha + \omega|$, for all $\alpha \geq 0$, or $I(\aleph_0, T) = 1$ and there is G a group of permutations of μ such that for $\alpha > 0$ $I(\aleph_\alpha, T) = (|\alpha + 1|^\mu)^*/G$, where $(|\alpha + 1|^\mu)^*$ is the number of sequences of length μ of ordinals $\leq \alpha$ at least one of which is α, and $\langle \beta_i ; i < \mu \rangle \sim \langle \gamma_i ; i < \mu \rangle$ if, and only if, $\beta_{\sigma(i)} = \gamma_i$ for each $i < \mu$, for some $\sigma \in G$.

3° If $\mu = \aleph_0$, then $I(\aleph_\alpha ; T) = |\alpha + 1|^{\aleph_0}$, for all $\alpha > 0$ and $I(\aleph_0, T) = 1$, $\aleph_0$ or $2^{\aleph_0}$.

A few final comments ; It can be shown fairly easily that if T is (ω-stable) and multidimensional, then for $\alpha > 0$, $I(\aleph_\alpha, T) \geq 2^{|\alpha|}$. Thus there is some content to the multidmensional/non-multidimensional dichotomy.

SHELAH has classified in a similar manner as above, the $F^a_{\aleph_0}$-saturated models of a superstable non-multidimensional theory.

The main result in this paper, and the main notions employed are due to S. SHELAH, "appearing" in [5]. The bulk of our section I parallels the development of the material in LASCAR [3] (sections 2 and 3). The important proposition III.5 is due to BOUSCAREN and LASCAR [1]. Some results on the spectrum were also obtained by LACHLAN [2].

REFERENCES

[1] BOUSCAREN (E.) and LASCAR (D.). - Countable models of non-multidimensional ω-stable theories (to appear).

[2] LACHLAN (A. H.). - Spectra of ω-stable theories, Z. für math. Logik, t. 24, 1978, p. 129-139.

[3] LASCAR (D.). - Ordre de Rudin-Keisler et poids dans les théories stables (to appear).

[4] LASCAR (D.) and POIZAT (B.). - An introduction to forking, J. of symb. Logic, t. 44, 1979, p. 330-350.

[5] SHELAH (S.). - Classification theory and the number of non-isomorphic models. - Amsterdam, New York, Oxford, North-Holland publishing Company, 1978 (Studies in Logic, 92).